

$\chi_{c1}(3872)$

$I^G(J^{PC}) = 0^+(1^{++})$

also known as $X(3872)$

This state shows properties different from a conventional $q\bar{q}$ state.
A candidate for an exotic structure. See the review on non- $q\bar{q}$ states.

First observed by CHOI 03 in $B \rightarrow K\pi^+\pi^- J/\psi(1S)$ decays as a narrow peak in the invariant mass distribution of the $\pi^+\pi^- J/\psi(1S)$ final state. Isovector hypothesis excluded by AUBERT 05B and CHOI 11.

AAIJ 13Q perform a full five-dimensional amplitude analysis of the angular correlations between the decay products in $B^+ \rightarrow \chi_{c1}(3872)K^+$ decays, where $\chi_{c1}(3872) \rightarrow J/\psi\pi^+\pi^-$ and $J/\psi \rightarrow \mu^+\mu^-$, which unambiguously gives the $J^{PC} = 1^{++}$ assignment under the assumption that the $\pi^+\pi^-$ and J/ψ are in an S -wave. AAIJ 15AO extend this analysis with more data to limit D -wave contributions to < 4% at 95% CL.

See the review on “Spectroscopy of Mesons Containing Two Heavy Quarks.”

$\chi_{c1}(3872)$ MASS FROM $J/\psi X$ MODE

VALUE (MeV)	EVTS			DOCUMENT ID	TECN	COMMENT
3871.65 ± 0.06 OUR AVERAGE						
3871.64 ± 0.06	± 0.01	19.8k		¹ AAIJ	20S LHCb	$B^+ \rightarrow J/\psi\pi^+\pi^- K^+$
3871.9	± 0.7	± 0.2	20	ABLIKIM	14 BES3	$e^+e^- \rightarrow J/\psi\pi^+\pi^-\gamma$
3871.95	± 0.48	± 0.12	0.6k	AAIJ	12H LHCb	$p\bar{p} \rightarrow J/\psi\pi^+\pi^- X$
3871.85	± 0.27	± 0.19	170	² CHOI	11 BELL	$B \rightarrow K\pi^+\pi^- J/\psi$
3873	+ 1.8	± 1.3	27	³ DEL-AMO-SA..10B	BABR	$B \rightarrow \omega J/\psi K$
3871.61	± 0.16	± 0.19	6k	^{3,4} AALTONEN	09AU CDF2	$p\bar{p} \rightarrow J/\psi\pi^+\pi^- X$
3871.4	± 0.6	± 0.1	93.4	AUBERT	08Y BABR	$B^+ \rightarrow K^+ J/\psi\pi^+\pi^-$
3868.7	± 1.5	± 0.4	9.4	AUBERT	08Y BABR	$B^0 \rightarrow K_S^0 J/\psi\pi^+\pi^-$
3871.8	± 3.1	± 3.0	522	^{3,5} ABAZOV	04F D0	$p\bar{p} \rightarrow J/\psi\pi^+\pi^- X$
• • • We do not use the following data for averages, fits, limits, etc. • • •						
3871.695	± 0.067	± 0.068	15.6k	⁶ AAIJ	20AD LHCb	$p\bar{p} \rightarrow J/\psi\pi^+\pi^- X$
3871.59	± 0.06	± 0.03	4.2k	⁷ AAIJ	20S LHCb	$B^+ \rightarrow J/\psi\pi^+\pi^- K^+$
3873.3	± 1.1	± 1.0	45	⁸ ABLIKIM	19V BES	$e^+e^- \rightarrow \gamma\omega J/\psi$
3860.0	± 10.4		13.6	^{3,9} AGHASYAN	18A COMP	$\gamma^* N \rightarrow X\pi^\pm N'$
3868.6	± 1.2	± 0.2	8	¹⁰ AUBERT	06 BABR	$B^0 \rightarrow K_S^0 J/\psi\pi^+\pi^-$

3871.3	± 0.6	± 0.1	61	¹⁰ AUBERT	06	BABR	$B^- \rightarrow K^- J/\psi \pi^+ \pi^-$
3873.4	± 1.4		25	¹¹ AUBERT	05R	BABR	$B^+ \rightarrow K^+ J/\psi \pi^+ \pi^-$
3871.3	± 0.7	± 0.4	730	^{3,12} ACOSTA	04	CDF2	$p\bar{p} \rightarrow J/\psi \pi^+ \pi^- X$
3872.0	± 0.6	± 0.5	36	¹³ CHOI	03	BELL	$B \rightarrow K\pi^+ \pi^- J/\psi$
3836	± 13		58	^{3,14} ANTONIAZZI	94	E705	$\pi^\pm Li \rightarrow J/\psi \pi^+ \pi^- X$

¹ Calculated from $m_{\chi_{c1}(3872)} - m_{\psi(2S)} = 185.54 \pm 0.06$ MeV obtained by combining the data with $\chi_{c1}(3872)$ produced in B^+ decays from AAJ 20S and inclusive b -hadron decays from AAJ 20AD and using $m_{\psi(2S)} = 3686.097$ MeV. Breit-Wigner parametrization.

² The mass difference for the $\chi_{c1}(3872)$ produced in B^+ and B^0 decays is $(-0.71 \pm 0.96 \pm 0.19)$ MeV.

³ Width consistent with detector resolution.

⁴ A possible equal mixture of two states with a mass difference greater than 3.6 MeV/c² is excluded at 95% CL.

⁵ Calculated from the corresponding $m_{\chi_{c1}(3872)} - m_{J/\psi}$ using $m_{J/\psi} = 3096.916$ MeV.

⁶ Using $\chi_{c1}(3872)$ produced in inclusive b -hadron decays and $m_{\psi(2S)} = 3686.097 \pm 0.010$ MeV. Breit-Wigner parametrization. Superseded by the combined value in AAJ 20S.

⁷ Using Breit-Wigner parametrization. Superseded by the combined value in AAJ 20S.

⁸ Fit with fixed width and including two resonances, $\chi_{c0}(3915)$ and $X(3960)$.

⁹ Could be a different state.

¹⁰ Calculated from the corresponding $m_{\chi_{c1}(3872)} - m_{\psi(2S)}$ using $m_{\psi(2S)} = 3686.093$ MeV. Superseded by AUBERT 08Y.

¹¹ Calculated from the corresponding $m_{\chi_{c1}(3872)} - m_{\psi(2S)}$ using $m_{\psi(2S)} = 3685.96$ MeV. Superseded by AUBERT 06.

¹² Superseded by AALTONEN 09AU.

¹³ Superseded by CHOI 11.

¹⁴ A lower mass value can be due to an incorrect momentum scale for soft pions.

$\chi_{c1}(3872)$ MASS FROM $\overline{D}^{*0} D^0$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
$3872.9^{+0.6+0.4}_{-0.4-0.5}$	50	^{1,2} AUSHEV	10	BELL $B \rightarrow \overline{D}^{*0} D^0 K$
$3875.1^{+0.7}_{-0.5} \pm 0.5$	33 ± 6	² AUBERT	08B	BABR $B \rightarrow \overline{D}^{*0} D^0 K$
$3875.2 \pm 0.7^{+0.9}_{-1.8}$	24 ± 6	^{2,3} GOKHROO	06	BELL $B \rightarrow D^0 \overline{D}^0 \pi^0 K$

¹ Calculated from the measured $m_{\chi_{c1}(3872)} - m_{D^{*0}} - m_{\overline{D}^0} = 1.1^{+0.6+0.1}_{-0.4-0.3}$ MeV.

² Experiments report $D^{*0}\overline{D}^0$ invariant mass above $D^{*0}\overline{D}^0$ threshold because D^{*0} decay products are kinematically constrained to the D^{*0} mass, even though the D^{*0} may decay off-shell.

³ Superseded by AUSHEV 10.

$m_{\chi_{c1}(3872)} - m_{J/\psi}$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$774.9 \pm 3.1 \pm 3.0$	522	ABAZOV	04F	$p\bar{p} \rightarrow J/\psi \pi^+ \pi^- X$

$m_{\chi_{c1}(3872)} - m_{\psi(2S)}$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
185.598 $\pm 0.067 \pm 0.068$	15.6k	¹ AAIJ	20AD LHCb	$p p \rightarrow J/\psi \pi^+ \pi^- X$
185.54 ± 0.06	19.8k	² AAIJ	20S LHCb	$p p \rightarrow J/\psi \pi^+ \pi^- X$
187.4 ± 1.4	25	³ AUBERT	05R BABR	$B^+ \rightarrow K^+ J/\psi \pi^+ \pi^-$
¹ Using $\chi_{c1}(3872)$ produced in inclusive b -hadron decays. Breit-Wigner parametrization. Superseded by the combined value in AAIJ 20S. ² Combining $m_{\chi_{c1}(3872)} - m_{\psi(2S)} = 185.49 \pm 0.06 \pm 0.03$ MeV from AAIJ 20S and the measured mass difference from AAIJ 20AD. Breit-Wigner parametrization. ³ Superseded by AUBERT 06.				

$\chi_{c1}(3872)$ WIDTH

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
1.19 ± 0.21 OUR AVERAGE		Error includes scale factor of 1.1.			
1.39 $\pm 0.24 \pm 0.10$	15.6k	¹ AAIJ	20AD LHCb	$p p \rightarrow J/\psi \pi^+ \pi^- X$	
0.96 $\pm 0.19 \pm 0.21$	4.2k	² AAIJ	20S LHCb	$B^+ \rightarrow J/\psi \pi^+ \pi^- K^+$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<2.4	90	ABLIKIM	14	BES3	$e^+ e^- \rightarrow J/\psi \pi^+ \pi^- \gamma$
<1.2	90	CHOI	11	BELL	$B \rightarrow K \pi^+ \pi^- J/\psi$
<3.3	90	AUBERT	08Y	BABR	$B^+ \rightarrow K^+ J/\psi \pi^+ \pi^-$
<4.1	90	69	AUBERT	06	$B \rightarrow K \pi^+ \pi^- J/\psi$
<2.3	90	36	³ CHOI	03	$B \rightarrow K \pi^+ \pi^- J/\psi$
¹ Using $\chi_{c1}(3872)$ produced in inclusive b -hadron decays. Breit-Wigner parametrization. ² Using Breit-Wigner parametrization. Partially overlapping dataset with that of AAIJ 20AD. ³ Superseded by CHOI 11.					

$\chi_{c1}(3872)$ WIDTH FROM $\bar{D}^{*0} D^0$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
3.9 $\pm 2.8 \pm 0.2$	50	¹ AUSHEV	10	BELL $B \rightarrow \bar{D}^{*0} D^0 K$	
3.0 $\pm 1.9 \pm 0.9$	33 ± 6	AUBERT	08B	BABR $B \rightarrow \bar{D}^{*0} D^0 K$	
¹ With a measured value of $B(B \rightarrow \chi_{c1}(3872) K) \times B(\chi_{c1}(3872) \rightarrow D^{*0} \bar{D}^0) = (0.80 \pm 0.20 \pm 0.10) \times 10^{-4}$, assumed to be equal for both charged and neutral modes.					

$\chi_{c1}(3872)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 $e^+ e^-$	$< 2.8 \times 10^{-6}$	90%
Γ_2 $\pi^+ \pi^- \pi^0$	$< 9 \times 10^{-3}$	90%
Γ_3 $\pi^+ \pi^- J/\psi(1S)$	(3.8 ± 1.2) %	
Γ_4 $\pi^+ \pi^- \pi^0 J/\psi(1S)$	not seen	
Γ_5 $\omega \eta_c(1S)$	$< 33 \%$	90%

Γ_6	$\omega J/\psi(1S)$	(4.3 \pm 2.1) %	
Γ_7	$\phi\phi$	not seen	
Γ_8	$D^0 \overline{D}^0 \pi^0$	(49 \pm 18 \pm 20) %	
Γ_9	$\overline{D}^{*0} D^0$	(37 \pm 9) %	
Γ_{10}	$\gamma\gamma$	< 11 %	90%
Γ_{11}	$D^0 \overline{D}^0$	< 29 %	90%
Γ_{12}	$D^+ D^-$	< 19 %	90%
Γ_{13}	$\pi^0 \chi_{c2}$	< 4 %	90%
Γ_{14}	$\pi^0 \chi_{c1}$	(3.4 \pm 1.6) %	
Γ_{15}	$\pi^0 \chi_{c0}$	< 14 %	90%
Γ_{16}	$\pi^+ \pi^- \eta_c(1S)$	< 14 %	90%
Γ_{17}	$\pi^0 \pi^0 \chi_{c0}$	< 7 %	90%
Γ_{18}	$\pi^+ \pi^- \chi_{c0}$	< 2.1 %	90%
Γ_{19}	$\pi^+ \pi^- \chi_{c1}$	< 7 $\times 10^{-3}$	90%
Γ_{20}	$p\bar{p}$	< 2.4 $\times 10^{-5}$	95%

Radiative decays

Γ_{21}	$\gamma D^+ D^-$	< 4 %	90%
Γ_{22}	$\gamma \overline{D}^0 D^0$	< 6 %	90%
Γ_{23}	$\gamma J/\psi$	(8 \pm 4) $\times 10^{-3}$	
Γ_{24}	$\gamma \chi_{c1}$	< 9 $\times 10^{-3}$	90%
Γ_{25}	$\gamma \chi_{c2}$	< 3.2 %	90%
Γ_{26}	$\gamma \psi(2S)$	(4.5 \pm 2.0) %	

C-violating decays

Γ_{27}	$\eta J/\psi$	< 1.8 %	90%
---------------	---------------	---------	-----

$\chi_{c1}(3872)$ PARTIAL WIDTHS

$\Gamma(e^+ e^-)$			Γ_1	
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 4.3	90	¹ ABLIKIM	15V BES3	4.0–4.4 $e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$

<280 90 ² YUAN 04 RVUE $e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$

¹ ABLIKIM 15V reports this limit from the measurement of $\Gamma(\chi_{c1}(3872) \rightarrow \pi^+ \pi^- J/\psi(1S)) \times \Gamma(\chi_{c1}(3872) \rightarrow e^+ e^-)/\Gamma < 0.13$ eV using $\Gamma(\chi_{c1}(3872) \rightarrow \pi^+ \pi^- J/\psi(1S))/\Gamma = 3\%$.

² Using BAI 98E data on $e^+ e^- \rightarrow \pi^+ \pi^- \ell^+ \ell^-$. Assuming that $\Gamma(\pi^+ \pi^- J/\psi)$ of $\chi_{c1}(3872)$ is the same as that of $\psi(2S)$ (85.4 keV).

$\Gamma(\pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$			Γ_2/Γ	
VALUE (%)	CL%	DOCUMENT ID	TECN	COMMENT
<0.9	90	1,2 ADACHI	23 BELL	$B^+ \rightarrow \chi_{c1}(3872) K^+$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.4	90	^{2,3} ADACHI	23	BELL	$B^0 \rightarrow \chi_{c1}(3872) K^0$	
¹ ADACHI 23 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \pi^+ \pi^- \pi^0) / \Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] < 1.9 \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) = 2.1 \times 10^{-4}$.						
² Assuming the decay products, $\pi^+ \pi^- \pi^0$, are uniformly distributed in phase space. The limit is the 90% "credible" upper limit (i.e. Bayesian).						
³ ADACHI 23 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \pi^+ \pi^- \pi^0) / \Gamma_{\text{total}}] \times [B(B^0 \rightarrow \chi_{c1}(3872) K^0)] < 1.5 \times 10^{-6}$ which we divide by our best value $B(B^0 \rightarrow \chi_{c1}(3872) K^0) = 1.1 \times 10^{-4}$.						

$\chi_{c1}(3872) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$

$$\Gamma(\pi^+ \pi^- J/\psi(1S)) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_3 \Gamma_1/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
< 0.13	90	ABLIKIM	15V	$BES3 \quad 4.0-4.4 \text{ e}^+ \text{e}^- \rightarrow \pi^+ \pi^- J/\psi$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
< 6.2	90	^{1,2} AUBERT	05D	$BABR \quad 10.6 \text{ e}^+ \text{e}^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$
< 8.3	90	² DOBBS	05	$CLE3 \quad e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$
< 10	90	³ YUAN	04	$RVUE \quad e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$

¹ Using $B(\chi_{c1}(3872) \rightarrow J/\psi \pi^+ \pi^-) \cdot B(J/\psi \rightarrow \mu^+ \mu^-) \cdot \Gamma(\chi_{c1}(3872) \rightarrow e^+ e^-) < 0.37$ eV from AUBERT 05D and $B(J/\psi \rightarrow \mu^+ \mu^-) = 0.0588 \pm 0.0010$ from the PDG 04.

² Assuming $\chi_{c1}(3872)$ has $J^{PC} = 1^{--}$.

³ Using BAI 98E data on $e^+ e^- \rightarrow \pi^+ \pi^- \ell^+ \ell^-$. From theoretical calculation of the production cross section and using $B(J/\psi \rightarrow \mu^+ \mu^-) = (5.88 \pm 0.10)\%$.

$\chi_{c1}(3872) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$$\Gamma(\pi^+ \pi^- J/\psi(1S)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_3 \Gamma_{10}/\Gamma$$

VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
$5.5^{+4.1}_{-3.8} \pm 0.7$	3	¹ TERAMOTO	21	BELL	$e^+ e^- \rightarrow \gamma^* \gamma$ at $\gamma(nS)$
< 12.9	90	² DOBBS	05	CLE3	$e^+ e^- \rightarrow \pi^+ \pi^- J/\psi \gamma$
$\bullet \bullet \bullet$ Measured in single-tag two-photon production assuming Q^2 dependence of a $c\bar{c}$ meson model. Here, $\Gamma(\chi_{c1}(3872) \rightarrow \gamma\gamma)$ is the reduced two-photon decay width, $\tilde{\Gamma}_{\gamma\gamma}$.					
² Assuming $\chi_{c1}(3872)$ has positive C parity and spin 0.					

$$\Gamma(\omega J/\psi(1S)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_6 \Gamma_{10}/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
< 1.7	90	¹ LEES	12AD BABR	$e^+ e^- \rightarrow e^+ e^- \omega J/\psi$
¹ Assuming $\chi_{c1}(3872)$ has spin 2.				

$$\Gamma(\pi^+ \pi^- \eta_c(1S)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{16} \Gamma_{10}/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
< 11.1	90	LEES	12AE BABR	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \eta_c$

$\chi_{c1}(3872)$ BRANCHING RATIOS

$\Gamma(\pi^+\pi^- J/\psi(1S))/\Gamma_{\text{total}}$	Γ_3/Γ
<u>VALUE</u>	<u>EVTS</u>
0.038±0.012 OUR AVERAGE	
0.038±0.002±0.012	¹ AAIJ
0.041±0.005±0.013	² CHOI
0.040±0.008±0.013	^{3,4} AUBERT
93	
• • • We do not use the following data for averages, fits, limits, etc. • • •	
seen	151
0.061±0.020±0.020	30
0.065±0.014±0.021	36
¹ AAIJ 20S reports $[\Gamma(\chi_{c1}(3872) \rightarrow \pi^+\pi^- J/\psi(1S))/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] = (7.95 \pm 0.15 \pm 0.33) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) = (2.1 \pm 0.7) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.	
² CHOI 11 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \pi^+\pi^- J/\psi(1S))/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] = (8.63 \pm 0.82 \pm 0.52) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) = (2.1 \pm 0.7) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.	
³ AUBERT 08Y reports $[\Gamma(\chi_{c1}(3872) \rightarrow \pi^+\pi^- J/\psi(1S))/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] = (8.4 \pm 1.5 \pm 0.7) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) = (2.1 \pm 0.7) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.	
⁴ superseded by LEES 20C	
⁵ BALA 15 reports $B(\chi_{c1}(3872) \rightarrow \pi^+\pi^- J/\psi) \times B(B^0 \rightarrow \chi_{c1}(3872) K^+\pi^-) = (7.9 \pm 1.3 \pm 0.4) \times 10^{-6}$ and $B(\chi_{c1}(3872) \rightarrow \pi^+\pi^- J/\psi) \times B(B^+ \rightarrow \chi_{c1}(3872) K^0\pi^+) = (10.6 \pm 3.0 \pm 0.9) \times 10^{-6}$.	
⁶ Superseded by AUBERT 08Y. AUBERT 05R reports $[\Gamma(\chi_{c1}(3872) \rightarrow \pi^+\pi^- J/\psi(1S))/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] = (1.28 \pm 0.41) \times 10^{-5}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) = (2.1 \pm 0.7) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.	
⁷ CHOI 03 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \pi^+\pi^- J/\psi(1S))/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] / [B(B^+ \rightarrow \psi(2S) K^+)] / [B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)] = 0.063 \pm 0.012 \pm 0.007$ which we multiply or divide by our best values $B(B^+ \rightarrow \chi_{c1}(3872) K^+) = (2.1 \pm 0.7) \times 10^{-4}$, $B(B^+ \rightarrow \psi(2S) K^+) = (6.24 \pm 0.20) \times 10^{-4}$, $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.68 \pm 0.30) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.	

$\Gamma(\pi^+\pi^-\pi^0 J/\psi(1S))/\Gamma_{\text{total}}$	Γ_4/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u>
not seen	¹ WANG
not seen	² SHEN
¹ WANG 11B reports $B(\gamma(2S) \rightarrow \gamma\chi_{c1}(3872)) \times B(\chi_{c1} \rightarrow \pi^+\pi^-\pi^0 J/\psi) < 2.4 \times 10^{-6}$ at 95% CL.	$\gamma(2S) \rightarrow \gamma X$
² SHEN 10A reports $B(\gamma(1S) \rightarrow \gamma\chi_{c1}(3872)) \times B(\chi_{c1} \rightarrow \pi^+\pi^-\pi^0 J/\psi) < 2.8 \times 10^{-6}$ at 95% CL.	$\gamma(1S) \rightarrow \gamma X$

$\Gamma(\omega\eta_c(1S))/\Gamma_{\text{total}}$					Γ_5/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.33	90	1 VINOKUROVA 15	BELL	$B^+ \rightarrow \omega\eta_c K^+$	
¹ VINOKUROVA 15 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \omega\eta_c(1S))/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] < 6.9 \times 10^{-5}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = 2.1 \times 10^{-4}$.					

$\Gamma(\omega J/\psi(1S))/\Gamma_{\text{total}}$					Γ_6/Γ
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
$0.029 \pm 0.011 \pm 0.009$	21 ± 7	1 DEL-AMO-SA..10B	BABR	$B^+ \rightarrow \omega J/\psi K^+$	
¹ DEL-AMO-SANCHEZ 10B reports $[\Gamma(\chi_{c1}(3872) \rightarrow \omega J/\psi(1S))/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] = (6 \pm 2 \pm 1) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = (2.1 \pm 0.7) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. DEL-AMO-SANCHEZ 10B also reports $B(B^0 \rightarrow \chi_{c1}(3872)K^0) \times B(\chi_{c1}(3872) \rightarrow J/\psi\omega) = (6 \pm 3 \pm 1) \times 10^{-6}$.					

$\Gamma(\omega J/\psi(1S))/\Gamma(\pi^+\pi^- J/\psi(1S))$					Γ_6/Γ_3
VALUE	DOCUMENT ID	TECN	COMMENT		
1.1±0.4 OUR AVERAGE Error includes scale factor of 1.7.					
$1.6^{+0.4}_{-0.3} \pm 0.2$	1 ABLIKIM	19V	BES	$e^+e^- \rightarrow \gamma\omega J/\psi$	
0.8±0.3	2 DEL-AMO-SA..10B	BABR	$B \rightarrow \omega J/\psi K$		
¹ Fit with fixed width and including two resonances, $\chi_{c0}(3915)$ and $X(3960)$.					
² Statistical and systematic errors added in quadrature. Uses the values of $B(B \rightarrow \chi_{c1}(3872)K) \times B(\chi_{c1}(3872) \rightarrow J/\psi\pi^+\pi^-)$ reported in AUBERT 08Y, taking into account the common systematics.					

$\Gamma(\phi\phi)/\Gamma_{\text{total}}$					Γ_7/Γ
VALUE	DOCUMENT ID	TECN	COMMENT		
not seen	1 AAIJ	17BB	LHCb	$p p$ at 7, 8 TeV	
¹ AAIJ 17BB reports $B(b \rightarrow \chi_{c1}(3872)\text{anything}) \times B(\chi_{c1}(3872) \rightarrow \phi\phi) < 4.5 \times 10^{-7}$ at 95% CL.					

$\Gamma(D^0\bar{D}^0\pi^0)/\Gamma_{\text{total}}$					Γ_8/Γ
VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$0.49^{+0.18}_{-0.20} \pm 0.16$	17	1 GOKHROO	06	BELL	$B^+ \rightarrow D^0\bar{D}^0\pi^0 K^+$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
<0.29	90	2 CHISTOV	04	BELL	Sup. by GOKHROO 06
¹ GOKHROO 06 reports $[\Gamma(\chi_{c1}(3872) \rightarrow D^0\bar{D}^0\pi^0)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] = (1.02 \pm 0.31^{+0.21}_{-0.29}) \times 10^{-4}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = (2.1 \pm 0.7) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					
² CHISTOV 04 reports $[\Gamma(\chi_{c1}(3872) \rightarrow D^0\bar{D}^0\pi^0)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] < 0.6 \times 10^{-4}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = 2.1 \times 10^{-4}$.					

$\Gamma(D^0\bar{D}^0\pi^0)/\Gamma(\pi^+\pi^- J/\psi(1S))$ Γ_8/Γ_3

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
<1.16	90	ABLIKIM	20W BES3	$e^+e^- \rightarrow \gamma\chi_{c1}(3872)$

 $\Gamma(\bar{D}^{*0}D^0)/\Gamma_{\text{total}}$ Γ_9/Γ

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.37±0.09±0.12	41 ⁺⁹ ₋₈	1 AUSHEV	10 BELL	$B^+ \rightarrow D^{*0}\bar{D}^0 K^+$

 $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ 0.80±0.28±0.26 27 ± 6 ² AUBERT 08B BABR $B^+ \rightarrow \bar{D}^{*0}D^0 K^+$

¹ AUSHEV 10 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \bar{D}^{*0}D^0)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] = (0.77 \pm 0.16 \pm 0.10) \times 10^{-4}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = (2.1 \pm 0.7) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² AUBERT 08B reports $[\Gamma(\chi_{c1}(3872) \rightarrow \bar{D}^{*0}D^0)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] = (1.67 \pm 0.36 \pm 0.47) \times 10^{-4}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = (2.1 \pm 0.7) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\bar{D}^{*0}D^0)/\Gamma(\pi^+\pi^- J/\psi(1S))$ Γ_9/Γ_3

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
11.77±3.09	50	ABLIKIM	20W BES3	$e^+e^- \rightarrow \gamma\chi_{c1}(3872)$

 $\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ Γ_{10}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.11	90	1 WICHT	08 BELL	$e^+e^- \rightarrow \gamma(4S)$

¹ WICHT 08 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \gamma\gamma)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] < 2.4 \times 10^{-5}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = 2.1 \times 10^{-4}$.

 $\Gamma(D^0\bar{D}^0)/\Gamma_{\text{total}}$ Γ_{11}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.29	90	1 CHISTOV	04 BELL	$B \rightarrow KD^0\bar{D}^0$

¹ CHISTOV 04 reports $[\Gamma(\chi_{c1}(3872) \rightarrow D^0\bar{D}^0)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] < 6 \times 10^{-5}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = 2.1 \times 10^{-4}$.

 $\Gamma(D^+D^-)/\Gamma_{\text{total}}$ Γ_{12}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.19	90	1 CHISTOV	04 BELL	$B \rightarrow KD^+D^-$

¹ CHISTOV 04 reports $[\Gamma(\chi_{c1}(3872) \rightarrow D^+D^-)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] < 4 \times 10^{-5}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = 2.1 \times 10^{-4}$.

 $\Gamma(\pi^0\chi_{c2})/\Gamma(\pi^+\pi^- J/\psi(1S))$ Γ_{13}/Γ_3

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.1	90	ABLIKIM	19U BES3	$e^+e^- \rightarrow \gamma\chi_{c1}(3872)$

$\Gamma(\pi^0 \chi_{c1})/\Gamma_{\text{total}}$ Γ_{14}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.04	90	¹ BHARDWAJ	19 BELL	$B^\pm \rightarrow \pi^0 \chi_{c1} K^\pm$
1 BHARDWAJ	19	reports [$\Gamma(\chi_{c1}(3872) \rightarrow \pi^0 \chi_{c1})/\Gamma_{\text{total}}$] $\times [B(B^+ \rightarrow \chi_{c1}(3872) K^+) < 8.1 \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) = 2.1 \times 10^{-4}$.		

 $\Gamma(\pi^0 \chi_{c1})/\Gamma(\pi^+ \pi^- J/\psi(1S))$ Γ_{14}/Γ_3

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$88^{+33}_{-27} \pm 10$	10.8	ABLIKIM	19U BES3	$e^+ e^- \rightarrow \gamma \chi_{c1}(3872)$

 $\Gamma(\pi^0 \pi^0 \chi_{c0})/\Gamma(\pi^+ \pi^- J/\psi(1S))$ Γ_{17}/Γ_3

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.7	90	ABLIKIM	22D BES3	$e^+ e^- \rightarrow \gamma \chi_{c1}(3872)$

 $\Gamma(\pi^+ \pi^- \chi_{c0})/\Gamma(\pi^+ \pi^- J/\psi(1S))$ Γ_{18}/Γ_3

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.56	90	ABLIKIM	22D BES3	$e^+ e^- \rightarrow \gamma \chi_{c1}(3872)$

 $\Gamma(\pi^0 \chi_{c0})/\Gamma(\pi^+ \pi^- J/\psi(1S))$ Γ_{15}/Γ_3

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 3.6	90	ABLIKIM	22D BES3	$e^+ e^- \rightarrow \gamma \chi_{c1}(3872)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<19	90	ABLIKIM	19U BES3	$e^+ e^- \rightarrow \gamma \chi_{c1}(3872)$

 $\Gamma(\pi^+ \pi^- \eta_c(1S))/\Gamma_{\text{total}}$ Γ_{16}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.14	90	¹ VINOKUROVA	15 BELL	$B^+ \rightarrow \pi^+ \pi^- \eta_c K^+$
1 VINOKUROVA 15 reports [$\Gamma(\chi_{c1}(3872) \rightarrow \pi^+ \pi^- \eta_c(1S))/\Gamma_{\text{total}}$] $\times [B(B^+ \rightarrow \chi_{c1}(3872) K^+) < 3.0 \times 10^{-5}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) = 2.1 \times 10^{-4}$.				

 $\Gamma(\pi^+ \pi^- \chi_{c1})/\Gamma_{\text{total}}$ Γ_{19}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<7 \times 10^{-3}$	90	¹ BHARDWAJ	16 BELL	$B^+ \rightarrow \pi^+ \pi^- \chi_{c1} K^+$
1 BHARDWAJ 16 reports [$\Gamma(\chi_{c1}(3872) \rightarrow \pi^+ \pi^- \chi_{c1})/\Gamma_{\text{total}}$] $\times [B(B^+ \rightarrow \chi_{c1}(3872) K^+) < 1.5 \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) = 2.1 \times 10^{-4}$.				

 $\Gamma(p\bar{p})/\Gamma_{\text{total}}$ Γ_{20}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<2.4 \times 10^{-5}$	95	¹ AAIJ	17AD LHCb	$B^+ \rightarrow p\bar{p} K^+$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<8 \times 10^{-5}$	95	² AAIJ	13S LHCb	$B^+ \rightarrow p\bar{p}K^+$	
¹ AAIJ 17AD reports $[\Gamma(\chi_{c1}(3872) \rightarrow p\bar{p})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] < 0.5 \times 10^{-8}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = 2.1 \times 10^{-4}$.					
² AAIJ 13S reports $[\Gamma(\chi_{c1}(3872) \rightarrow p\bar{p})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] < 1.7 \times 10^{-8}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = 2.1 \times 10^{-4}$.					

———— Radiative decays ——

$\Gamma(\gamma D^+ D^-)/\Gamma(\pi^+ \pi^- J/\psi(1S))$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{21}/Γ_3
<0.99	90	ABLIKIM	20W	$e^+ e^- \rightarrow \gamma \chi_{c1}(3872)$	

$\Gamma(\gamma \bar{D}^0 D^0)/\Gamma(\pi^+ \pi^- J/\psi(1S))$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{22}/Γ_3
<1.58	90	ABLIKIM	20W	$e^+ e^- \rightarrow \gamma \chi_{c1}(3872)$	

$\Gamma(\gamma J/\psi)/\Gamma_{\text{total}}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{23}/Γ
$0.0085^{+0.0024}_{-0.0022} \pm 0.0027$		¹ BHARDWAJ	11	BELL $B^\pm \rightarrow \gamma J/\psi K^\pm$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.013 $\pm 0.004 \pm 0.004$	20	² AUBERT	09B BABR	$B^+ \rightarrow \gamma J/\psi K^+$
0.016 $\pm 0.005 \pm 0.005$	19	³ AUBERT,BE	06M BABR	$B^+ \rightarrow \gamma J/\psi K^+$

¹ BHARDWAJ 11 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \gamma J/\psi)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] = (1.78^{+0.48}_{-0.44} \pm 0.12) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = (2.1 \pm 0.7) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² AUBERT 09B reports $[\Gamma(\chi_{c1}(3872) \rightarrow \gamma J/\psi)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] = (2.8 \pm 0.8 \pm 0.1) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = (2.1 \pm 0.7) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ Superseded by AUBERT 09B. AUBERT,BE 06M reports $[\Gamma(\chi_{c1}(3872) \rightarrow \gamma J/\psi)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] = (3.3 \pm 1.0 \pm 0.3) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = (2.1 \pm 0.7) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\gamma J/\psi)/\Gamma(\pi^+ \pi^- J/\psi(1S))$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{23}/Γ_3
0.79 ± 0.28	ABLIKIM	20W	$e^+ e^- \rightarrow \gamma \chi_{c1}(3872)$	

$\Gamma(\gamma \chi_{c1})/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{24}/Γ
$<9 \times 10^{-3}$	90	¹ BHARDWAJ	13	BELL $B^\pm \rightarrow \chi_{c1}\gamma K^\pm$	

¹ BHARDWAJ 13 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \gamma \chi_{c1})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] < 1.9 \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = 2.1 \times 10^{-4}$.

$\Gamma(\gamma\chi_{c1})/\Gamma(\pi^+\pi^- J/\psi(1S))$

Γ_{24}/Γ_3

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.89	90	CHOI	03	BELL $B \rightarrow K\pi^+\pi^- J/\psi$

$\Gamma(\gamma\chi_{c2})/\Gamma_{\text{total}}$

Γ_{25}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.032	90	¹ BHARDWAJ	13	BELL $B^\pm \rightarrow \chi_{c2}\gamma K^\pm$

¹ BHARDWAJ 13 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \gamma\chi_{c2})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] < 6.7 \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = 2.1 \times 10^{-4}$.

$\Gamma(\gamma\psi(2S))/\Gamma_{\text{total}}$

Γ_{26}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.045±0.013±0.015	25 ± 7	¹ AUBERT	09B BABR	$B^+ \rightarrow \gamma\psi(2S)K^+$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
seen	36 ± 9	² AAIJ	14AH LHCb	$B^+ \rightarrow \gamma\psi(2S)K^+$
not seen		³ BHARDWAJ	11 BELL	$B^+ \rightarrow \gamma\psi(2S)K^+$

¹ AUBERT 09B reports $[\Gamma(\chi_{c1}(3872) \rightarrow \gamma\psi(2S))/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] = (9.5 \pm 2.7 \pm 0.6) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = (2.1 \pm 0.7) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² From 36.4 ± 9.0 events of $\chi_{c1}(3872) \rightarrow J/\psi\gamma$ decays with a statistical significance of 4.4σ .

³ BHARDWAJ 11 reports $B(B^+ \rightarrow K^+\chi_{c1}(3872)) \times B(\chi_{c1} \rightarrow \gamma\psi(2S)) < 3.45 \times 10^{-6}$ at 90% CL.

$\Gamma(\gamma\psi(2S))/\Gamma(\pi^+\pi^- J/\psi(1S))$

Γ_{26}/Γ_3

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.42	90	ABLIKIM	20W BES3	$e^+e^- \rightarrow \gamma\chi_{c1}(3872)$

$\Gamma(\gamma\psi(2S))/\Gamma(\gamma J/\psi)$

Γ_{26}/Γ_{23}

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
2.6 ±0.6 OUR AVERAGE					

$2.46 \pm 0.64 \pm 0.29$ 36 ± 9 ¹ AAIJ 14AH LHCb $B^+ \rightarrow \gamma\psi(2S)K^+$
 3.4 ± 1.4 AUBERT 09B BABR $B^+ \rightarrow \gamma c\bar{c}K'$

• • • We do not use the following data for averages, fits, limits, etc. **• • •**

<2.1 90 BHARDWAJ 11 BELL $B^+ \rightarrow \gamma\psi(2S)K^+$

¹ From 36.4 ± 9.0 events of $\chi_{c1}(3872) \rightarrow J/\psi\gamma$ decays with a statistical significance of 4.4σ .

— C-violating decays —

$\Gamma(\eta J/\psi)/\Gamma_{\text{total}}$

Γ_{27}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.018	90	^{1,2} IWASHITA	14 BELL	$B \rightarrow K\eta J/\psi$

• • • We do not use the following data for averages, fits, limits, etc. **• • •**

<0.04 90 ³ AUBERT 04Y BABR $B \rightarrow K\eta J/\psi$

¹ IWASHITA 14 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \eta J/\psi)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] < 3.8 \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = 2.1 \times 10^{-4}$.

² IWASHITA 14 also scans the $\eta J/\psi$ mass range 3.8–4.75 GeV and sets upper limits for $B(B^\pm \rightarrow \chi_{c1}(3872) K^\pm) \times B(\chi_{c1}(3872) \rightarrow \eta J/\psi)$ in 5 MeV intervals.

³ AUBERT 04Y reports $[\Gamma(\chi_{c1}(3872) \rightarrow \eta J/\psi) / \Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] < 7.7 \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) = 2.1 \times 10^{-4}$.

$\chi_{c1}(3872)$ REFERENCES

ADACHI	23	PR D107 052004	I. Adachi <i>et al.</i>	(BELLE Collab.)
ABLIKIM	22D	PR D105 072009	M. Ablikim <i>et al.</i>	(BESIII Collab.)
TERAMOTO	21	PRL 126 122001	Y. Teramoto <i>et al.</i>	(BELLE Collab.)
AAIJ	20AD	PR D102 092005	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	20S	JHEP 2008 123	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	20W	PRL 124 242001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
LEES	20C	PRL 124 152001	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	19U	PRL 122 202001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19V	PRL 122 232002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
BHARDWAJ	19	PR D99 111101	V. Bhardwaj <i>et al.</i>	(BELLE Collab.)
AGHASYAN	18A	PL B783 334	M. Aghasyan <i>et al.</i>	(COMPASS Collab.)
AAIJ	17AD	PL B769 305	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	17BB	EPJ C77 609	R. Aaij <i>et al.</i>	(LHCb Collab.)
BHARDWAJ	16	PR D93 052016	V. Bhardwaj <i>et al.</i>	(BELLE Collab.)
AAIJ	15AO	PR D92 011102	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	15V	PL B749 414	M. Ablikim <i>et al.</i>	(BESIII Collab.)
BALA	15	PR D91 051101	A. Bala <i>et al.</i>	(BELLE Collab.)
VINOKUROVA	15	JHEP 1506 132	A. Vinokurova <i>et al.</i>	(BELLE Collab.)
Also		JHEP 1702 088 (errat.)	A. Vinokurava <i>et al.</i>	(BELLE Collab.)
AAIJ	14AH	NP B886 665	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	14	PRL 112 092001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
IWASHITA	14	PTEP 2014 043C01	T. Iwashita <i>et al.</i>	(BELLE Collab.)
AAIJ	13Q	PRL 110 222001	R. Aaij <i>et al.</i>	(LHCb Collab.) JP
AAIJ	13S	EPJ C73 2462	R. Aaij <i>et al.</i>	(LHCb Collab.)
BHARDWAJ	13	PRL 111 032001	V. Bhardwaj <i>et al.</i>	(BELLE Collab.)
AAIJ	12H	EPJ C72 1972	R. Aaij <i>et al.</i>	(LHCb Collab.)
LEES	12AD	PR D86 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	12AE	PR D86 092005	J.P. Lees <i>et al.</i>	(BABAR Collab.)
BHARDWAJ	11	PRL 107 091803	V. Bhardwaj <i>et al.</i>	(BELLE Collab.)
CHOI	11	PR D84 052004	S.-K. Choi <i>et al.</i>	(BELLE Collab.)
WANG	11B	PR D84 071107	X.L. Wang <i>et al.</i>	(BELLE Collab.)
AUSHEV	10	PR D81 031103	T. Aushev <i>et al.</i>	(BELLE Collab.)
DEL-AMO-SA...	10B	PR D82 011101	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
SHEN	10A	PR D82 051504	C.P. Shen <i>et al.</i>	(BELLE Collab.)
AALTONEN	09AU	PRL 103 152001	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AUBERT	09B	PRL 102 132001	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08B	PR D77 011102	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08Y	PR D77 111101	B. Aubert <i>et al.</i>	(BABAR Collab.)
WICHT	08	PL B662 323	J. Wicht <i>et al.</i>	(BELLE Collab.)
AUBERT	06	PR D73 011101	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,BE	06M	PR D74 071101	B. Aubert <i>et al.</i>	(BABAR Collab.)
GOKHROO	06	PRL 97 162002	G. Gokhroo <i>et al.</i>	(BELLE Collab.)
AUBERT	05B	PR D71 031501	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	05D	PR D71 052001	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	05R	PR D71 071103	B. Aubert <i>et al.</i>	(BABAR Collab.)
DOBBS	05	PRL 94 032004	S. Dobbs <i>et al.</i>	(CLEO Collab.)
ABAZOV	04F	PRL 93 162002	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ACOSTA	04	PRL 93 072001	D. Acosta <i>et al.</i>	(CDF Collab.)
AUBERT	04Y	PRL 93 041801	B. Aubert <i>et al.</i>	(BABAR Collab.)
CHISTOV	04	PRL 93 051803	R. Chistov <i>et al.</i>	(BELLE Collab.)
PDG	04	PL B592 1	S. Eidelman <i>et al.</i>	(PDG Collab.)
YUAN	04	PL B579 74	C.Z. Yuan <i>et al.</i>	(BELLE Collab.)
CHOI	03	PRL 91 262001	S.-K. Choi <i>et al.</i>	(BES Collab.)
BAI	98E	PR D57 3854	J.Z. Bai <i>et al.</i>	(E705 Collab.)
ANTONIAZZI	94	PR D50 4258	L. Antoniazzi <i>et al.</i>	