

$\psi(4415)$

$I^G(J^{PC}) = 0^-(1^{--})$

$\psi(4415)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
4421 ± 4 OUR ESTIMATE			
4415.1± 7.9	¹ ABLIKIM	08D BES2	$e^+e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •			
4412 ± 15	² MO	10 RVUE	$e^+e^- \rightarrow$ hadrons
4411 ± 7	³ PAKHLOVA	08A BELL	$10.6 e^+e^- \rightarrow D^0 D^- \pi^+\gamma$
4425 ± 6	⁴ SETH	05A RVUE	$e^+e^- \rightarrow$ hadrons
4429 ± 9	⁵ SETH	05A RVUE	$e^+e^- \rightarrow$ hadrons
4417 ± 10	BRANDELIK	78C DASP	e^+e^-
4414 ± 7	SIEGRIST	76 MRK1	e^+e^-

¹ Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = (234 \pm 88)^\circ$.

² Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the $\psi(4040)$, $\psi(4160)$ and $\psi(4415)$ resonances and including interference effects.

³ Systematic uncertainties not estimated.

⁴ From a fit to Crystal Ball (OSTERHELD 86) data.

⁵ From a fit to BES (BAI 02C) data.

$\psi(4415)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
62 ±20 OUR ESTIMATE			
71.5±19.0	⁶ ABLIKIM	08D BES2	$e^+e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •			
118 ± 32	⁷ MO	10 RVUE	$e^+e^- \rightarrow$ hadrons
77 ± 20	⁸ PAKHLOVA	08A BELL	$10.6 e^+e^- \rightarrow D^0 D^- \pi^+\gamma$
119 ± 16	⁹ SETH	05A RVUE	$e^+e^- \rightarrow$ hadrons
118 ± 35	¹⁰ SETH	05A RVUE	$e^+e^- \rightarrow$ hadrons
66 ± 15	BRANDELIK	78C DASP	e^+e^-
33 ± 10	SIEGRIST	76 MRK1	e^+e^-

⁶ Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = (234 \pm 88)^\circ$.

⁷ Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the $\psi(4040)$, $\psi(4160)$ and $\psi(4415)$ resonances and including interference effects.

⁸ Systematic uncertainties not estimated.

⁹ From a fit to Crystal Ball (OSTERHELD 86) data.

¹⁰ From a fit to BES (BAI 02C) data.

$\psi(4415)$ DECAY MODES

Due to the complexity of the $c\bar{c}$ threshold region, in this listing, “seen” (“not seen”) means that a cross section for the mode in question has been measured at effective \sqrt{s} near this particle’s central mass value, more (less) than 2σ above zero, without regard to any peaking behavior in \sqrt{s} or absence thereof. See mode listing(s) for details and references.

Mode	Fraction (Γ_i/Γ)	Confidence level	
$\Gamma_1 D\overline{D}$		seen	
$\Gamma_2 D^0\overline{D}^0$		seen	
$\Gamma_3 D^+D^-$		seen	
$\Gamma_4 D^*\overline{D} + \text{c.c.}$		seen	
$\Gamma_5 D^*(2007)^0\overline{D}^0 + \text{c.c.}$		seen	
$\Gamma_6 D^*(2010)^+D^- + \text{c.c.}$		seen	
$\Gamma_7 D^*\overline{D}^*$		seen	
$\Gamma_8 D^*(2007)^0\overline{D}^*(2007)^0 + \text{c.c.}$		seen	
$\Gamma_9 D^*(2010)^+D^*(2010)^- + \text{c.c.}$		seen	
$\Gamma_{10} D^0 D^- \pi^+ (\text{excl. } D^*(2007)^0\overline{D}^0 + \text{c.c.}, D^*(2010)^+D^- + \text{c.c.})$	< 2.3 %		90%
$\Gamma_{11} D\overline{D}_2^*(2460) \rightarrow D^0 D^- \pi^+ + \text{c.c.}$	(10 ± 4) %		
$\Gamma_{12} D^0 D^{*-} \pi^+ + \text{c.c.}$	< 11 %		90%
$\Gamma_{13} D_1(2420)\overline{D} + \text{c.c.}$		possibly seen	
$\Gamma_{14} D_s^+ D_s^-$		not seen	
$\Gamma_{15} \omega \chi_{c2}$		possibly seen	
$\Gamma_{16} D_s^{*+} D_s^- + \text{c.c.}$		seen	
$\Gamma_{17} D_s^{*+} D_s^{*-}$		not seen	
$\Gamma_{18} \psi_2(3823)\pi^+\pi^-$		possibly seen	
$\Gamma_{19} \psi(3770)\pi^+\pi^-$		possibly seen	
$\Gamma_{20} J/\psi\eta$	< 6 × 10 ⁻³		90%
$\Gamma_{21} \chi_{c1}\gamma$	< 8 × 10 ⁻⁴		90%
$\Gamma_{22} \chi_{c2}\gamma$	< 4 × 10 ⁻³		90%
$\Gamma_{23} \Lambda\overline{\Lambda}$	< 3.1 × 10 ⁻⁶		90%
$\Gamma_{24} \omega\pi^0$		not seen	
$\Gamma_{25} \omega\eta$		not seen	
$\Gamma_{26} e^+e^-$	(9.4 ± 3.2) × 10 ⁻⁶		
$\Gamma_{27} \mu^+\mu^-$	(2.0 ± 1.0) × 10 ⁻⁵		

$\psi(4415)$ PARTIAL WIDTHS

$\Gamma(e^+e^-)$	Γ_{26}
0.58 ± 0.07 OUR ESTIMATE	
0.35 ± 0.12	11 ABLIKIM 08D BES2 $e^+e^- \rightarrow \text{hadrons}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.4 to 0.8	¹² MO	10	RVUE	$e^+ e^- \rightarrow$	hadrons
0.72 ± 0.11	¹³ SETH	05A	RVUE	$e^+ e^- \rightarrow$	hadrons
0.64 ± 0.23	¹⁴ SETH	05A	RVUE	$e^+ e^- \rightarrow$	hadrons
0.49 ± 0.13	BRANDELIK	78C	DASP	$e^+ e^-$	
0.44 ± 0.14	SIEGRIST	76	MRK1	$e^+ e^-$	

¹¹ Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = (234 \pm 88)^\circ$.

¹² Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the $\psi(4040)$, $\psi(4160)$ and $\psi(4415)$ resonances and including interference effects. Four sets of solutions are obtained with the same fit quality, mass and total width, but with different $e^+ e^-$ partial widths. We quote only the range of values.

¹³ From a fit to Crystal Ball (OSTERHELD 86) data.

¹⁴ From a fit to BES (BAI 02c) data.

$\Gamma(\mu^+ \mu^-)$	Γ_{27}		
VALUE (keV)	DOCUMENT ID	TECN	COMMENT
$1.25 \pm 0.28 \pm 0.35$	^{15,16} ABLIKIM	20AG BES3	$e^+ e^- \rightarrow \mu^+ \mu^-$
¹⁵ From a fit to the $e^+ e^- \rightarrow \mu^+ \mu^-$ cross section between 3.8 and 4.6 GeV to the coherent sum of four resonant amplitudes assuming $\Gamma(\mu^+ \mu^-) = \Gamma(e^+ e^-)$.			
¹⁶ From solution 1 of 8 with equal fit quality. Other solutions range from $1.24 \pm 0.28 \pm 0.35$ to $1.27 \pm 0.41 \pm 0.36$ keV.			

$\psi(4415) \Gamma(i) \times \Gamma(e^+ e^-)/\Gamma(\text{total})$

$\Gamma(J/\psi \eta) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_{20}\Gamma_{26}/\Gamma$			
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<3.6	90	WANG	13B	$e^+ e^- \rightarrow J/\psi \eta \gamma$

$\Gamma(\chi_{c1} \gamma) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_{21}\Gamma_{26}/\Gamma$			
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<0.47	90	¹⁷ HAN	15	$10.58 e^+ e^- \rightarrow \chi_{c1} \gamma$

¹⁷ Using $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$.

$\Gamma(\chi_{c2} \gamma) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_{22}\Gamma_{26}/\Gamma$			
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<2.3	90	¹⁸ HAN	15	$10.58 e^+ e^- \rightarrow \chi_{c2} \gamma$

¹⁸ Using $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$.

$\Gamma(\Lambda\bar{\Lambda}) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_{23}\Gamma_{26}/\Gamma$			
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
$<1.8 \times 10^{-3}$	90	¹⁹ ABLIKIM	21AS BES3	$e^+ e^- \rightarrow \psi(4415)$

¹⁹ From a measurement of the $e^+ e^- \rightarrow \Lambda\bar{\Lambda}$ cross section between 3.5 and 4.6 GeV.

$\psi(4415)$ BRANCHING RATIOS

$\Gamma(D^0\bar{D}^0)/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_2/Γ
seen	PAKHLOVA 08	BELL	$e^+ e^- \rightarrow D^0\bar{D}^0\gamma$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
not seen	AUBERT 09M BABR	$e^+ e^- \rightarrow D^0\bar{D}^0\gamma$		

$\Gamma(D^+\bar{D}^-)/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_3/Γ
seen	PAKHLOVA 08	BELL	$e^+ e^- \rightarrow D^+\bar{D}^-\gamma$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
not seen	AUBERT 09M BABR	$e^+ e^- \rightarrow D^+\bar{D}^-\gamma$		

$\Gamma(D\bar{D})/\Gamma(D^*\bar{D}^*)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_1/Γ_7
0.14±0.12±0.03	AUBERT 09M BABR	$e^+ e^- \rightarrow \gamma D^{(*)}\bar{D}^{(*)}$		

$\Gamma(D^*(2007)^0\bar{D}^0 + \text{c.c.})/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_5/Γ
seen	AUBERT 09M BABR	$e^+ e^- \rightarrow D^{*0}\bar{D}^0\gamma$		

$\Gamma(D^*(2010)^+ D^- + \text{c.c.})/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_6/Γ
seen	20 ZHUKOVA 18	BELL	$e^+ e^- \rightarrow D^{*+}D^-\gamma$	
seen	AUBERT 09M BABR	$e^+ e^- \rightarrow D^{*+}D^-\gamma$		
• • • We do not use the following data for averages, fits, limits, etc. • • •				
seen	PAKHLOVA 07	BELL	$e^+ e^- \rightarrow D^{*+}D^-\gamma$	

20 Supersedes PAKHLOVA 07.

$\Gamma(D^*\bar{D} + \text{c.c.})/\Gamma(D^*\bar{D}^*)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_4/Γ_7
0.17±0.25±0.03	AUBERT 09M BABR	$e^+ e^- \rightarrow \gamma D^{(*)}\bar{D}^{(*)}$		

$\Gamma(D^*(2007)^0\bar{D}^*(2007)^0 + \text{c.c.})/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_8/Γ
seen	AUBERT 09M BABR	$e^+ e^- \rightarrow D^{*0}\bar{D}^{*0}\gamma$		

$\Gamma(D^*(2010)^+ D^*(2010)^- + \text{c.c.})/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_9/Γ
seen	21 ZHUKOVA 18	BELL	$e^+ e^- \rightarrow D^{*+}D^{*-}\gamma$	
seen	AUBERT 09M BABR	$e^+ e^- \rightarrow D^{*+}D^{*-}\gamma$		

• • • We do not use the following data for averages, fits, limits, etc. • • •

<u>seen</u>	<u>PAKHLOVA 07</u>	<u>BELL</u>	$e^+ e^- \rightarrow D^{*+}D^{*-}\gamma$
21	Supersedes PAKHLOVA 07.		

$\Gamma(D\bar{D}_2^*(2460) \rightarrow D^0 D^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$

Γ_{11}/Γ

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
$10.5 \pm 2.4 \pm 3.8$	22 PAKHLOVA 08A BELL		$10.6 e^+ e^- \rightarrow D^0 D^- \pi^+ \gamma$

22 Using 4421 ± 4 MeV for the mass and 62 ± 20 MeV for the width of $\psi(4415)$.

$\Gamma(D^0 D^- \pi^+ (\text{excl. } D^*(2007)^0 \bar{D}^0 + \text{c.c.}, D^*(2010)^+ D^- + \text{c.c.}) /$

$\Gamma(D\bar{D}_2^*(2460) \rightarrow D^0 D^- \pi^+ + \text{c.c.})$

Γ_{10}/Γ_{11}

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.22	90	23 PAKHLOVA 08A BELL		$10.6 e^+ e^- \rightarrow D^0 D^- \pi^+ \gamma$

23 Using 4421 ± 4 MeV for the mass and 62 ± 20 MeV for the width of $\psi(4415)$.

$\Gamma(D^0 D^{*-} \pi^+ + \text{c.c.})/\Gamma_{\text{total}} \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$

$\Gamma_{12}/\Gamma \times \Gamma_{26}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<0.99 \times 10^{-6}$	90	24 PAKHLOVA 09 BELL		$e^+ e^- \rightarrow D^0 D^{*-} \pi^+$

24 Using 4421 ± 4 MeV for the mass of $\psi(4415)$.

$\Gamma(D_1(2420)\bar{D} + \text{c.c.})/\Gamma_{\text{total}}$

Γ_{13}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
possibly seen	25 ABLIKIM 19AR BES3		$e^+ e^- \rightarrow \pi^+ \pi^- D\bar{D}$

25 Evidence for $e^+ e^- \rightarrow D_1(2420)\bar{D} + \text{c.c.}$ between $\sqrt{s} = 4.3$ and 4.6 GeV, not necessarily resonant.

$\Gamma(D_s^+ D_s^-)/\Gamma_{\text{total}}$

Γ_{14}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	PAKHLOVA 11 BELL		$e^+ e^- \rightarrow D_s^+ D_s^- \gamma$
not seen	DEL-AMO-SA..10N BABR		$e^+ e^- \rightarrow D_s^+ D_s^- \gamma$

$\Gamma(\omega \chi_{c2})/\Gamma_{\text{total}}$

Γ_{15}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
possibly seen	ABLIKIM 16A BES3		$e^+ e^- \rightarrow \gamma \pi^+ \pi^- \pi^0 \ell^+ \ell^-$

$\Gamma(D_s^{*+} D_s^- + \text{c.c.})/\Gamma_{\text{total}}$

Γ_{16}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	PAKHLOVA 11 BELL		$e^+ e^- \rightarrow D_s^{*+} D_s^- \gamma$
seen	DEL-AMO-SA..10N BABR		$e^+ e^- \rightarrow D_s^{*+} D_s^- \gamma$

$\Gamma(D_s^{*+} D_s^{*-})/\Gamma_{\text{total}}$

Γ_{17}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	PAKHLOVA 11 BELL		$e^+ e^- \rightarrow D_s^{*+} D_s^{*-} \gamma$
not seen	DEL-AMO-SA..10N BABR		$e^+ e^- \rightarrow D_s^{*+} D_s^{*-} \gamma$

$\Gamma(\psi(3770)\pi^+ \pi^-)/\Gamma_{\text{total}}$

Γ_{19}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
possibly seen	26 ABLIKIM 19AR BES3		$e^+ e^- \rightarrow \pi^+ \pi^- D\bar{D}$

26 Observe $e^+ e^- \rightarrow \pi^+ \pi^- \psi(3770)$ at $\sqrt{s} = 4.26, 4.36$, and 4.42 GeV but cannot establish if continuum or resonant.

$\Gamma(\psi_2(3823)\pi^+\pi^-)/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{18}/Γ
possibly seen	19	27	ABLIKIM	15S BES3	$e^+e^- \rightarrow \pi^+\pi^-\chi_{c1}\gamma$
27 From a fit of $e^+e^- \rightarrow \pi^+\pi^-\psi_2(3823)$, $\psi_2(3823) \rightarrow \chi_{c1}\gamma$ cross sections taken at \sqrt{s} values of 4.23, 4.26, 4.36, 4.42, and 4.60 GeV to the $\psi(4415)$ line shape.					

 $\Gamma(\omega\pi^0)/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{24}/Γ
not seen	ABLIKIM	22K	BES3	$e^+e^- \rightarrow \omega\pi^0$

 $\Gamma(\omega\eta)/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{25}/Γ
not seen	ABLIKIM	22K	BES3	$e^+e^- \rightarrow \omega\eta$

 $\psi(4415)$ REFERENCES

ABLIKIM	22K	JHEP 2207 064	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21AS	PR D104 L091104	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	20AG	PR D102 112009	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19AR	PR D100 032005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ZHUKOVA	18	PR D97 012002	V. Zhukova <i>et al.</i>	(BELLE Collab.)
ABLIKIM	16A	PR D93 011102	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	15S	PRL 115 011803	M. Ablikim <i>et al.</i>	(BESIII Collab.)
HAN	15	PR D92 012011	Y.L. Han <i>et al.</i>	(BELLE Collab.)
WANG	13B	PR D87 051101	X.L. Wang <i>et al.</i>	(BELLE Collab.)
PAKHLOVA	11	PR D83 011101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
DEL-AMO-SA...	10N	PR D82 052004	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
MO	10	PR D82 077501	X.H. Mo, C.Z. Yuan, P. Wang	(BHEP)
AUBERT	09M	PR D79 092001	B. Aubert <i>et al.</i>	(BABAR Collab.)
PAKHLOVA	09	PR D80 091101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
ABLIKIM	08D	PL B660 315	M. Ablikim <i>et al.</i>	(BES Collab.)
PAKHLOVA	08	PR D77 011103	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
PAKHLOVA	08A	PRL 100 062001	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
PAKHLOVA	07	PRL 98 092001	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
SETH	05A	PR D72 017501	K.K. Seth	
BAI	02C	PRL 88 101802	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	00	PRL 84 594	J.Z. Bai <i>et al.</i>	(BES Collab.)
OSTERHELD	86	SLAC-PUB-4160	A. Osterheld <i>et al.</i>	(SLAC Crystal Ball Collab.)
BRANDELIK	78C	PL 76B 361	R. Brandelik <i>et al.</i>	(DASP Collab.)
SIEGRIST	76	PRL 36 700	J.L. Siegrist <i>et al.</i>	(LBL, SLAC)