



$$I(J^P) = 0(0^-)$$


The angular distributions of the decays of the ϕ and $\bar{K}^*(892)^0$ in the $\phi\pi^+$ and $K^+\bar{K}^*(892)^0$ modes strongly indicate that the spin is zero. The parity given is that expected of a $c\bar{s}$ ground state.

D_s^\pm MASS

The fit includes D^\pm , D^0 , D_s^\pm , $D^{*\pm}$, D^{*0} , $D_s^{*\pm}$, $D_1(2420)^0$, $D_2^*(2460)^0$, and $D_{s1}(2536)^\pm$ mass and mass difference measurements. Measurements of the D_s^\pm mass with an error greater than 10 MeV are omitted from the fit and average. A number of early measurements have been omitted altogether.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1968.35 ± 0.07 OUR FIT				
1969.0 ± 1.4 OUR AVERAGE		Error includes scale factor of 1.5. See the ideogram below.		
1967.0 ± 1.0 ± 1.0	54	BARLAG	90C	ACCM π^- Cu 230 GeV
1969.3 ± 1.4 ± 1.4		ALBRECHT	88	ARG e^+e^- 9.4–10.6 GeV
1972.7 ± 1.5 ± 1.0	21	BECKER	87B	SILI 200 GeV π, K, p
1972.4 ± 3.7 ± 3.7	27	BLAYLOCK	87	MRK3 e^+e^- 4.14 GeV
1963 ± 3 ± 3	30	DERRICK	85B	HRS e^+e^- 29 GeV
1970 ± 5 ± 5	104	CHEN	83C	CLEO e^+e^- 10.5 GeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1968.3 ± 0.7 ± 0.7	290	¹ ANJOS	88	E691 Photoproduction
1980 ± 15	6	USHIDA	86	EMUL ν wideband
1973.6 ± 2.6 ± 3.0	163	ALBRECHT	85D	ARG e^+e^- 10 GeV
1948 ± 28 ± 10	65	AIHARA	84D	TPC e^+e^- 29 GeV
1975 ± 9 ± 10	49	ALTHOFF	84	TASS e^+e^- 14–25 GeV
1975 ± 4	3	BAILEY	84	ACCM hadron ⁺ Be → $\phi\pi^+X$

¹ ANJOS 88 enters the fit via $m_{D_s^\pm} - m_{D^\pm}$ (see below).



$m_{D_s^\pm} - m_{D^\pm}$

The fit includes D^\pm , D^0 , D_s^\pm , $D^{*\pm}$, D^{*0} , $D_s^{*\pm}$, $D_1(2420)^0$, $D_2^*(2460)^0$, and $D_{s1}(2536)^\pm$ mass and mass difference measurements.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
98.69±0.05 OUR FIT				
98.69±0.05 OUR AVERAGE				
98.68±0.03±0.04		AAIJ	13V LHCb	$D_s^+ \rightarrow K^+ K^- \pi^+$
99.41±0.38±0.21		ACOSTA	03D CDF2	$\bar{p}p$, $\sqrt{s}=1.96$ TeV
98.4 ±0.1 ±0.3	48k	AUBERT	02G BABR	$e^+e^- \approx \Upsilon(4S)$
99.5 ±0.6 ±0.3		BROWN	94 CLE2	$e^+e^- \approx \Upsilon(4S)$
98.5 ±1.5	555	CHEN	89 CLEO	e^+e^- 10.5 GeV
99.0 ±0.8	290	ANJOS	88 E691	Photoproduction

D_s^\pm MEAN LIFE

Measurements with an error greater than 100×10^{-15} s or with fewer than 100 events have been omitted from the Listings.

VALUE (10^{-15} s)	EVTS	DOCUMENT ID	TECN	COMMENT
501.2± 2.2 OUR AVERAGE				Error includes scale factor of 1.3. See the ideogram below.
499.5± 1.7± 0.9	116k	ADACHI	23G BEL2	$D_s^+ \rightarrow \phi \pi^+$
506.4± 3.0± 1.7±1.7		¹ AAIJ	17AN LHCb	pp at 7, 8 TeV
507.4± 5.5± 5.1	13.6k	LINK	05J FOCS	$\phi \pi^+$ and $\bar{K}^{*0} K^+$


• • • We do not use the following data for averages, fits, limits, etc. • • •

472.5 ± 17.2 ± 6.6	760	IORI	01	SELX	600 GeV Σ^- , π^- , p
518 ± 14 ± 7	1662	AITALA	99	E791	π^- nucleus, 500 GeV
486.3 ± 15.0 ⁺ _{-5.1}	2167	² BONVICINI	99	CLE2	$e^+e^- \approx \gamma(4S)$
475 ± 20 ± 7	900	FRABETTI	93F	E687	γBe , $\phi\pi^+$
500 ± 60 ± 30	104	FRABETTI	90	E687	γBe , $\phi\pi^+$
470 ± 40 ± 20	228	RAAB	88	E691	Photoproduction

¹ This AAIJ 17AN value is derived from the difference between the D_s^- and D^- widths.

The 3rd uncertainty, $\pm 1.7 \times 10^{-15}$ s, arises from the uncertainty of the D^- width.

² BONVICINI 99 obtains 1.19 ± 0.04 for the ratio of D_s^+ to D^0 lifetimes.



D_s^+ DECAY MODES

Unless otherwise noted, the branching fractions for modes with a resonance in the final state include all the decay modes of the resonance. D_s^- modes are charge conjugates of the modes below.

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Inclusive modes		
Γ_1 e^+ semileptonic	[a] (6.33 ± 0.15) %	
Γ_2 π^+ anything	(119.3 ± 1.4) %	
Γ_3 π^- anything	(43.2 ± 0.9) %	

Γ_4	π^0 anything	(123 \pm 7)%	
Γ_5	K^- anything	(18.7 \pm 0.5)%	
Γ_6	K^+ anything	(28.9 \pm 0.7)%	
Γ_7	K_S^0 anything	(19.0 \pm 1.1)%	
Γ_8	η anything	[b] (29.9 \pm 2.8)%	
Γ_9	ω anything	(6.1 \pm 1.4)%	
Γ_{10}	η' anything	[c] (10.3 \pm 1.4)%	S=1.1
Γ_{11}	$f_0(980)$ anything, $f_0 \rightarrow \pi^+\pi^-$	< 1.3 %	CL=90%
Γ_{12}	ϕ anything	(15.7 \pm 1.0)%	
Γ_{13}	K^+K^- anything	(15.8 \pm 0.7)%	
Γ_{14}	$K_S^0K^+$ anything	(5.8 \pm 0.5)%	
Γ_{15}	$K_S^0K^-$ anything	(1.9 \pm 0.4)%	
Γ_{16}	$2K_S^0$ anything	(1.70 \pm 0.32)%	
Γ_{17}	$2K^+$ anything	< 2.6 $\times 10^{-3}$	CL=90%
Γ_{18}	$2K^-$ anything	< 6 $\times 10^{-4}$	CL=90%
Γ_{19}	$2\pi^+\pi^- +$ anything	(32.8 \pm 0.7)%	

Leptonic and semileptonic modes

Γ_{20}	$e^+\nu_e$	< 8.3 $\times 10^{-5}$	CL=90%
Γ_{21}	$\mu^+\nu_\mu$	(5.37 \pm 0.11) $\times 10^{-3}$	
Γ_{22}	$\tau^+\nu_\tau$	(5.39 \pm 0.09)%	
Γ_{23}	$\gamma e^+\nu_e$	< 1.3 $\times 10^{-4}$	CL=90%
Γ_{24}	$K^+K^- e^+\nu_e$	—	
Γ_{25}	$K_S^0K_S^0 e^+\nu_e$	< 3.8 $\times 10^{-4}$	CL=90%
Γ_{26}	$\phi e^+\nu_e$	[d] (2.34 \pm 0.12)%	S=1.2
Γ_{27}	$K_1(1270)^0 e^+\nu_e$	< 4.1 $\times 10^{-4}$	CL=90%
Γ_{28}	$b_1(1235)^0 e^+\nu_e$, $b_1^0 \rightarrow \omega\pi^0$	< 6.4 $\times 10^{-4}$	CL=90%
Γ_{29}	$\phi\mu^+\nu_\mu$	(2.24 \pm 0.11)%	
Γ_{30}	$\eta e^+\nu_e$	[d] (2.27 \pm 0.06)%	
Γ_{31}	$\eta'(958) e^+\nu_e$	[d] (8.1 \pm 0.4) $\times 10^{-3}$	
Γ_{32}	$\eta\mu^+\nu_\mu$	(2.24 \pm 0.07)%	
Γ_{33}	$\eta'(958)\mu^+\nu_\mu$	(8.0 \pm 0.6) $\times 10^{-3}$	
Γ_{34}	$\omega e^+\nu_e$	[e] < 2.0 $\times 10^{-3}$	CL=90%
Γ_{35}	$K^0 e^+\nu_e$	(2.88 \pm 0.26) $\times 10^{-3}$	S=1.2
Γ_{36}	$K^*(892)^0 e^+\nu_e$	[d] (2.05 \pm 0.20) $\times 10^{-3}$	
Γ_{37}	$f_0(500) e^+\nu_e$, $f_0 \rightarrow \pi^0\pi^0$	< 7.3 $\times 10^{-4}$	CL=90%
Γ_{38}	$f_0(500) e^+\nu_e$, $f_0 \rightarrow \pi^+\pi^-$	< 3.3 $\times 10^{-4}$	CL=90%
Γ_{39}	$f_0(980) e^+\nu_e$, $f_0 \rightarrow \pi^0\pi^0$	(7.9 \pm 1.5) $\times 10^{-4}$	
Γ_{40}	$f_0(980) e^+\nu_e$, $f_0 \rightarrow \pi^+\pi^-$	(1.64 \pm 0.13) $\times 10^{-3}$	
Γ_{41}	$f_0(980)\mu^+\nu_\mu$, $f_0 \rightarrow K^+K^-$	< 5.45 $\times 10^{-4}$	CL=90%
Γ_{42}	$a_0(980)^0 e^+\nu_e$, $a_0^0 \rightarrow \pi^0\eta$	< 1.2 $\times 10^{-4}$	CL=90%
Γ_{43}	$\pi^0 e^+\nu_e$	< 6.4 $\times 10^{-5}$	CL=90%

Hadronic modes with a $K\bar{K}$ pair

Γ_{44}	$K^+ K_S^0$		(1.500 ± 0.014) %	
Γ_{45}	$K^+ K_L^0$		(1.49 ± 0.06) %	
Γ_{46}	$K_S^0 K_L^0 \pi^+$		(1.86 ± 0.07) %	
Γ_{47}	$\phi(1020) \pi^+, \phi \rightarrow K_L^0 K_S^0$		(1.31 ± 0.06) %	
Γ_{48}	$K_L^0 K^*(892)^+, K^{*+} \rightarrow K_S^0 \pi^+$		(3.59 ± 0.32) × 10 ⁻³	
Γ_{49}	$K_S^0 K^*(892)^+, K^{*+} \rightarrow K_L^0 \pi^+$		(2.68 ± 0.33) × 10 ⁻³	
Γ_{50}	$\phi(1680) \pi^+, \phi \rightarrow K_L^0 K_S^0$		(6.1 ± 2.1) × 10 ⁻⁴	
Γ_{51}	$K^+ \bar{K}^0$		(2.95 ± 0.14) %	
Γ_{52}	$K^+ K^- \pi^+$	[f]	(5.45 ± 0.08) %	S=1.3
Γ_{53}	$\phi \pi^+$	[d,g]	(4.5 ± 0.4) %	
Γ_{54}	$\phi \pi^+, \phi \rightarrow K^+ K^-$	[g]	(2.25 ± 0.05) %	
Γ_{55}	$K^+ \bar{K}^*(892)^0$		(12.7 ^{+4.0} _{-3.1}) %	
Γ_{56}	$K^+ \bar{K}^*(892)^0, \bar{K}^{*0} \rightarrow K^- \pi^+$		(2.61 ± 0.05) %	
Γ_{57}	$K^+ \bar{K}^*(892)^0, \bar{K}^{*0} \rightarrow K_S^0 \pi^0$		(4.8 ± 0.4) × 10 ⁻³	
Γ_{58}	$f_0(980) \pi^+, f_0 \rightarrow K^+ K^-$		(1.12 ± 0.19) %	
Γ_{59}	$f_0(1370) \pi^+, f_0 \rightarrow K^+ K^-$		(7.2 ± 3.0) × 10 ⁻⁴	
Γ_{60}	$f_0(1710) \pi^+, f_0 \rightarrow K^+ K^-$		(6.8 ± 2.8) × 10 ⁻⁴	
Γ_{61}	$a_0(980)^+ \pi^0, a_0^+ \rightarrow K^+ K_S^0$		(1.1 ± 0.4) × 10 ⁻³	
Γ_{62}	$a_0(1710)^+ \pi^0, a_0^+ \rightarrow K^+ K_S^0$		(3.5 ± 0.6) × 10 ⁻³	
Γ_{63}	$K^+ \bar{K}_0^*(1430)^0, \bar{K}_0^* \rightarrow K^- \pi^+$		(1.79 ± 0.26) × 10 ⁻³	
Γ_{64}	$K^+ \bar{K}_0^*(1430)^0, \bar{K}_0^* \rightarrow K_S^0 \pi^0$		(8.8 ± 2.8) × 10 ⁻⁴	
Γ_{65}	$K^+ K_S^0 \pi^0$		(1.471 ± 0.028) %	
Γ_{66}	$K^*(892)^+ K_S^0, K^{*+} \rightarrow K^+ \pi^0$		(2.04 ± 0.32) × 10 ⁻³	
Γ_{67}	$2K_S^0 \pi^+$		(7.32 ± 0.14) × 10 ⁻³	
Γ_{68}	$f_0(980) \pi^+, f_0 \rightarrow K_S^0 K_S^0$	<	1.8 × 10 ⁻⁴	CL=90%
Γ_{69}	$f_0(1710) \pi^+, f_0 \rightarrow K_S^0 K_S^0$		(3.39 ± 0.31) × 10 ⁻³	
Γ_{70}	$K^*(892)^+ K_S^0, K^{*+} \rightarrow K_S^0 \pi^+$		(3.19 ± 0.29) × 10 ⁻³	
Γ_{71}	$K^0 \bar{K}^0 \pi^+$		—	
Γ_{72}	$K^*(892)^+ \bar{K}^0$	[d]	(5.4 ± 1.2) %	
Γ_{73}	$\phi \rho^+$		(4.9 ± 0.8) %	S=3.0
Γ_{74}	$K^+ K^- \pi^+ \pi^0$		(5.53 ± 0.15) %	S=1.3
Γ_{75}	$\bar{K}_1(1270)^0 K^+, \bar{K}_1(1270)^0 \rightarrow K^- \rho^+$		(5.7 ± 0.6) × 10 ⁻³	

Γ_{76}	$\bar{K}_1(1270)^0 K^+,$ $\bar{K}_1(1270)^0 \rightarrow K^*(892)\pi$	(1.31 \pm 0.25) %	
Γ_{77}	$\bar{K}_1(1400)^0 K^+,$ $\bar{K}_1(1400)^0 \rightarrow K^*(892)\pi$	(2.0 \pm 0.4) %	
Γ_{78}	$a_0(980)^0 \rho^+, a_0^0 \rightarrow K^+ K^-$	(1.9 \pm 0.4) $\times 10^{-3}$	
Γ_{79}	$f_1(1420)^0 \pi^+, f_1(1420)^0 \rightarrow$ $K^*(892)^\mp K^\pm$	(3.9 \pm 0.7) $\times 10^{-3}$	
Γ_{80}	$f_1(1420)^0 \pi^+, f_1^0 \rightarrow$ $a_0(980)^0 \pi^0, a_0^0 \rightarrow$ $K^+ K^-$	(4.0 \pm 1.4) $\times 10^{-4}$	
Γ_{81}	$\eta(1475)\pi^+, \eta \rightarrow$ $a_0(980)^0 \pi^0, a_0^0 \rightarrow$ $K^+ K^-$	(7.0 \pm 2.8) $\times 10^{-4}$	
Γ_{82}	$K_S^0 K^- 2\pi^+$	(1.569 \pm 0.028) %	
Γ_{83}	$K^+ K^- K_S^0 \pi^+$	(1.27 \pm 0.15) $\times 10^{-4}$	
Γ_{84}	$K^*(892)^+ \bar{K}^*(892)^0$	[d] (5.64 \pm 0.35) %	
Γ_{85}	$\eta(1475) K_S^0, \eta \rightarrow$ $K^*(892)^0 \pi^+, K^{*0} \rightarrow$ $K^- \pi^+$	(3.5 \pm 1.0) $\times 10^{-4}$	
Γ_{86}	$\eta(1475)\pi^+, \eta \rightarrow$ $\bar{K}^*(892)^+ K^-, \bar{K}^{*+} \rightarrow$ $K_S^0 \pi^+$	(3.5 \pm 1.0) $\times 10^{-4}$	
Γ_{87}	$\eta(1475)\pi^+, \eta \rightarrow$ $a_0(980)^- \pi^+, a_0^- \rightarrow$ $K_S^0 K^-$	(1.7 \pm 0.9) $\times 10^{-3}$	
Γ_{88}	$f_1(1285)\pi^+, f_1 \rightarrow$ $a_0(980)^- \pi^+, a_0^- \rightarrow$ $K_S^0 K^-$	(3.5 \pm 0.8) $\times 10^{-4}$	
Γ_{89}	$K^+ K_S^0 \pi^+ \pi^-$	(9.34 \pm 0.22) $\times 10^{-3}$	
Γ_{90}	$K^+ K^- 2\pi^+ \pi^-$	(6.6 \pm 0.6) $\times 10^{-3}$	
Γ_{91}	$\phi 2\pi^+ \pi^-$	[d] (1.21 \pm 0.16) %	
Γ_{92}	$\phi \rho^0 \pi^+, \phi \rightarrow K^+ K^-$	(4.9 \pm 0.7) $\times 10^{-3}$	
Γ_{93}	$\phi a_1(1260)^+, \phi \rightarrow$ $K^+ K^-, a_1^+ \rightarrow$ $\rho^0 \pi^+$	(7.5 \pm 1.2) $\times 10^{-3}$	
Γ_{94}	$\phi 2\pi^+ \pi^- \text{ non-}\rho, \phi \rightarrow$ $K^+ K^-$	(1.4 \pm 0.5) $\times 10^{-3}$	
Γ_{95}	$K^+ K^- \rho^0 \pi^+ \text{ non-}\phi$	< 2.0 $\times 10^{-4}$ CL=90%	
Γ_{96}	$K^+ K^- 2\pi^+ \pi^- \text{ nonresonant}$	(1.0 \pm 0.4) $\times 10^{-3}$	
Γ_{97}	$2K_S^0 2\pi^+ \pi^-$	(8.0 \pm 3.3) $\times 10^{-4}$	
Hadronic modes without K's			
Γ_{98}	$\pi^+ \pi^0$	< 1.2 $\times 10^{-4}$ CL=90%	
Γ_{99}	$2\pi^+ \pi^-$	(1.090 \pm 0.014) %	

Γ_{100}	$\rho^0 \pi^+$		$(1.14 \pm 0.16) \times 10^{-4}$	
Γ_{101}	$\omega \pi^+, \omega \rightarrow \pi^+ \pi^-$		$(3.9 \pm 0.4) \times 10^{-5}$	
Γ_{102}	$\pi^+ (\pi^+ \pi^-)_{S\text{-wave}}$	[h]	$(9.23 \pm 0.13) \times 10^{-3}$	
Γ_{103}	$f_0(980) \pi^+, f_0 \rightarrow \pi^+ \pi^-$			
Γ_{104}	$f_0(1370) \pi^+, f_0 \rightarrow \pi^+ \pi^-$			
Γ_{105}	$f_0(1500) \pi^+, f_0 \rightarrow \pi^+ \pi^-$			
Γ_{106}	$f_2(1270) \pi^+, f_2 \rightarrow \pi^+ \pi^-$		$(1.42 \pm 0.10) \times 10^{-3}$	
Γ_{107}	$f_2'(1525)^0 \pi^+, f_2' \rightarrow \pi^+ \pi^-$		$(5.8 \pm 2.0) \times 10^{-6}$	
Γ_{108}	$\rho(1450)^0 \pi^+, \rho^0 \rightarrow \pi^+ \pi^-$		$(1.8 \pm 0.6) \times 10^{-4}$	
Γ_{109}	$\rho(1700)^0 \pi^+, \rho^0 \rightarrow \pi^+ \pi^-$		$(4 \pm 4) \times 10^{-5}$	
Γ_{110}	$\pi^+ 2\pi^0$		$(5.2 \pm 0.5) \times 10^{-3}$	S=1.1
Γ_{111}	$f_0(980) \pi^+, f_0 \rightarrow \pi^0 \pi^0$		$(2.9 \pm 0.6) \times 10^{-3}$	
Γ_{112}	$f_0(1370) \pi^+, f_0 \rightarrow \pi^0 \pi^0$		$(1.3 \pm 0.6) \times 10^{-3}$	
Γ_{113}	$f_2(1270) \pi^+, f_2 \rightarrow \pi^0 \pi^0$		$(5.0 \pm 3.5) \times 10^{-4}$	
Γ_{114}	$2\pi^+ \pi^- \pi^0$		—	
Γ_{115}	$\omega \pi^+$	[d]	$(1.93 \pm 0.30) \times 10^{-3}$	
Γ_{116}	$\eta \pi^+$	[d]	$(1.686 \pm 0.027) \%$	
Γ_{117}	$(2\pi^+ \pi^- \pi^0)_{\text{non-}\eta}$		$(2.04 \pm 0.09) \%$	
Γ_{118}	$f_0(1370) \rho^+, f_0 \rho^+ \rightarrow$ $2\pi^+ \pi^- \pi^0$		$(5.1 \pm 0.9) \times 10^{-3}$	
Γ_{119}	$f_0(980)^0 \rho^+, f_0^0 \rho^+ \rightarrow$ $2\pi^+ \pi^- \pi^0$		$(2.6 \pm 0.5) \times 10^{-3}$	
Γ_{120}	$f_2(1270)^0 \rho^+,$ $f_2(1270)^0 \rho^+ \rightarrow$ $2\pi^+ \pi^- \pi^0$		$(1.9 \pm 0.4) \times 10^{-3}$	
Γ_{121}	$(\rho^+ \rho^0)_{S\text{-wave}} \rightarrow$ $2\pi^+ \pi^- \pi^0$		$(7.1 \pm 2.8) \times 10^{-4}$	
Γ_{122}	$(\rho(1450)^+ \rho^0)_{S\text{-wave}} \rightarrow$ $2\pi^+ \pi^- \pi^0$		$(9.4 \pm 3.1) \times 10^{-4}$	
Γ_{123}	$(\rho^+ \rho(1450)^0)_{P\text{-wave}} \rightarrow$ $2\pi^+ \pi^- \pi^0$		$(1.75 \pm 0.29) \times 10^{-3}$	
Γ_{124}	$\phi \pi^+, \phi \rightarrow \rho \pi$		$(5.08 \pm 0.35) \times 10^{-3}$	
Γ_{125}	$\omega \pi^+, \omega \rightarrow \rho \pi$		$(1.41 \pm 0.19) \times 10^{-3}$	
Γ_{126}	$a_1(1260)^+ \pi^0, a_1^+ \rightarrow$ $(\rho^0 \pi^+)_{S\text{-wave}}$		$(2.6 \pm 0.4) \times 10^{-3}$	
Γ_{127}	$a_1(1260)^0 \pi^+, a_1^0 \rightarrow$ $(\rho \pi)_{S\text{-wave}}$		$(1.3 \pm 0.5) \times 10^{-3}$	
Γ_{128}	$\pi(1300)^0 \pi^+, \pi^0 \rightarrow$ $(\rho \pi)_{P\text{-wave}}$		$(2.4 \pm 0.7) \times 10^{-3}$	
Γ_{129}	$3\pi^+ 2\pi^-$		$(8.0 \pm 0.8) \times 10^{-3}$	
Γ_{130}	$2\pi^+ \pi^- 2\pi^0$		$(4.41 \pm 0.20) \%$	
Γ_{131}	$\omega \rho^+, \omega \rightarrow \pi^+ \pi^- \pi^0$		$(8.8 \pm 0.9) \times 10^{-3}$	
Γ_{132}	$\phi \rho^+, \phi \rightarrow \pi^+ \pi^- \pi^0$		$(6.1 \pm 0.6) \times 10^{-3}$	
Γ_{133}	$\rho(1450)^+ \pi^0, \rho(1450)^+ \rightarrow$ $\omega \pi^+, \omega \rightarrow \pi^+ \pi^- \pi^0$		$(3.5 \pm 0.5) \times 10^{-3}$	

Γ_{134}	$a_1(1260)^0 \rho^+, a_1(1260)^0 \rightarrow \rho^+ \pi^-$	(5.0 \pm 0.4) $\times 10^{-3}$	
Γ_{135}	$a_1(1260)^0 \rho^+, a_1(1260)^0 \rightarrow \rho^- \pi^+$	(3.26 \pm 0.27) $\times 10^{-3}$	
Γ_{136}	$a_1(1260)^+ \rho^0, a_1(1260)^+ \rightarrow \rho^+ \pi^0$	(7.3 \pm 1.0) $\times 10^{-3}$	
Γ_{137}	$b_1(1235)^+ \pi^0, b_1(1235)^+ \rightarrow \omega \pi^+, \omega \rightarrow \pi^+ \pi^- \pi^0$	(4.8 \pm 0.6) $\times 10^{-3}$	
Γ_{138}	$b_1(1235)^0 \pi^+, b_1(1235)^0 \rightarrow \omega \pi^0, \omega \rightarrow \pi^+ \pi^- \pi^0$	(6.4 \pm 0.6) $\times 10^{-3}$	
Γ_{139}	$\eta \rho^+$	[d] (8.9 \pm 0.8) %	
Γ_{140}	$\eta \pi^+ \pi^0$	(9.10 \pm 0.17) %	
Γ_{141}	$\eta(\pi^+ \pi^0)_{P\text{-wave}}$	(4.9 \pm 3.0) $\times 10^{-3}$	
Γ_{142}	$a_0(980)^{+0} \pi^{0+}, a_0(980)^{+0} \rightarrow \eta \pi^{+0}$	(2.1 \pm 0.4) %	
Γ_{143}	$\omega \pi^+ \pi^0$	[d] (2.8 \pm 0.7) %	
Γ_{144}	$2\pi^+ \pi^- \eta$	(3.08 \pm 0.08) %	
Γ_{145}	$a_1(1260)^+ \eta, a_1^+ \rightarrow \rho(770)^0 \pi^+, \rho^0 \rightarrow \pi^+ \pi^-$	(1.71 \pm 0.14) %	
Γ_{146}	$a_1(1260)^+ \eta, a_1^+ \rightarrow f_0(500) \pi^+, f_0 \rightarrow \pi^+ \pi^-$	(2.5 \pm 0.9) $\times 10^{-3}$	
Γ_{147}	$a_0(980)^+ \rho(770)^0, a_0^+ \rightarrow \eta \pi^+$	(2.1 \pm 0.9) $\times 10^{-3}$	
Γ_{148}	$\eta(1405) \pi^+, \eta \rightarrow a_0(980)^- \pi^+, a_0^- \rightarrow \eta \pi^-$	(2.2 \pm 0.7) $\times 10^{-4}$	
Γ_{149}	$\eta(1405) \pi^+, \eta \rightarrow a_0(980)^+ \pi^-, a_0^+ \rightarrow \eta \pi^+$	(2.2 \pm 0.7) $\times 10^{-4}$	
Γ_{150}	$f_1(1420) \pi^+, f_1 \rightarrow a_0(980)^- \pi^+, a_0^- \rightarrow \eta \pi^-$	(5.9 \pm 1.8) $\times 10^{-4}$	
Γ_{151}	$f_1(1420) \pi^+, f_1 \rightarrow a_0(980)^+ \pi^-, a_0^+ \rightarrow \eta \pi^+$	(5.2 \pm 1.8) $\times 10^{-4}$	
Γ_{152}	$3\pi^+ 2\pi^- \pi^0$	(4.9 \pm 3.2) %	
Γ_{153}	$\omega 2\pi^+ \pi^-$	[d] (1.6 \pm 0.5) %	
Γ_{154}	$\eta'(958) \pi^+$	[c,d] (3.95 \pm 0.08) %	
Γ_{155}	$3\pi^+ 2\pi^- 2\pi^0$	—	
Γ_{156}	$\omega \eta \pi^+$	[d] (5.4 \pm 1.3) $\times 10^{-3}$	
Γ_{157}	$\eta'(958) \rho^+$	[c,d] (5.8 \pm 1.5) %	
Γ_{158}	$\eta'(958) \pi^+ \pi^0$	(6.14 \pm 0.18) %	
Γ_{159}	$\eta'(958) \pi^+ \pi^0$ nonresonant	< 5.1 %	CL=90%

Modes with one or three K's

Γ_{160}	$K^+ \pi^0$	(7.5 \pm 0.5) $\times 10^{-4}$	
Γ_{161}	$K_S^0 \pi^+$	(1.22 \pm 0.04) $\times 10^{-3}$	
Γ_{162}	$K^+ \eta$	[d] (1.76 \pm 0.08) $\times 10^{-3}$	

Γ_{163}	$K^+\omega$	[d]	$(9.9 \pm 1.5) \times 10^{-4}$	
Γ_{164}	$K^+\eta'(958)$	[d]	$(2.68 \pm 0.24) \times 10^{-3}$	
Γ_{165}	$K^+\pi^+\pi^-$		$(6.23 \pm 0.10) \times 10^{-3}$	
Γ_{166}	$K^+\rho^0$		$(2.18 \pm 0.25) \times 10^{-3}$	
Γ_{167}	$K^+\rho(1450)^0, \rho^0 \rightarrow \pi^+\pi^-$		$(7.3 \pm 1.7) \times 10^{-4}$	
Γ_{168}	$K^+f_0(500), f_0 \rightarrow \pi^+\pi^-$		$(4.5 \pm 3.0) \times 10^{-4}$	
Γ_{169}	$K^+f_0(980), f_0 \rightarrow \pi^+\pi^-$		$(2.8 \pm 1.1) \times 10^{-4}$	
Γ_{170}	$K^+f_0(1370), f_0 \rightarrow \pi^+\pi^-$		$(1.2 \pm 0.6) \times 10^{-3}$	
Γ_{171}	$K^*(892)^0\pi^+, K^{*0} \rightarrow$		$(1.68 \pm 0.26) \times 10^{-3}$	
Γ_{172}	$K^+\pi^-$ $K^*(1410)^0\pi^+, K^{*0} \rightarrow$		$(6 \pm 4) \times 10^{-4}$	
Γ_{173}	$K^+\pi^-$ $K^*(1430)^0\pi^+, K^{*0} \rightarrow$		$(9.4 \pm 3.2) \times 10^{-4}$	
Γ_{174}	$K^+\pi^+\pi^-$ nonresonant		$(9.9 \pm 3.2) \times 10^{-4}$	
Γ_{175}	$K^0\pi^+\pi^0$			
Γ_{176}	$K_S^0\pi^+\pi^0$		$(5.09 \pm 0.22) \times 10^{-3}$	
Γ_{177}	$K_S^0\rho(770)^+, \rho^+ \rightarrow \pi^+\pi^0$		$(2.6 \pm 0.4) \times 10^{-3}$	
Γ_{178}	$K_S^0\rho(1450)^+, \rho^+ \rightarrow \pi^+\pi^0$		$(1.04 \pm 0.32) \times 10^{-3}$	
Γ_{179}	$K^*(892)^0\pi^+, K^{*0} \rightarrow K_S^0\pi^0$		$(4.3 \pm 1.2) \times 10^{-4}$	
Γ_{180}	$K^*(892)^+\pi^0, K^{*+} \rightarrow$		$(2.3 \pm 0.7) \times 10^{-4}$	
Γ_{181}	$K_S^0\pi^+$ $K^*(1410)^0\pi^+, K^{*0} \rightarrow$		$(1.7 \pm 0.9) \times 10^{-4}$	
Γ_{182}	$K_S^0\pi^0$ $K_S^0 2\pi^+\pi^-$		$(2.8 \pm 1.0) \times 10^{-3}$	
Γ_{183}	$K^+\pi^+\pi^-\pi^0$		$(9.7 \pm 0.6) \times 10^{-3}$	
Γ_{184}	$K^*(892)^0\rho^+, K^{*0} \rightarrow K^+\pi^-$		$(3.9 \pm 0.4) \times 10^{-3}$	
Γ_{185}	$K^*(892)^+\rho^0, K^{*+} \rightarrow K^+\pi^0$		$(4.2 \pm 1.2) \times 10^{-4}$	
Γ_{186}	$K_1(1270)^0\pi^+, K_1^0 \rightarrow K^+\rho^-$		$(3.9 \pm 1.3) \times 10^{-4}$	
Γ_{187}	$K_1(1400)^0\pi^+, K_1^0 \rightarrow$		$(5.4 \pm 0.9) \times 10^{-4}$	
Γ_{188}	$K^*(890)^+\pi^-, K^{*+} \rightarrow$ $K^+\pi^0$ $K_1(1400)^0\pi^+, K_1^0 \rightarrow$ $K^*(890)^0\pi^0, K^{*0} \rightarrow$		$(5.9 \pm 1.0) \times 10^{-4}$	
Γ_{189}	$K^+a_1(1260)^0, a_1 \rightarrow \rho^+\pi^-$		$(1.8 \pm 1.1) \times 10^{-4}$	
Γ_{190}	$K^+a_1(1260)^0, a_1 \rightarrow \rho^-\pi^+$		$(1.8 \pm 1.1) \times 10^{-4}$	
Γ_{191}	$K^+\pi^+\pi^-\pi^0$ nonresonant		$(9.2 \pm 2.4) \times 10^{-4}$	
Γ_{192}	$(K^+\pi^0)_{P\text{-wave}}\rho^0$		$(1.01 \pm 0.21) \times 10^{-3}$	
Γ_{193}	$K^+\omega\pi^0$	[d]	$< 8.2 \times 10^{-3}$	CL=90%
Γ_{194}	$K^+\omega\pi^+\pi^-$	[d]	$< 5.4 \times 10^{-3}$	CL=90%
Γ_{195}	$K^+\omega\eta$	[d]	$< 7.9 \times 10^{-3}$	CL=90%
Γ_{196}	$2K^+K^-$		$(2.18 \pm 0.20) \times 10^{-4}$	
Γ_{197}	$\phi K^+, \phi \rightarrow K^+K^-$		$(8.9 \pm 2.0) \times 10^{-5}$	

Radiative decays

$$\Gamma_{198} \rho(770)^+ \gamma < 6.1 \times 10^{-4} \text{ CL}=90\%$$

Doubly Cabibbo-suppressed modes

$$\Gamma_{199} 2K^+ \pi^- \quad (1.293 \pm 0.027) \times 10^{-4} \quad S=1.1$$

$$\Gamma_{200} K^+ K^*(892)^0, K^{*0} \rightarrow (6.1 \pm 3.4) \times 10^{-5}$$

$$\Gamma_{201} 2K^+ \pi^- \pi^0 < 1.7 \times 10^{-4} \text{ CL}=90\%$$

Baryon-antibaryon mode

$$\Gamma_{202} \rho \bar{n} \quad (1.22 \pm 0.11) \times 10^{-3}$$

$$\Gamma_{203} \rho \bar{p} e^+ \nu_e < 2.0 \times 10^{-4} \text{ CL}=90\%$$

 **$\Delta C = 1$ weak neutral current (C1) modes,
Lepton family number (LF), or
Lepton number (L) violating modes**

$$\Gamma_{204} \pi^+ e^+ e^- \quad [i] < 5.5 \times 10^{-6} \text{ CL}=90\%$$

$$\Gamma_{205} \pi^+ \phi, \phi \rightarrow e^+ e^- \quad [j] (1.17 \pm 0.22) \times 10^{-5}$$

$$\Gamma_{206} \pi^+ \pi^0 e^+ e^- < 7.0 \times 10^{-5} \text{ CL}=90\%$$

$$\Gamma_{207} \rho^+ \phi, \phi \rightarrow e^+ e^- \quad (2.4 \pm 0.7) \times 10^{-5}$$

$$\Gamma_{208} \pi^+ \mu^+ \mu^- \quad [i] < 1.8 \times 10^{-7} \text{ CL}=90\%$$

$$\Gamma_{209} K^+ e^+ e^- \quad C1 < 3.7 \times 10^{-6} \text{ CL}=90\%$$

$$\Gamma_{210} K^+ \pi^0 e^+ e^- < 7.1 \times 10^{-5} \text{ CL}=90\%$$

$$\Gamma_{211} K_S^0 \pi^+ e^+ e^- < 8.1 \times 10^{-5} \text{ CL}=90\%$$

$$\Gamma_{212} K^+ \mu^+ \mu^- \quad C1 < 1.4 \times 10^{-7} \text{ CL}=90\%$$

$$\Gamma_{213} K^*(892)^+ \mu^+ \mu^- \quad C1 < 1.4 \times 10^{-3} \text{ CL}=90\%$$

$$\Gamma_{214} \pi^+ e^+ \mu^- \quad LF < 1.1 \times 10^{-6} \text{ CL}=90\%$$

$$\Gamma_{215} \pi^+ e^- \mu^+ \quad LF < 9.4 \times 10^{-7} \text{ CL}=90\%$$

$$\Gamma_{216} K^+ e^+ \mu^- \quad LF < 7.9 \times 10^{-7} \text{ CL}=90\%$$

$$\Gamma_{217} K^+ e^- \mu^+ \quad LF < 5.6 \times 10^{-7} \text{ CL}=90\%$$

$$\Gamma_{218} \pi^- 2e^+ \quad L < 1.4 \times 10^{-6} \text{ CL}=90\%$$

$$\Gamma_{219} \pi^- 2\mu^+ \quad L < 8.6 \times 10^{-8} \text{ CL}=90\%$$

$$\Gamma_{220} \pi^- e^+ \mu^+ \quad L < 6.3 \times 10^{-7} \text{ CL}=90\%$$

$$\Gamma_{221} K^- 2e^+ \quad L < 7.7 \times 10^{-7} \text{ CL}=90\%$$

$$\Gamma_{222} \pi^- \pi^0 e^+ e^+ < 2.9 \times 10^{-5} \text{ CL}=90\%$$

$$\Gamma_{223} K^- \pi^0 e^+ e^+ < 3.4 \times 10^{-5} \text{ CL}=90\%$$

$$\Gamma_{224} K_S^0 \pi^- e^+ e^+ < 1.3 \times 10^{-5} \text{ CL}=90\%$$

$$\Gamma_{225} K_S^0 K^- e^+ e^+ < 2.9 \times 10^{-5} \text{ CL}=90\%$$

$$\Gamma_{226} \phi \pi^- e^+ e^+ < 6.9 \times 10^{-5} \text{ CL}=90\%$$

$$\Gamma_{227} \phi K^- e^+ e^+ < 9.9 \times 10^{-5} \text{ CL}=90\%$$

$$\Gamma_{228} K^- 2\mu^+ \quad L < 2.6 \times 10^{-8} \text{ CL}=90\%$$

$$\Gamma_{229} K^- e^+ \mu^+ \quad L < 2.6 \times 10^{-7} \text{ CL}=90\%$$

$$\Gamma_{230} K^*(892)^- 2\mu^+ \quad L < 1.4 \times 10^{-3} \text{ CL}=90\%$$

[a] This is the purely e^+ semileptonic branching fraction: the e^+ fraction from τ^+ decays has been subtracted off. The sum of our (non- τ) e^+

exclusive fractions — an $e^+ \nu_e$ with an η , η' , ϕ , K^0 , or K^{*0} — is 5.99 ± 0.31 %.

- [b] This fraction includes η from η' decays.
- [c] The sum of our exclusive η' fractions — $\eta' e^+ \nu_e$, $\eta' \mu^+ \nu_\mu$, $\eta' \pi^+$, $\eta' \rho^+$, and $\eta' K^+$ — is 11.8 ± 1.6 %.
- [d] This branching fraction includes all the decay modes of the final-state resonance.
- [e] A test for $u\bar{u}$ or $d\bar{d}$ content in the D_s^+ . Neither Cabibbo-favored nor Cabibbo-suppressed decays can contribute, and ω - ϕ mixing is an unlikely explanation for any fraction above about 2×10^{-4} .
- [f] The branching fraction for this mode may differ from the sum of the submodes that contribute to it, due to interference effects. See the relevant papers.
- [g] We decouple the $D_s^+ \rightarrow \phi \pi^+$ branching fraction obtained from mass projections (and used to get some of the other branching fractions) from the $D_s^+ \rightarrow \phi \pi^+$, $\phi \rightarrow K^+ K^-$ branching fraction obtained from the Dalitz-plot analysis of $D_s^+ \rightarrow K^+ K^- \pi^+$. That is, the ratio of these two branching fractions is not exactly the $\phi \rightarrow K^+ K^-$ branching fraction 0.491.
- [h] This is the average of a model-independent and a K -matrix parametrization of the $\pi^+ \pi^-$ S -wave and is a sum over several f_0 mesons.
- [i] This mode is not a useful test for a $\Delta C=1$ weak neutral current because both quarks must change flavor in this decay.
- [j] This is *not* a test for the $\Delta C=1$ weak neutral current, but leads to the $\pi^+ \ell^+ \ell^-$ final state.

FIT INFORMATION

An overall fit to 18 branching ratios uses 32 measurements to determine 12 parameters. The overall fit has a $\chi^2 = 12.1$ for 20 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$.

x52	11										
x74	-1	1									
x82	1	2	2								
x89	0	1	1	1							
x99	1	1	0	1	0						
x115	0	0	0	0	0	0					
x116	2	2	-1	0	0	0	1				
x163	0	0	0	0	0	0	0	0			
x165	0	2	1	1	1	0	0	0	0		
x183	0	0	0	0	0	0	0	0	26	0	
x199	8	72	1	2	0	1	0	1	0	1	
	x44	x52	x74	x82	x89	x99	x115	x116	x163	x165	
x199	0										
	x183										

D_s^+ BRANCHING RATIOS

A number of older, now obsolete results have been omitted. They may be found in earlier editions.

Inclusive modes

$\Gamma(e^+ \text{ semileptonic})/\Gamma_{\text{total}}$

Γ_1/Γ

This is the purely e^+ semileptonic branching fraction: the e^+ fraction from τ^+ decays has been subtracted off.

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
6.33±0.15 OUR AVERAGE				
6.30±0.13±0.10	17k	^{1,2} ABLIKIM	21AC BES3	e^+e^- at 4.178–4.230 GeV
6.52±0.39±0.15	0.5k	³ ASNER	10 CLEO	e^+e^- at 3774 MeV

¹ ABLIKIM 21AC finds that the ratio of the D_s^+ and D^0 semielectronic widths is $0.790 \pm 0.016 \pm 0.020$.

² ABLIKIM 21AC reports a value of $(6.30 \pm 0.13 \pm 0.09 \pm 0.04) \times 10^{-2}$, where the last uncertainty is an external systematic from $B(D_s^+ \rightarrow \tau\nu)$. We have added the systematic uncertainties in quadrature.

³ Using the D_s^+ and D^0 lifetimes, ASNER 10 finds that the ratio of the D_s^+ and D^0 semileptonic widths is $0.828 \pm 0.051 \pm 0.025$.

$\Gamma(\pi^+ \text{ anything})/\Gamma_{\text{total}}$

Γ_2/Γ

Events with two π^+ 's count twice, etc. But π^+ 's from $K_S^0 \rightarrow \pi^+\pi^-$ are not included.

VALUE (%)	DOCUMENT ID	TECN	COMMENT
119.3±1.2±0.7	DOBBS	09	CLEO e^+e^- at 4170 MeV

$\Gamma(\pi^- \text{ anything})/\Gamma_{\text{total}}$ Γ_3/Γ

Events with two π^- 's count twice, etc. But π^- 's from $K_S^0 \rightarrow \pi^+\pi^-$ are not included.

VALUE (%)	DOCUMENT ID	TECN	COMMENT
43.2±0.9±0.3	DOBBS	09	CLEO e^+e^- at 4170 MeV

$\Gamma(\pi^0 \text{ anything})/\Gamma_{\text{total}}$ Γ_4/Γ

Events with two π^0 's count twice, etc. But π^0 's from $K_S^0 \rightarrow 2\pi^0$ are not included.

VALUE (%)	DOCUMENT ID	TECN	COMMENT
123.4±3.8±5.3	DOBBS	09	CLEO e^+e^- at 4170 MeV

$\Gamma(K^- \text{ anything})/\Gamma_{\text{total}}$ Γ_5/Γ

VALUE (%)	DOCUMENT ID	TECN	COMMENT
18.7±0.5±0.2	DOBBS	09	CLEO e^+e^- at 4170 MeV

$\Gamma(K^+ \text{ anything})/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE (%)	DOCUMENT ID	TECN	COMMENT
28.9±0.6±0.3	DOBBS	09	CLEO e^+e^- at 4170 MeV

$\Gamma(K_S^0 \text{ anything})/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE (%)	DOCUMENT ID	TECN	COMMENT
19.0±1.0±0.4	DOBBS	09	CLEO e^+e^- at 4170 MeV

$\Gamma(\eta \text{ anything})/\Gamma_{\text{total}}$ Γ_8/Γ

This ratio includes η particles from η' decays.

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
29.9±2.2±1.7		DOBBS	09	CLEO e^+e^- at 4170 MeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

23.5±3.1±2.0	674 ± 91	HUANG	06B	CLEO See DOBBS 09
--------------	----------	-------	-----	-------------------

$\Gamma(\omega \text{ anything})/\Gamma_{\text{total}}$ Γ_9/Γ

VALUE (%)	DOCUMENT ID	TECN	COMMENT
6.1±1.4±0.3	DOBBS	09	CLEO e^+e^- at 4170 MeV

$\Gamma(\eta' \text{ anything})/\Gamma_{\text{total}}$ Γ_{10}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
10.3±1.4 OUR AVERAGE		Error includes scale factor of 1.1.		

8.8±1.8±0.5	68	ABLIKIM	15Z	BES3 482 pb ⁻¹ , 4009 MeV
-------------	----	---------	-----	--------------------------------------

11.7±1.7±0.7		DOBBS	09	CLEO e^+e^- at 4170 MeV
--------------	--	-------	----	---------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

8.7±1.9±0.8	68	HUANG	06B	CLEO See DOBBS 09
-------------	----	-------	-----	-------------------

$\Gamma(f_0(980) \text{ anything}, f_0 \rightarrow \pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE (%)	CL%	DOCUMENT ID	TECN	COMMENT
<1.3	90	DOBBS	09	CLEO e^+e^- at 4170 MeV

$\Gamma(\phi \text{ anything})/\Gamma_{\text{total}}$ Γ_{12}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
15.7±0.8±0.6		DOBBS	09	CLEO e^+e^- at 4170 MeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

16.1±1.2±1.1 398 ± 27 HUANG 06B CLEO See DOBBS 09

$\Gamma(K^+ K^- \text{ anything})/\Gamma_{\text{total}}$ **Γ_{13}/Γ**

VALUE (%)		DOCUMENT ID	TECN	COMMENT
15.8±0.6±0.3		DOBBS	09	CLEO $e^+ e^-$ at 4170 MeV

$\Gamma(K_S^0 K^+ \text{ anything})/\Gamma_{\text{total}}$ **Γ_{14}/Γ**

VALUE (%)		DOCUMENT ID	TECN	COMMENT
5.8±0.5±0.1		DOBBS	09	CLEO $e^+ e^-$ at 4170 MeV

$\Gamma(K_S^0 K^- \text{ anything})/\Gamma_{\text{total}}$ **Γ_{15}/Γ**

VALUE (%)		DOCUMENT ID	TECN	COMMENT
1.9±0.4±0.1		DOBBS	09	CLEO $e^+ e^-$ at 4170 MeV

$\Gamma(K_S^0 K_L^0 \pi^+)/\Gamma_{\text{total}}$ **Γ_{46}/Γ**

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
1.86±0.06±0.03	2.35k	ABLIKIM	25BR BES3	$e^+ e^-$ at 4.128–4.226 GeV

$\Gamma(\phi(1020)\pi^+, \phi \rightarrow K_L^0 K_S^0)/\Gamma(K_S^0 K_L^0 \pi^+)$ **Γ_{47}/Γ_{46}**

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
70.2±1.4±1.2	2.35k	ABLIKIM	25BR BES3	$e^+ e^-$ at 4.128–4.226 GeV

$\Gamma(K_L^0 K^{*(892)+}, K^{*+} \rightarrow K_S^0 \pi^+)/\Gamma(K_S^0 K_L^0 \pi^+)$ **Γ_{48}/Γ_{46}**

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
19.3±1.2±1.0	2.35k	ABLIKIM	25BR BES3	$e^+ e^-$ at 4.128–4.226 GeV

$\Gamma(K_S^0 K^{*(892)+}, K^{*+} \rightarrow K_L^0 \pi^+)/\Gamma(K_S^0 K_L^0 \pi^+)$ **Γ_{49}/Γ_{46}**

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
14.4±1.2±1.2	2.35k	ABLIKIM	25BR BES3	$e^+ e^-$ at 4.128–4.226 GeV

$\Gamma(\phi(1680)\pi^+, \phi \rightarrow K_L^0 K_S^0)/\Gamma(K_S^0 K_L^0 \pi^+)$ **Γ_{50}/Γ_{46}**

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
3.3±0.9±0.7	2.35k	ABLIKIM	25BR BES3	$e^+ e^-$ at 4.128–4.226 GeV

$\Gamma(2K_S^0 \text{ anything})/\Gamma_{\text{total}}$ **Γ_{16}/Γ**

VALUE (%)		DOCUMENT ID	TECN	COMMENT
1.7±0.3±0.1		DOBBS	09	CLEO $e^+ e^-$ at 4170 MeV

$\Gamma(2K^+ \text{ anything})/\Gamma_{\text{total}}$ **Γ_{17}/Γ**

VALUE (%)	CL%	DOCUMENT ID	TECN	COMMENT
<0.26	90	DOBBS	09	CLEO $e^+ e^-$ at 4170 MeV

$\Gamma(2K^- \text{ anything})/\Gamma_{\text{total}}$ **Γ_{18}/Γ**

VALUE (%)	CL%	DOCUMENT ID	TECN	COMMENT
<0.06	90	DOBBS	09	CLEO $e^+ e^-$ at 4170 MeV

$\Gamma(2\pi^+\pi^- + \text{anything})/\Gamma_{\text{total}}$ Γ_{19}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
32.81 ± 0.35 ± 0.63	25k	¹ ABLIKIM	23AV BES3	e^+e^- at 4.178 GeV

¹ Charged pions from K_S^0 meson decays are excluded from this measurement

———— Leptonic and semileptonic modes ————

See the related review(s):

[Leptonic Decays of Charged Pseudoscalar Mesons](#)

$\Gamma(e^+\nu_e)/\Gamma_{\text{total}}$ Γ_{20}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 0.83 × 10⁻⁴	90	¹ ZUPANC	13 BELL	e^+e^- at $\Upsilon(4S), \Upsilon(5S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
< 2.3 × 10 ⁻⁴	90	DEL-AMO-SA..10J	BABR	e^+e^- , 10.58 GeV
< 1.2 × 10 ⁻⁴	90	ALEXANDER 09	CLEO	e^+e^- at 4170 MeV
< 1.3 × 10 ⁻⁴	90	PEDLAR	07A CLEO	See ALEXANDER 09

¹ ZUPANC 13 also gives the limit as < 1.0 × 10⁻⁴ at 95% CL.

$\Gamma(\mu^+\nu_\mu)/\Gamma_{\text{total}}$ Γ_{21}/Γ

See the note on "Decay Constants of Charged Pseudoscalar Mesons."

VALUE (units 10 ⁻³)	EVTS	DOCUMENT ID	TECN	COMMENT
5.37 ± 0.11 OUR AVERAGE				
5.47 ± 0.26 ± 0.16	0.5k	ABLIKIM	24CG BES3	e^+e^- at 4.237–4.699 GeV
5.294 ± 0.108 ± 0.085	2.5k	ABLIKIM	23BR BES3	e^+e^- at 4.128–4.226 GeV
5.17 ± 0.75 ± 0.21	69	¹ ABLIKIM	160 BES3	e^+e^- at 4.009 GeV
5.31 ± 0.28 ± 0.20	490	² ZUPANC	13 BELL	e^+e^- at $\Upsilon(4S), \Upsilon(5S)$
6.02 ± 0.38 ± 0.34	270	³ DEL-AMO-SA..10J	BABR	e^+e^- , 10.58 GeV
5.65 ± 0.45 ± 0.17	230	ALEXANDER 09	CLEO	e^+e^- at 4.170 GeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
5.35 ± 0.13 ± 0.16	2.2k	⁴ ABLIKIM	21BE BES3	e^+e^- , 4.178, 4.226 GeV
5.49 ± 0.16 ± 0.15	1.1k	⁴ ABLIKIM	19E BES3	e^+e^- at 4.178 GeV
6.44 ± 0.76 ± 0.57	170	⁵ WIDHALM	08 BELL	See ZUPANC 13
5.94 ± 0.66 ± 0.31	88	⁶ PEDLAR	07A CLEO	See ALEXANDER 09
6.8 ± 1.1 ± 1.8	553	⁷ HEISTER	02I ALEP	Z decays

¹ ABLIKIM 160 also reports that when constrained by the Standard Model ratio of $\Gamma(D_s^+ \rightarrow \tau^+\nu_\tau)/\Gamma(D_s^+ \rightarrow \mu^+\nu_\mu) = 9.76$, the branching fraction is found to be $(0.495 \pm 0.067 \pm 0.026)\%$. The constrained value is used to obtain the decay constant, $f_{D_s^+} = (241.0 \pm 16.3 \pm 6.6)$ MeV.

² ZUPANC 13 uses both $\mu^+\nu$ and $\tau^+\nu$ events to get $f_{D_s} = (255.5 \pm 4.2 \pm 5.1)$ MeV.

³ DEL-AMO-SANCHEZ 10J uses $\mu^+\nu_\mu$ and $\tau^+\nu_\tau$ events together to get $f_{D_s} = (258.6 \pm 6.4 \pm 7.5)$ MeV.

⁴ Superseded by ABLIKIM 23BR.

⁵ WIDHALM 08 gets $f_{D_s} = (275 \pm 16 \pm 12)$ MeV from the branching fraction.

⁶ PEDLAR 07A also fits μ^+ and τ^+ events together and gets an effective $\mu^+\nu_\mu$ branching fraction of $(6.38 \pm 0.59 \pm 0.33) \times 10^{-3}$

⁷ This HEISTER 02I result is not actually an independent measurement of the absolute $\mu^+\nu_\mu$ branching fraction, but is in fact based on our $\phi\pi^+$ branching fraction of 3.6 ±

0.9%, so it cannot be included in our overall fit. HEISTER 02I combines its $D_s^+ \rightarrow \tau^+ \nu_\tau$ and $\mu^+ \nu_\mu$ branching fractions to get $f_{D_s} = (285 \pm 19 \pm 40)$ MeV.

$\Gamma(\mu^+ \nu_\mu)/\Gamma(\phi\pi^+)$

Γ_{21}/Γ_{53}

See the note on “Decay Constants of Charged Pseudoscalar Mesons” above.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.143 \pm 0.018 \pm 0.006$	489 ± 55	¹ AUBERT	07V BABR	$e^+ e^- \approx \Upsilon(4S)$
$0.23 \pm 0.06 \pm 0.04$	18	² ALEXANDROV	00 BEAT	π^- nucleus, 350 GeV
$0.173 \pm 0.023 \pm 0.035$	182	³ CHADHA	98 CLE2	$e^+ e^- \approx \Upsilon(4S)$
$0.245 \pm 0.052 \pm 0.074$	39	⁴ ACOSTA	94 CLE2	See CHADHA 98

¹ AUBERT 07V gets $f_{D_s^+} = (283 \pm 17 \pm 16)$ MeV, using $\Gamma(D_s^+ \rightarrow \phi\pi^+)/\Gamma(\text{total}) = (4.71 \pm 0.46)\%$.

² ALEXANDROV 00 uses $f_D^2/f_{D_s}^2 = 0.82 \pm 0.09$ from a lattice-gauge-theory calculation to get the relative numbers of $D^+ \rightarrow \mu^+ \nu_\mu$ and $D_s^+ \rightarrow \mu^+ \nu_\mu$ events. The present result leads to $f_{D_s} = (323 \pm 44 \pm 36)$ MeV.

³ CHADHA 98 obtains $f_{D_s} = (280 \pm 19 \pm 28 \pm 34)$ MeV from this measurement, using $\Gamma(D_s^+ \rightarrow \phi\pi^+)/\Gamma(\text{total}) = 0.036 \pm 0.009$.

⁴ ACOSTA 94 obtains $f_{D_s} = (344 \pm 37 \pm 52 \pm 42)$ MeV from this measurement, using $\Gamma(D_s^+ \rightarrow \phi\pi^+)/\Gamma(\text{total}) = 0.037 \pm 0.009$.

$\Gamma(\tau^+ \nu_\tau)/\Gamma_{\text{total}}$

Γ_{22}/Γ

See the note on “Decay Constants of Charged Pseudoscalar Mesons” above.

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

5.39 ± 0.09 OUR AVERAGE

$5.60 \pm 0.16 \pm 0.20$	2.8k	¹ ABLIKIM	24CG BES3	$e^+ e^-$ at 4.237–4.699 GeV
$5.44 \pm 0.17 \pm 0.13$	2.4k	² ABLIKIM	23BP BES3	$e^+ e^-$ at 4.128–4.226 GeV
$5.37 \pm 0.17 \pm 0.15$	2.3k	³ ABLIKIM	23BX BES3	$e^+ e^-$ at 4.128–4.226 GeV
$5.29 \pm 0.25 \pm 0.20$	1.7k	⁴ ABLIKIM	21AF BES3	$e^+ e^-$ at 4.178, 4.226 GeV
$5.27 \pm 0.10 \pm 0.12$	4.9k	⁵ ABLIKIM	21AZ BES3	$e^+ e^-$ at 4.178, 4.226 GeV
$3.28 \pm 1.83 \pm 0.37$	33	⁶ ABLIKIM	160 BES3	$e^+ e^-$ at 4.009 GeV
$5.70 \pm 0.21^{+0.31}_{-0.30}$	2.2k	⁷ ZUPANC	13 BELL	$e^+ e^-$ at $\Upsilon(4S)$, $\Upsilon(5S)$
$4.96 \pm 0.37 \pm 0.57$	748	⁸ DEL-AMO-SA..10J	BABR	$e^- \bar{\nu}_e \nu_\tau, \mu^- \bar{\nu}_\mu \nu_\tau$
$6.42 \pm 0.81 \pm 0.18$	126	⁹ ALEXANDER	09 CLEO	$\tau^+ \rightarrow \pi^+ \bar{\nu}_\tau$
$5.52 \pm 0.57 \pm 0.21$	155	⁹ NAIK	09A CLEO	$\tau^+ \rightarrow \rho^+ \bar{\nu}_\tau$
$5.30 \pm 0.47 \pm 0.22$	181	⁹ ONYISI	09 CLEO	$\tau^+ \rightarrow e^+ \nu_e \bar{\nu}_\tau$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$5.21 \pm 0.25 \pm 0.17$	950	¹⁰ ABLIKIM	21BE BES3	$e^+ e^-$ at 4.178, 4.226 GeV
$6.17 \pm 0.71 \pm 0.34$	102	¹¹ ECKLUND	08 CLEO	See ONYISI 09
$8.0 \pm 1.3 \pm 0.4$	47	¹¹ PEDLAR	07A CLEO	See ALEXANDER 09
$5.79 \pm 0.77 \pm 1.84$	881	¹² HEISTER	02I ALEP	Z decays
$7.0 \pm 2.1 \pm 2.0$	22	¹³ ABBIENDI	01L OPAL	$D_s^{*+} \rightarrow \gamma D_s^+$ from Z's
$7.4 \pm 2.8 \pm 2.4$	16	¹⁴ ACCIARRI	97F L3	$D_s^{*+} \rightarrow \gamma D_s^+$ from Z's

¹ ABLIKIM 24CG uses $\tau^+ \rightarrow (e^+ \nu_e, \mu^+ \nu_\mu, \pi^+, \rho^+) \bar{\nu}_\tau$ decays.

² ABLIKIM 23BP uses $\tau^+ \rightarrow \pi^+ \bar{\nu}_\tau$ decays.

- ³ ABLIKIM 23BX uses $\tau^+ \rightarrow \mu^+ \nu_\mu \bar{\nu}_\tau$ decays.
⁴ ABLIKIM 21AF uses $\tau^+ \rightarrow \pi^+ \pi^0 \bar{\nu}_\tau$ decays.
⁵ ABLIKIM 21AZ uses $\tau^+ \rightarrow e^+ \nu_e \bar{\nu}_\tau$ decays.
⁶ ABLIKIM 160 also reports that when constrained by the Standard Model ratio of $\Gamma(D_s^+ \rightarrow \tau^+ \nu_\tau)/\Gamma(D_s^+ \rightarrow \mu^+ \nu_\mu) = 9.76$; the branching fraction is found to be $(4.83 \pm 0.65 \pm 0.26)\%$.
⁷ ZUPANC 13 uses both $\mu^+ \nu$ and $\tau^+ \nu$ events to get $f_{D_s} = (255.5 \pm 4.2 \pm 5.1)$ MeV.
⁸ DEL-AMO-SANCHEZ 10J (with a small correction; see LEES 15D) uses $\mu^+ \nu_\mu$ and $\tau^+ \nu_\tau$ events together to get $f_{D_s} = (259.9 \pm 6.6 \pm 7.6)$ MeV.
⁹ ALEXANDER 09, NAIK 09A, and ONYISI 09 use different τ decay modes and are independent. The three papers combined give $f_{D_s} = (259.7 \pm 7.8 \pm 3.4)$ MeV.
¹⁰ ABLIKIM 21BE uses $\tau^+ \rightarrow \pi^+ \bar{\nu}_\tau$ decays. When constrained by the Standard Model ratio of $\Gamma(D_s^+ \rightarrow \tau^+ \nu_\tau)/\Gamma(D_s^+ \rightarrow \mu^+ \nu_\mu) = 9.75$, the branching fraction is found to be $(5.22 \pm 0.10 \pm 0.14)\%$. Superseded by ABLIKIM 23BP.
¹¹ ECKLUND 08 and PEDLAR 07A are independent: ECKLUND 08 uses $\tau^+ \rightarrow e^+ \nu_e \bar{\nu}_\tau$ events, PEDLAR 07A uses $\tau^+ \rightarrow \pi^+ \bar{\nu}_\tau$ events.
¹² HEISTER 02I combines its $D_s^+ \rightarrow \tau^+ \nu_\tau$ and $\mu^+ \nu_\mu$ branching fractions to get $f_{D_s} = (285 \pm 19 \pm 40)$ MeV.
¹³ This ABBIENDI 01L value gives a decay constant f_{D_s} of $(286 \pm 44 \pm 41)$ MeV.
¹⁴ The second ACCIARRI 97F error here combines in quadrature systematic (0.016) and normalization (0.018) errors. The branching fraction gives $f_{D_s} = (309 \pm 58 \pm 33 \pm 38)$ MeV.

$\Gamma(\tau^+ \nu_\tau)/\Gamma(\mu^+ \nu_\mu)$

Γ_{22}/Γ_{21}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$10.73 \pm 0.69^{+0.56}_{-0.53}$	2.2k/492	¹ ZUPANC	13	BELL $e^+ e^-$ at $\Upsilon(4S), \Upsilon(5S)$
$11.0 \pm 1.4 \pm 0.6$	102	² ECKLUND	08	CLEO See ONYISI 09

¹ This ZUPANC 13 ratio is not independent of the separate $\tau\nu$ and $\mu\nu$ fractions listed above.

² This ECKLUND 08 value also uses results from PEDLAR 07A, and it is not independent of other results in these Listings. Combined with earlier CLEO results, the decay constant f_{D_s} is $274 \pm 10 \pm 5$ MeV.

$\Gamma(\gamma e^+ \nu_e)/\Gamma_{\text{total}}$

Γ_{23}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 1.3 \times 10^{-4}$	90	ABLIKIM	19AD BES3	for $E_\gamma > 10$ MeV

$\Gamma(K^+ K^- e^+ \nu_e)/\Gamma(K^+ K^- \pi^+)$

Γ_{24}/Γ_{52}

VALUE	DOCUMENT ID	TECN	COMMENT
$0.558 \pm 0.007 \pm 0.016$	¹ AUBERT	08AN BABR	$e^+ e^-$ at $\Upsilon(4S)$

¹ This AUBERT 08AN ratio is only for the $K^+ K^-$ mass in the range 1.01–to–1.03 GeV in the numerator and 1.0095–to–1.0295 GeV in the denominator.

$\Gamma(K_S^0 K_S^0 e^+ \nu_e)/\Gamma_{\text{total}}$

Γ_{25}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 3.8 \times 10^{-4}$	90	ABLIKIM	22J BES3	$e^+ e^-$ at 4.178–4.226 GeV

$\Gamma(\phi e^+ \nu_e)/\Gamma_{\text{total}}$ Γ_{26}/Γ

See the end of the D_s^+ Listings for measurements of $D_s^+ \rightarrow \phi e^+ \nu_e$ form factors.
Unseen decay modes of the ϕ are included.

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
2.34±0.12 OUR AVERAGE		Error includes scale factor of 1.2.		
2.21±0.16±0.11	350	ABLIKIM	24CI BES3	$e^+ e^-$ at 4237–4699 MeV
2.26±0.45±0.09	26	ABLIKIM	18A BES3	$e^+ e^-$ at 4009 MeV
2.14±0.17±0.08	207	HIETALA	15	Uses CLEO data
2.61±0.03±0.17	25k	AUBERT	08AN BABR	$e^+ e^-$ at $\Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
2.36±0.23±0.13	106	ECKLUND	09 CLEO	See HIETALA 15
2.29±0.37±0.11	45	YELTON	09 CLEO	See ECKLUND 09

 $\Gamma(\phi e^+ \nu_e)/\Gamma(\phi \pi^+)$ Γ_{26}/Γ_{53}

As noted in the comment column, most of these measurements use $\phi \mu^+ \nu_\mu$ events in addition to or instead of $\phi e^+ \nu_e$ events.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
0.540±0.033±0.048	793	LINK	02J FOCS	Uses $\phi \mu^+ \nu_\mu$
0.54 ±0.05 ±0.04	367	BUTLER	94 CLE2	Uses $\phi e^+ \nu_e$ and $\phi \mu^+ \nu_\mu$
0.58 ±0.17 ±0.07	97	FRABETTI	93G E687	Uses $\phi \mu^+ \nu_\mu$
0.57 ±0.15 ±0.15	104	ALBRECHT	91 ARG	Uses $\phi e^+ \nu_e$
0.49 ±0.10 ^{+0.10} _{-0.14}	54	ALEXANDER	90B CLEO	Uses $\phi e^+ \nu_e$ and $\phi \mu^+ \nu_\mu$

 $\Gamma(\phi \mu^+ \nu_\mu)/\Gamma_{\text{total}}$ Γ_{29}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
2.24±0.11 OUR AVERAGE				
2.25±0.09±0.07	1.7k	ABLIKIM	23BZ BES3	$e^+ e^-$ at 4.128–4.226 GeV
1.94±0.53±0.09	22	ABLIKIM	18A BES3	$e^+ e^-$ at 4.009 GeV

 $\Gamma(\eta e^+ \nu_e)/\Gamma_{\text{total}}$ Γ_{30}/Γ

Unseen decay modes of the η are included.

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
2.27 ±0.06 OUR AVERAGE				
2.35 ±0.11 ±0.10	716	ABLIKIM	24CI BES3	$e^+ e^-$ at 4237–4699 MeV
2.255±0.039±0.051	4k	ABLIKIM	23BO BES3	$e^+ e^-$ at 4128–4226 MeV
2.30 ±0.31 ±0.08	63	ABLIKIM	16T BES3	$e^+ e^-$ at 4009 MeV
2.28 ±0.14 ±0.19	358	¹ HIETALA	15	Uses CLEO data
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
2.323±0.063±0.063	1.8k	² ABLIKIM	19S BES3	$e^+ e^-$ at 4178 MeV
2.48 ±0.29 ±0.13	82	YELTON	09 CLEO	See HIETALA 15

¹ Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.

² Superseded by ABLIKIM 23BO

 $\Gamma(\eta e^+ \nu_e)/\Gamma(\phi e^+ \nu_e)$ Γ_{30}/Γ_{26}

Unseen decay modes of the η and the ϕ are included.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1.24±0.12±0.15	440	¹ BRANDENB...	95 CLE2	See HIETALA 15

¹ BRANDENBURG 95 uses both e^+ and μ^+ events and makes a phase-space adjustment to use the μ^+ events as e^+ events.

$\Gamma(\eta'(958)e^+\nu_e)/\Gamma_{\text{total}}$ **Γ_{31}/Γ**

Unseen decay modes of the $\eta'(958)$ are included.

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.81 ± 0.04	OUR AVERAGE			
0.82 ± 0.09 ± 0.04	134	ABLIKIM 24CI	BES3	e^+e^- at 4237–4699 MeV
0.810 ± 0.038 ± 0.024	675	ABLIKIM 23BO	BES3	e^+e^- at 4128–4226 MeV
0.93 ± 0.30 ± 0.05	14	ABLIKIM 16T	BES3	e^+e^- at 4009 MeV
0.68 ± 0.15 ± 0.06	20	¹ HIETALA 15		Uses CLEO data
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.824 ± 0.073 ± 0.027	261	² ABLIKIM 19S	BES3	e^+e^- at 4178 MeV
0.91 ± 0.33 ± 0.05	7.5	YELTON 09	CLEO	See HIETALA 15

¹ Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.

² Superseded by ABLIKIM 23BO

$\Gamma(\eta'(958)e^+\nu_e)/\Gamma(\phi e^+\nu_e)$ **Γ_{31}/Γ_{26}**

Unseen decay modes of the resonances are included.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.43 ± 0.11 ± 0.07	29	¹ BRANDENB... 95	CLE2	See HIETALA 15

¹ BRANDENBURG 95 uses both e^+ and μ^+ events and makes a phase-space adjustment to use the μ^+ events as e^+ events.

$\Gamma(\eta\mu^+\nu_\mu)/\Gamma_{\text{total}}$ **Γ_{32}/Γ**

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
2.24 ± 0.07	OUR AVERAGE			
2.235 ± 0.051 ± 0.052	3.1k	ABLIKIM 24AQ	BES3	e^+e^- , 4.128–4.266 GeV
2.42 ± 0.46 ± 0.11	44	ABLIKIM 18A	BES3	e^+e^- at 4.009 GeV

$\Gamma(\eta'(958)\mu^+\nu_\mu)/\Gamma_{\text{total}}$ **Γ_{33}/Γ**

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.80 ± 0.06	OUR AVERAGE			
0.801 ± 0.055 ± 0.028	390	ABLIKIM 24AQ	BES3	e^+e^- , 4.128–4.266 GeV
1.06 ± 0.54 ± 0.07	10	ABLIKIM 18A	BES3	e^+e^- at 4.009 GeV

$\Gamma(\omega e^+\nu_e)/\Gamma_{\text{total}}$ **Γ_{34}/Γ**

A test for $u\bar{u}$ or $d\bar{d}$ content in the D_s^+ . Neither Cabibbo-favored nor Cabibbo-suppressed decays can contribute, and $\omega - \phi$ mixing is an unlikely explanation for any fraction above about 2×10^{-4} .

VALUE (%)	CL%	DOCUMENT ID	TECN	COMMENT
<0.20	90	MARTIN 11	CLEO	e^+e^- at 4170 MeV

$\Gamma(K^0 e^+\nu_e)/\Gamma_{\text{total}}$ **Γ_{35}/Γ**

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.288 ± 0.026	OUR AVERAGE	Error includes scale factor of 1.2.		
0.298 ± 0.023 ± 0.012	228	¹ ABLIKIM 24CH	BES3	e^+e^- at 4128–4226 MeV
0.24 ± 0.04 ± 0.01	51	ABLIKIM 24CI	BES3	e^+e^- at 4237–4699 MeV
0.39 ± 0.08 ± 0.03	42	HIETALA 15		Uses CLEO data

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.325 ± 0.038 ± 0.016 117 ^{1,2} ABLIKIM 19D BES3 e^+e^- at 4178 MeV

0.37 ± 0.10 ± 0.02 14 YELTON 09 CLEO See HIETALA 15

¹ K^0 reconstructed via $K^0 \rightarrow K_S^0 \rightarrow \pi^+\pi^-$ decays.

² Superseded by ABLIKIM 24CH.

$\Gamma(K_1(1270)^0 e^+ \nu_e) / \Gamma_{\text{total}}$ Γ_{27}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 4.1 \times 10^{-4}$	90	¹ ABLIKIM 23BS	BES3	e^+e^- at 4.128–4.226 GeV

¹ ABLIKIM 23BS uses $K_1(1270)^0 \rightarrow K^- \pi^+ \pi^0$ decays.

$\Gamma(K^*(892)^0 e^+ \nu_e) / \Gamma_{\text{total}}$ Γ_{36}/Γ

Unseen decay modes of the $K^*(892)^0$ are included.

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.205 ± 0.020 OUR AVERAGE				

0.19 ± 0.03 ± 0.01 65 ABLIKIM 24CI BES3 e^+e^- at 4237–4699 MeV

0.237 ± 0.026 ± 0.020 155 ABLIKIM 19D BES3 e^+e^- at 4178 MeV

0.18 ± 0.04 ± 0.01 32 ¹ HIETALA 15 e^+e^- at 4170 MeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.18 ± 0.07 ± 0.01 7.5 YELTON 09 CLEO See HIETALA 15

¹ Uses CLEO data, but not authored by the CLEO collaboration

$\Gamma(f_0(500) e^+ \nu_e, f_0 \rightarrow \pi^0 \pi^0) / \Gamma_{\text{total}}$ Γ_{37}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 7.3 \times 10^{-4}$	90	ABLIKIM 22J	BES3	e^+e^- at 4.178–4.226 GeV

$\Gamma(f_0(500) e^+ \nu_e, f_0 \rightarrow \pi^+ \pi^-) / \Gamma_{\text{total}}$ Γ_{38}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 3.3 \times 10^{-4}$	90	ABLIKIM 24AT	BES3	e^+e^- at 4.128–4.226 GeV

$\Gamma(f_0(980) e^+ \nu_e, f_0 \rightarrow \pi^0 \pi^0) / \Gamma_{\text{total}}$ Γ_{39}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
7.9 ± 1.4 ± 0.4	55	¹ ABLIKIM 22J	BES3	e^+e^- at 4.178–4.226 GeV

¹ Assuming $B(f_0 \rightarrow \pi^0 \pi^0) = 1/3$ via the isospin limit, this result implies $B(D_s^+ \rightarrow f_0(980) e^+ \nu_e) = (2.4 \pm 0.4) \times 10^{-3}$.

$\Gamma(f_0(980) e^+ \nu_e, f_0 \rightarrow \pi^+ \pi^-) / \Gamma_{\text{total}}$ Γ_{40}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.64 ± 0.13 OUR AVERAGE				

1.72 ± 0.13 ± 0.10 0.4k ABLIKIM 24AT BES3 e^+e^- at 4128–4226 MeV

1.5 ± 0.2 ± 0.1 91 ABLIKIM 24CI BES3 e^+e^- at 4237–4699 MeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.3 ± 0.3 ± 0.1 42 ¹ HIETALA 15 Uses CLEO data

2.0 ± 0.3 ± 0.1 44 ECKLUND 09 CLEO See HIETALA 15

1.3 ± 0.4 ± 0.1 13 YELTON 09 CLEO See ECKLUND 09

¹ HIETALA 15 uses a tighter cut on the reconstructed $\pi^+\pi^-$ mass (± 60 MeV around the f^0) than ECKLUND 09. It finds that applying the same tight cut to both analyses gives consistent results.

$\Gamma(f_0(980)\mu^+\nu_\mu, f_0 \rightarrow K^+K^-)/\Gamma_{\text{total}}$ Γ_{41}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5.45 \times 10^{-4}$	90	¹ ABLIKIM	23BZ BES3	e^+e^- at 4.128–4.226 GeV

¹ Partial wave analysis of 939 $D_s^+ \rightarrow K^+K^-\mu^+\nu_\mu$ events, assuming K^+K^- S-wave is 100% $f_0(980)$.

 $\Gamma(a_0(980)^0e^+\nu_e, a_0^0 \rightarrow \pi^0\eta)/\Gamma_{\text{total}}$ Γ_{42}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.2 \times 10^{-4}$	90	ABLIKIM	21Y BES3	e^+e^- at 4.178–4.226 GeV

 $\Gamma(\pi^0e^+\nu_e)/\Gamma_{\text{total}}$ Γ_{43}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<6.4 \times 10^{-5}$	90	ABLIKIM	22BH BES3	6.32 fb^{-1} of e^+e^- at 4.178–4.226 GeV

 $\Gamma(b_1(1235)^0e^+\nu_e, b_1^0 \rightarrow \omega\pi^0)/\Gamma_{\text{total}}$ Γ_{28}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<6.4 \times 10^{-4}$	90	ABLIKIM	23BS BES3	e^+e^- at 4.128–4.226 GeV

————— Hadronic modes with a $K\bar{K}$ pair ————— $\Gamma(K^+K_S^0)/\Gamma_{\text{total}}$ Γ_{44}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

1.500±0.014 OUR FIT**1.503±0.015 OUR AVERAGE**

$1.502 \pm 0.012 \pm 0.009$		ABLIKIM	24BD BES3	e^+e^- , 4.128–4.226 GeV
$1.52 \pm 0.05 \pm 0.03$		ONYISI	13 CLEO	e^+e^- at 4.17 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.425 \pm 0.038 \pm 0.031$	1.8k	¹ ABLIKIM	19AMBES3	e^+e^- at 4178 MeV
$1.49 \pm 0.07 \pm 0.05$		² ALEXANDER	08 CLEO	See ONYISI 13

¹ Superseded by ABLIKIM 24BD.

² ALEXANDER 08 uses single- and double-tagged events in an overall fit.

 $\Gamma(K^+K_S^0)/\Gamma(K^+K^-\pi^+)$ Γ_{44}/Γ_{52}

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
$27.55 \pm 0.18 \pm 0.50$	40k	ABLIKIM	20R BES3	e^+e^- , 4178 ~ 4226 MeV

 $\Gamma(K^+K_L^0)/\Gamma_{\text{total}}$ Γ_{45}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$1.485 \pm 0.039 \pm 0.046$	2.3k	ABLIKIM	19AMBES3	e^+e^- at 4178 MeV

 $\Gamma(K^+\bar{K}^0)/\Gamma_{\text{total}}$ Γ_{51}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$2.95 \pm 0.11 \pm 0.09$	2.0k	¹ ZUPANC	13 BELL	e^+e^- at $\Upsilon(4S), \Upsilon(5S)$

¹ ZUPANC 13 finds the \bar{K}^0 from its missing-mass squared, not from $K_S^0 \rightarrow \pi^+\pi^-$.

The DCS ($D_s^+ \rightarrow K^+K^0$) contribution to this fraction is estimated to be an order of magnitude below the statistical uncertainty.

$\Gamma(K^+ K^- \pi^+)/\Gamma_{\text{total}}$ Γ_{52}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
5.45±0.08 OUR FIT	Error includes scale factor of 1.3.			
5.48±0.08 OUR AVERAGE	Error includes scale factor of 1.1.			
5.49±0.04±0.07		ABLIKIM	24BD BES3	$e^+ e^-$, 4.128–4.226 GeV
5.55±0.14±0.13		ONYISI	13 CLEO	$e^+ e^-$ at 4.17 GeV
5.06±0.15±0.21	4.1k	ZUPANC	13 BELL	$e^+ e^-$ at $\Upsilon(4S), \Upsilon(5S)$
5.78±0.20±0.30		DEL-AMO-SA..10J	BABR	$e^+ e^-$, 10.58 GeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
5.47±0.08±0.13	5.1k	¹ ABLIKIM	21AE BES3	$e^+ e^-$ at 4.178 GeV
5.50±0.23±0.16		² ALEXANDER	08 CLEO	See ONYISI 13

¹ Superseded by ABLIKIM 24BD.

² ALEXANDER 08 uses single- and double-tagged events in an overall fit.

$\Gamma(\phi\pi^+)/\Gamma_{\text{total}}$ Γ_{53}/Γ

The results here are model-independent. For earlier, model-dependent results, see our PDG 06 edition. We decouple the $D_s^+ \rightarrow \phi\pi^+$ branching fraction obtained from mass projections (and used to get some of the other branching fractions) from the $D_s^+ \rightarrow \phi\pi^+, \phi \rightarrow K^+ K^-$ branching fraction obtained from the Dalitz-plot analysis of $D_s^+ \rightarrow K^+ K^- \pi^+$. That is, the ratio of these two branching fractions is not exactly the $\phi \rightarrow K^+ K^-$ branching fraction 0.491.

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
4.5 ±0.4 OUR AVERAGE				
4.62±0.36±0.51		¹ AUBERT	06N BABR	$e^+ e^-$ at $\Upsilon(4S)$
4.81±0.52±0.38	212 ± 19	² AUBERT	05V BABR	$e^+ e^- \approx \Upsilon(4S)$
3.59±0.77±0.48		³ ARTUSO	96 CLE2	$e^+ e^-$ at $\Upsilon(4S)$

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

3.9 ^{+5.1} _{-1.9} ^{+1.8} _{-1.1}		⁴ BAI	95C BES	$e^+ e^-$ 4.03 GeV
---	--	------------------	---------	--------------------

¹ This AUBERT 06N measurement uses $\bar{B}^0 \rightarrow D_s^{(*)-} D^{(*)+}$ and $B^- \rightarrow D_s^{(*)-} D^{(*)0}$ decays, including some from other papers. However, the result is independent of AUBERT 05V.

² AUBERT 05V uses the ratio of $B^0 \rightarrow D^{*-} D_s^{*+}$ events seen in two different ways, in both of which the $D^{*-} \rightarrow \bar{D}^0 \pi^-$ decay is fully reconstructed: (1) The $D_s^{*+} \rightarrow D_s^+ \gamma$, $D_s^+ \rightarrow \phi\pi^+$ decay is fully reconstructed. (2) The number of events in the D_s^+ peak in the missing mass spectrum against the $D^{*-} \gamma$ is measured.

³ ARTUSO 96 uses partially reconstructed $\bar{B}^0 \rightarrow D^{*+} D_s^{*-}$ decays to get a model-independent value for $\Gamma(D_s^- \rightarrow \phi\pi^-)/\Gamma(D^0 \rightarrow K^- \pi^+)$ of $0.92 \pm 0.20 \pm 0.11$.

⁴ BAI 95C uses $e^+ e^- \rightarrow D_s^+ D_s^-$ events in which one or both of the D_s^\pm are observed to obtain the first model-independent measurement of the $D_s^+ \rightarrow \phi\pi^+$ branching fraction, without assumptions about $\sigma(D_s^\pm)$. However, with only two “doubly-tagged” events, the statistical error is very large.

$\Gamma(\phi\pi^+, \phi \rightarrow K^+K^-)/\Gamma(K^+K^-\pi^+)$ Γ_{54}/Γ_{52}

This is the “fit fraction” from the Dalitz-plot analysis. We decouple the $D_s^+ \rightarrow \phi\pi^+$ branching fraction obtained from mass projections (and used to get some of the other branching fractions) from the $D_s^+ \rightarrow \phi\pi^+, \phi \rightarrow K^+K^-$ branching fraction obtained from the Dalitz-plot analysis of $D_s^+ \rightarrow K^+K^-\pi^+$. That is, the ratio of these two branching fractions is not exactly the $\phi \rightarrow K^+K^-$ branching fraction 0.491.

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
41.2±0.7 OUR AVERAGE				
40.5±0.7±0.9	18.6k	ABLIKIM	21AE BES3	e^+e^- at 4.178 GeV
41.4±0.8±0.5		DEL-AMO-SA..11G	BABR	Dalitz fit, 96k evts
42.2±1.6±0.3		MITCHELL	09A CLEO	Dalitz fit, 12k evts
• • • We do not use the following data for averages, fits, limits, etc. • • •				
39.6±3.3±4.7		FRABETTI	95B E687	Dalitz fit, 701 evts

 $\Gamma(K^+\bar{K}^*(892)^0)/\Gamma(K^*(892)^+\bar{K}^0)$ Γ_{55}/Γ_{72}

VALUE	DOCUMENT ID	TECN	COMMENT
2.35^{+0.42}_{-0.23}±0.10	ABLIKIM	22AH BES3	Dalitz plot fit to 990 $D_s^\pm \rightarrow K^\pm K_S^0 \pi^0$ evts

 $\Gamma(K^+\bar{K}^*(892)^0, \bar{K}^{*0} \rightarrow K^-\pi^+)/\Gamma(K^+K^-\pi^+)$ Γ_{56}/Γ_{52}

This is the “fit fraction” from the Dalitz-plot analysis.

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
47.9±0.6 OUR AVERAGE				
48.3±0.9±0.6	18.6k	ABLIKIM	21AE BES3	e^+e^- at 4.178 GeV
47.9±0.5±0.5		DEL-AMO-SA..11G	BABR	Dalitz fit, 96k evts
47.4±1.5±0.4		MITCHELL	09A CLEO	Dalitz fit, 12k evts
• • • We do not use the following data for averages, fits, limits, etc. • • •				
47.8±4.6±4.0		FRABETTI	95B E687	Dalitz fit, 701 evts


 $\Gamma(K^+\bar{K}^*(892)^0, \bar{K}^{*0} \rightarrow K_S^0\pi^0)/\Gamma(K^+K_S^0\pi^0)$ Γ_{57}/Γ_{65}

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
32.7±2.2±1.9	ABLIKIM	22AH BES3	Dalitz plot fit, 990 evts

 $\Gamma(f_0(980)\pi^+, f_0 \rightarrow K^+K^-)/\Gamma(K^+K^-\pi^+)$ Γ_{58}/Γ_{52}

This is the “fit fraction” from the Dalitz-plot analysis. This is likely a superposition of $D_s^+ \rightarrow f_0(980)\pi$ and $D_s^+ \rightarrow a_0(980)\pi$ which are indistinguishable in such an analysis.

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
20.6±3.5 OUR AVERAGE Error includes scale factor of 2.5. See the ideogram below.				
19.3±1.7±2.0	18.6k	ABLIKIM	21AE BES3	e^+e^- at 4.178 GeV
16.4±0.7±2.0		DEL-AMO-SA..11G	BABR	Dalitz fit, 96k evts
28.2±1.9±1.8		MITCHELL	09A CLEO	Dalitz fit, 12k evts
• • • We do not use the following data for averages, fits, limits, etc. • • •				
11.0±3.5±2.6		FRABETTI	95B E687	Dalitz fit, 701 evts




$\Gamma(f_0(1370)\pi^+, f_0 \rightarrow K^+K^-) / \Gamma(K^+K^-\pi^+)$

$\Gamma_{59} / \Gamma_{52}$

This is the "fit fraction" from the Dalitz-plot analysis.

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
1.3 ± 0.5 OUR AVERAGE	Error includes scale factor of 2.8. See the ideogram below.			
$1.2 \pm 0.4 \pm 0.2$	18.6k	ABLIKIM	21AE BES3	e^+e^- at 4.178 GeV
$1.1 \pm 0.1 \pm 0.2$		DEL-AMO-SA..11G	BABR	Dalitz fit, 96k evts
$4.3 \pm 0.6 \pm 0.5$		MITCHELL	09A CLEO	Dalitz fit, 12k evts



$\Gamma(f_0(1710)\pi^+, f_0 \rightarrow K^+K^-)/\Gamma(K^+K^-\pi^+)$ Γ_{60}/Γ_{52}

This is the “fit fraction” from the Dalitz-plot analysis. This is likely a superposition of $D_s^+ \rightarrow f_0(1710)\pi$ and $D_s^+ \rightarrow a_0(1710)\pi$ which are indistinguishable in such an analysis.

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.3±0.5 OUR AVERAGE				Error includes scale factor of 3.8.
1.9±0.4±0.6	18.6k	ABLIKIM	21AE BES3	e^+e^- at 4.178 GeV
1.1±0.1±0.1		DEL-AMO-SA..11G	BABR	Dalitz fit, 96k evts
3.4±0.5±0.3		MITCHELL	09A CLEO	Dalitz fit, 12k evts
• • • We do not use the following data for averages, fits, limits, etc. • • •				
3.4±2.3±3.5		FRABETTI	95B E687	Dalitz fit, 701 evts

$\Gamma(a_0(980)^+\pi^0, a_0^+ \rightarrow K^+K_S^0)/\Gamma(K^+K_S^0\pi^0)$ Γ_{61}/Γ_{65}

<u>VALUE (units 10^{-2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
7.7±1.7±1.8	ABLIKIM	22AH BES3	Dalitz plot fit, 990 evts

$\Gamma(a_0(1710)^+\pi^0, a_0^+ \rightarrow K^+K_S^0)/\Gamma(K^+K_S^0\pi^0)$ Γ_{62}/Γ_{65}

<u>VALUE (units 10^{-2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
23.6±3.4±2.0	¹ ABLIKIM	22AH BES3	Dalitz plot fit, 990 evts

¹ ABLIKIM 22AH observe an a_0 -like state with mass $m_{a_0} = 1.817 \pm 0.008 \pm 0.020$ GeV, and name the intermediate resonance $a_0(1817)$. We interpret this as the $a_0(1710)$ observed by LEES 21A.

$\Gamma(K^+\bar{K}_0^*(1430)^0, \bar{K}_0^* \rightarrow K^-\pi^+)/\Gamma(K^+K^-\pi^+)$ Γ_{63}/Γ_{52}

This is the “fit fraction” from the Dalitz-plot analysis.

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
3.3±0.5 OUR AVERAGE				
3.0±0.6±0.5	18.6k	ABLIKIM	21AE BES3	e^+e^- at 4.178 GeV
2.4±0.3±1.0		DEL-AMO-SA..11G	BABR	Dalitz fit, 96k evts
3.9±0.5±0.5		MITCHELL	09A CLEO	Dalitz fit, 12k evts
• • • We do not use the following data for averages, fits, limits, etc. • • •				
9.3±3.2±3.2		FRABETTI	95B E687	Dalitz fit, 701 evts

$\Gamma(K^+\bar{K}^*(1410)^0, \bar{K}_0^* \rightarrow K_S^0\pi^0)/\Gamma(K^+K_S^0\pi^0)$ Γ_{64}/Γ_{65}

<u>VALUE (units 10^{-2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
6.0±1.4±1.3	ABLIKIM	22AH BES3	Dalitz plot fit, 990 evts

$\Gamma(K^+K_S^0\pi^0)/\Gamma_{\text{total}}$ Γ_{65}/Γ

<u>VALUE (%)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.471±0.028 OUR AVERAGE				
1.47 ±0.02 ±0.02		ABLIKIM	24BD BES3	e^+e^- , 4.128–4.226 GeV
1.52 ±0.09 ±0.20		ONYISI	13 CLEO	e^+e^- at 4.17 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.46 ±0.06 ±0.05	990	¹ ABLIKIM	22AH BES3	e^+e^- at 4.178-4.226 GeV

¹ Superseded by ABLIKIM 24BD.

$\Gamma(K^*(892)^+K_S^0, K^{*+} \rightarrow K^+\pi^0)/\Gamma(K^+K_S^0\pi^0)$ Γ_{66}/Γ_{65}

<u>VALUE (units 10^{-2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
13.9±1.7±1.3	ABLIKIM	22AH BES3	Dalitz plot fit, 990 evts

$\Gamma(2K_S^0\pi^+)/\Gamma_{\text{total}}$ Γ_{67}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

0.732±0.014 OUR AVERAGE

0.73 ±0.01 ±0.01		ABLIKIM	24BD BES3	e^+e^- , 4.128–4.226 GeV
------------------	--	---------	-----------	----------------------------

0.77 ±0.05 ±0.03		ONYISI	13 CLEO	e^+e^- at 4.17 GeV
------------------	--	--------	---------	----------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.68 ±0.04 ±0.01	370	¹ ABLIKIM	22F BES3	e^+e^- , 4.178–4.226 GeV
------------------	-----	----------------------	----------	----------------------------

¹Superseded by ABLIKIM 24BD.

$\Gamma(f_0(980)\pi^+, f_0 \rightarrow K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{68}/Γ

This is the “fit fraction” from the Dalitz-plot analysis. This is likely a superposition of $D_s^+ \rightarrow f_0(980)\pi$ and $D_s^+ \rightarrow a_0(980)\pi$ which are indistinguishable in such an analysis.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<1.8 × 10⁻⁴	90	¹ ABLIKIM	22F BES3	Dalitz plot fit
----------------------------------	----	----------------------	----------	-----------------

¹Based on isospin considerations, the authors interpret the suppression in the observed rate of this mode compared to $D_s^+ \rightarrow f_0(980)\pi^+$, $f_0 \rightarrow K^+K^-$ as likely due to the destructive interference between $a_0(980)$ and $f_0(980)$ in decays to $K_S^0 K_S^0$.

$\Gamma(f_0(1710)\pi^+, f_0 \rightarrow K_S^0 K_S^0)/\Gamma(2K_S^0\pi^+)$ Γ_{69}/Γ_{67}

This is the “fit fraction” from the Dalitz-plot analysis. This is likely a superposition of $D_s^+ \rightarrow f_0(1710)\pi$ and $D_s^+ \rightarrow a_0(1700)\pi$ which are indistinguishable in such an analysis.

VALUE (units 10 ⁻²)	DOCUMENT ID	TECN	COMMENT
---------------------------------	-------------	------	---------

46.3±4.0±1.2	ABLIKIM	22F BES3	Dalitz plot fit, 400 evts
---------------------	---------	----------	---------------------------

$\Gamma(K^*(892)^+ K_S^0, K^{*+} \rightarrow K_S^0\pi^+)/\Gamma(2K_S^0\pi^+)$ Γ_{70}/Γ_{67}

VALUE (units 10 ⁻²)	DOCUMENT ID	TECN	COMMENT
---------------------------------	-------------	------	---------

43.5±3.9±0.5	ABLIKIM	22F BES3	Dalitz plot fit, 400 evts
---------------------	---------	----------	---------------------------

$\Gamma(K^*(892)^+ \bar{K}^0)/\Gamma(\phi\pi^+)$ Γ_{72}/Γ_{53}

Unseen decay modes of the resonances are included.

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

1.20±0.21±0.13	CHEN	89 CLEO	e^+e^- 10 GeV
-----------------------	------	---------	-----------------

$\Gamma(\phi\rho^+)/\Gamma_{\text{total}}$ Γ_{73}/Γ

VALUE (units 10 ⁻²)	EVTS	DOCUMENT ID	TECN	COMMENT
---------------------------------	------	-------------	------	---------

4.9 ±0.8 OUR AVERAGE Error includes scale factor of 3.0.

4.0 ±0.4 ±0.1	2k	^{1,2} ABLIKIM	25F BES3	e^+e^- at 4.128–4.226 GeV, $\phi \rightarrow \pi^+\pi^-\pi^0$
---------------	----	------------------------	----------	--

5.59±0.15±0.30	3k	³ ABLIKIM	21U BES3	e^+e^- at 4.178–4.226 GeV, $\phi \rightarrow K^+K^-$
----------------	----	----------------------	----------	---

¹See $\Gamma(D_s^+ \rightarrow \phi\rho^+, \phi \rightarrow \pi^+\pi^-\pi^0)/\Gamma(D_s^+ \rightarrow 2\pi^+\pi^-2\pi^0)$, $\Gamma_{132}/\Gamma_{130}$. ABLIKIM 25F reports $(3.98 \pm 0.33^{+0.21}_{-0.19}) \times 10^{-2}$ from a measurement of $[\Gamma(D_s^+ \rightarrow \phi\rho^+)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow \rho\pi + \pi^+\pi^-\pi^0)]$ assuming $B(\phi(1020) \rightarrow \rho\pi + \pi^+\pi^-\pi^0) = (14.9 \pm 0.4) \times 10^{-2}$, which we rescale to our best (shown rounded) value $B(\phi(1020) \rightarrow \rho\pi + \pi^+\pi^-\pi^0) = (14.7 \pm 0.4) \times 10^{-2}$. Our first error is their experiment’s error and our second error is the systematic error from using our best (shown rounded) value.

² Amplitude analysis of $D_s^+ \rightarrow 2\pi^+ \pi^- 2\pi^0$ with 14 components.

³ ABLIKIM 21U uses an amplitude analysis of $D_s^+ \rightarrow K^+ K^- \pi^+ \pi^0$ with 9 components.

$\Gamma(\phi\rho^+)/\Gamma(\phi\pi^+)$					Γ_{73}/Γ_{53}
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
$1.86 \pm 0.26^{+0.29}_{-0.40}$	253	EVERY	92	CLE2	$e^+e^- \simeq 10.5$ GeV

$\Gamma(K^+ K^- \pi^+ \pi^0)/\Gamma_{\text{total}}$					Γ_{74}/Γ
VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT	
5.53 ± 0.15 OUR FIT	Error includes scale factor of 1.3.				
5.53 ± 0.12 OUR AVERAGE					

5.50 ± 0.05 ± 0.11 ABLIKIM 24BD BES3 e^+e^- , 4.128–4.226 GeV

6.37 ± 0.21 ± 0.56 ONYISI 13 CLEO e^+e^- at 4.17 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

5.42 ± 0.10 ± 0.17 3k ¹ ABLIKIM 21U BES3 e^+e^- , 4.178–4.226 GeV

5.65 ± 0.29 ± 0.40 ² ALEXANDER 08 CLEO See ONYISI 13

¹ ABLIKIM 21U uses an amplitude analysis of $D_s^+ \rightarrow K^+ K^- \pi^+ \pi^0$ with 9 components.
Superseded by ABLIKIM 24BD.

² ALEXANDER 08 uses single- and double-tagged events in an overall fit.

$\Gamma(\bar{K}_1(1270)^0 K^+, \bar{K}_1(1270)^0 \rightarrow K^- \rho^+)/\Gamma_{\text{total}}$					Γ_{75}/Γ
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	
$0.57 \pm 0.05 \pm 0.04$	3k	¹ ABLIKIM	21U	BES3	e^+e^- at 4.178–4.226 GeV

¹ ABLIKIM 21U uses an amplitude analysis of $D_s^+ \rightarrow K^+ K^- \pi^+ \pi^0$ with 9 components.

$\Gamma(\bar{K}_1(1270)^0 K^+, \bar{K}_1(1270)^0 \rightarrow K^*(892)\pi)/\Gamma_{\text{total}}$					Γ_{76}/Γ
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	
$1.31 \pm 0.18 \pm 0.18$	3k	^{1,2} ABLIKIM	21U	BES3	e^+e^- at 4.178–4.226 GeV

¹ ABLIKIM 21U uses an amplitude analysis of $D_s^+ \rightarrow K^+ K^- \pi^+ \pi^0$ with 9 components.

² $\bar{K}_1(1270)^0 \rightarrow K^*(892)\pi$ denotes a sum over $\bar{K}(892)^0 \pi^0$ and $K(892)^- \pi^+$ final states, which are assumed to have relative branching ratio 1/2, as per isospin.

$\Gamma(\bar{K}_1(1400)^0 K^+, \bar{K}_1(1400)^0 \rightarrow K^*(892)\pi)/\Gamma_{\text{total}}$					Γ_{77}/Γ
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	
$1.98 \pm 0.27 \pm 0.32$	3k	¹ ABLIKIM	21U	BES3	e^+e^- at 4.178–4.226 GeV

¹ $\bar{K}_1(1400)^0 \rightarrow K^*(892)\pi$ denotes a sum over $\bar{K}(892)^0 \pi^0$ and $K(892)^- \pi^+$ final states, which are assumed to have relative branching ratio 1/2, as per isospin.

$\Gamma(a_0(980)^0 \rho^+, a_0^0 \rightarrow K^+ K^-)/\Gamma_{\text{total}}$					Γ_{78}/Γ
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	
$0.19 \pm 0.03 \pm 0.03$	3k	¹ ABLIKIM	21U	BES3	e^+e^- at 4.178–4.226 GeV

¹ ABLIKIM 21U uses an amplitude analysis of $D_s^+ \rightarrow K^+ K^- \pi^+ \pi^0$ with 9 components.

$\Gamma(f_1(1420)^0 \pi^+, f_1(1420)^0 \rightarrow K^*(892)^\mp K^\pm)/\Gamma_{\text{total}}$					Γ_{79}/Γ
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	
$0.39 \pm 0.06 \pm 0.03$	3k	¹ ABLIKIM	21U	BES3	e^+e^- at 4.178–4.226 GeV

¹ ABLIKIM 21U uses an amplitude analysis of $D_s^+ \rightarrow K^+ K^- \pi^+ \pi^0$ with 9 components.

$\Gamma(f_1(1420)^0 \pi^+, f_1^0 \rightarrow a_0(980)^0 \pi^0, a_0^0 \rightarrow K^+ K^-) / \Gamma_{\text{total}}$ Γ_{80} / Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
0.04±0.01±0.01	3k	¹ ABLIKIM	21U BES3	$e^+ e^-$ at 4.178–4.226 GeV
¹ ABLIKIM 21U uses an amplitude analysis of $D_s^+ \rightarrow K^+ K^- \pi^+ \pi^0$ with 9 components.				

$\Gamma(\eta(1475) \pi^+, \eta \rightarrow a_0(980)^0 \pi^0, a_0^0 \rightarrow K^+ K^-) / \Gamma_{\text{total}}$ Γ_{81} / Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
0.07±0.02±0.02	3k	¹ ABLIKIM	21U BES3	$e^+ e^-$ at 4.178–4.226 GeV
¹ ABLIKIM 21U uses an amplitude analysis of $D_s^+ \rightarrow K^+ K^- \pi^+ \pi^0$ with 9 components.				

$\Gamma(K_S^0 K^- 2\pi^+) / \Gamma_{\text{total}}$ Γ_{82} / Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
1.569±0.028 OUR FIT				
1.569±0.032 OUR AVERAGE				Error includes scale factor of 1.2.
1.56 ±0.02 ±0.02		ABLIKIM	24BD BES3	$e^+ e^-$, 4.128–4.226 GeV
1.69 ±0.07 ±0.08		ONYISI	13 CLEO	$e^+ e^-$ at 4.17 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.46 ±0.05 ±0.05	1.3k	¹ ABLIKIM	21K BES3	$e^+ e^-$, 4.178–4.226 GeV
1.64 ±0.10 ±0.07		² ALEXANDER	08 CLEO	See ONYISI 13
¹ Superseded by ABLIKIM 24BD.				
² ALEXANDER 08 uses single- and double-tagged events in an overall fit.				

$\Gamma(K^+ K^- K_S^0 \pi^+) / \Gamma(K^+ K_S^0 \pi^+ \pi^-)$ $\Gamma_{83} / \Gamma_{89}$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
1.36±0.15±0.04	645	MOON	23 BELL	980 fb ⁻¹ at $\sim \gamma(4S)$

$\Gamma(K^*(892)^+ \bar{K}^*(892)^0) / \Gamma_{\text{total}}$ Γ_{84} / Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
5.64±0.23±0.27	3k	¹ ABLIKIM	21U BES3	$e^+ e^-$ at 4.178–4.226 GeV
¹ ABLIKIM 21U uses an amplitude analysis of $D_s^+ \rightarrow K^+ K^- \pi^+ \pi^0$ with 9 components.				

$\Gamma(K^*(892)^+ \bar{K}^*(892)^0) / \Gamma(\phi \pi^+)$ $\Gamma_{84} / \Gamma_{53}$

Unseen decay modes of the resonances are included.

VALUE	DOCUMENT ID	TECN	COMMENT
1.6±0.4±0.4	ALBRECHT	92B ARG	$e^+ e^- \simeq 10.4$ GeV

$\Gamma(K^*(892)^+ \bar{K}^*(892)^0) / \Gamma(K_S^0 K^- 2\pi^+)$ $\Gamma_{84} / \Gamma_{82}$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
40.6±2.9±4.9	1.3k	^{1,2} ABLIKIM	21K BES3	$e^+ e^-$ at 4.178–4.226 GeV
¹ Predominantly S -wave, with a significant D -wave component.				
² $D_s^+ \rightarrow K_S^0 K^- 2\pi^+$ amplitude analysis with 13 components.				

$\Gamma(\eta(1475) K_S^0, \eta \rightarrow K^*(892)^0 \pi^+, K^{*0} \rightarrow K^- \pi^+) / \Gamma(K_S^0 K^- 2\pi^+)$ $\Gamma_{85} / \Gamma_{82}$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
2.2±0.6±0.2	1.3k	¹ ABLIKIM	21K BES3	$e^+ e^-$ at 4.178–4.226 GeV
¹ $D_s^+ \rightarrow K_S^0 K^- 2\pi^+$ amplitude analysis with 13 components.				

$$\Gamma(\eta(1475)\pi^+, \eta \rightarrow \bar{K}^*(892)^+ K^-, \bar{K}^{*+} \rightarrow K_S^0 \pi^+)/\Gamma(K_S^0 K^- 2\pi^+) \quad \Gamma_{86}/\Gamma_{82}$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
2.2±0.6±0.2	1.3k	¹ ABLIKIM	21k BES3	$e^+ e^-$ at 4.178–4.226 GeV
¹ $D_s^+ \rightarrow K_S^0 K^- 2\pi^+$ amplitude analysis with 13 components.				

$$\Gamma(\eta(1475)\pi^+, \eta \rightarrow a_0(980)^- \pi^+, a_0^- \rightarrow K_S^0 K^-)/\Gamma(K_S^0 K^- 2\pi^+) \quad \Gamma_{87}/\Gamma_{82}$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
10.8±2.6±5.2	1.3k	¹ ABLIKIM	21k BES3	$e^+ e^-$ at 4.178–4.226 GeV
¹ $D_s^+ \rightarrow K_S^0 K^- 2\pi^+$ amplitude analysis with 13 components.				

$$\Gamma(f_1(1285)\pi^+, f_1 \rightarrow a_0(980)^- \pi^+, a_0^- \rightarrow K_S^0 K^-)/\Gamma(K_S^0 K^- 2\pi^+) \quad \Gamma_{88}/\Gamma_{82}$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
2.2±0.5±0.2	1.3k	¹ ABLIKIM	21k BES3	$e^+ e^-$ at 4.178–4.226 GeV
¹ $D_s^+ \rightarrow K_S^0 K^- 2\pi^+$ amplitude analysis with 13 components.				

$$\Gamma(K^+ K_S^0 \pi^+ \pi^-)/\Gamma_{\text{total}} \quad \Gamma_{89}/\Gamma$$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
0.934±0.022 OUR FIT			
0.935±0.022 OUR AVERAGE			
0.93 ±0.02 ±0.01	ABLIKIM	24BD BES3	$e^+ e^-$, 4.128–4.226 GeV
1.03 ±0.06 ±0.08	ONYISI	13 CLEO	$e^+ e^-$ at 4.17 GeV

$$\Gamma(K^+ K_S^0 \pi^+ \pi^-)/\Gamma(K_S^0 K^- 2\pi^+) \quad \Gamma_{89}/\Gamma_{82}$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.595±0.017 OUR FIT				
0.586±0.052±0.043	476	LINK	01C FOCS	γ A, $\bar{E}_\gamma \approx 180$ GeV

$$\Gamma(K^+ K^- 2\pi^+ \pi^-)/\Gamma_{\text{total}} \quad \Gamma_{90}/\Gamma$$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
6.60±0.47±0.38	309	ABLIKIM	22AB BES3	$e^+ e^-$ at 4.178–4.226 GeV

$$\Gamma(K^+ K^- 2\pi^+ \pi^-)/\Gamma(K^+ K^- \pi^+) \quad \Gamma_{90}/\Gamma_{52}$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.160±0.027 OUR AVERAGE				
0.150±0.019±0.025	240	LINK	03D FOCS	γ A, $\bar{E}_\gamma \approx 180$ GeV
0.188±0.036±0.040	75	FRABETTI	97C E687	γ Be, $\bar{E}_\gamma \approx 200$ GeV

$$\Gamma(\phi 2\pi^+ \pi^-)/\Gamma(\phi \pi^+) \quad \Gamma_{91}/\Gamma_{53}$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.269±0.027 OUR AVERAGE				
0.249±0.024±0.021	136	LINK	03D FOCS	γ A, $\bar{E}_\gamma \approx 180$ GeV
0.28 ±0.06 ±0.01	40	FRABETTI	97C E687	γ Be, $\bar{E}_\gamma \approx 200$ GeV
0.58 ±0.21 ±0.10	21	FRABETTI	92 E687	γ Be
0.42 ±0.13 ±0.07	19	ANJOS	88 E691	Photoproduction
1.11 ±0.37 ±0.28	62	ALBRECHT	85D ARG	$e^+ e^-$ 10 GeV

$\Gamma(\phi\rho^0\pi^+, \phi \rightarrow K^+K^-)/\Gamma(K^+K^-2\pi^+\pi^-)$ Γ_{92}/Γ_{90}

VALUE	DOCUMENT ID	TECN	COMMENT
0.75±0.06±0.04	LINK	03D FOCS	γ A, $\bar{E}_\gamma \approx 180$ GeV

$\Gamma(\phi a_1(1260)^+, \phi \rightarrow K^+K^-, a_1^+ \rightarrow \rho^0\pi^+)/\Gamma(K^+K^-\pi^+)$ Γ_{93}/Γ_{52}

VALUE	DOCUMENT ID	TECN	COMMENT
0.137±0.019±0.011	LINK	03D FOCS	γ A, $\bar{E}_\gamma \approx 180$ GeV

$\Gamma(\phi a_1(1260)^+, \phi \rightarrow K^+K^-, a_1^+ \rightarrow \rho^0\pi^+)/\Gamma(K^+K^-2\pi^+\pi^-)$ Γ_{93}/Γ_{90}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.781±0.029±0.016	235	ABLIKIM	22AB BES3	e^+e^- at 4.178–4.226 GeV

$\Gamma(\phi 2\pi^+\pi^- \text{ non-}\rho, \phi \rightarrow K^+K^-)/\Gamma(K^+K^-2\pi^+\pi^-)$ Γ_{94}/Γ_{90}

VALUE	DOCUMENT ID	TECN	COMMENT
0.21±0.05±0.06	LINK	03D FOCS	γ A, $\bar{E}_\gamma \approx 180$ GeV

$\Gamma(K^+K^-\rho^0\pi^+ \text{ non-}\phi)/\Gamma(K^+K^-2\pi^+\pi^-)$ Γ_{95}/Γ_{90}

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.03	90	LINK	03D FOCS	γ A, $\bar{E}_\gamma \approx 180$ GeV

$\Gamma(K^+K^-2\pi^+\pi^- \text{ nonresonant})/\Gamma(K^+K^-2\pi^+\pi^-)$ Γ_{96}/Γ_{90}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.15 ±0.06 OUR AVERAGE				
0.218±0.029±0.08	235	ABLIKIM	22AB BES3	e^+e^- at 4.178–4.226 GeV
0.10 ±0.06 ±0.05		LINK	03D FOCS	γ A, $\bar{E}_\gamma \approx 180$ GeV

$\Gamma(2K_S^0 2\pi^+\pi^-)/\Gamma(K_S^0 K^- 2\pi^+)$ Γ_{97}/Γ_{82}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.051±0.015±0.015	37 ± 10	LINK	04D FOCS	γ A, $\bar{E}_\gamma \approx 180$ GeV

———— Pionic modes ————

$\Gamma(\pi^+\pi^0)/\Gamma_{\text{total}}$ Γ_{98}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.2 × 10⁻⁴	90	¹ GUAN	21 BELL	$e^+e^- \approx \Upsilon(4,5S)$

¹ Uses $B(D_S^+ \rightarrow \pi^+\phi, \phi \rightarrow K^+K^-) = (2.24 \pm 0.08)\%$.

$\Gamma(\pi^+\pi^0)/\Gamma(K^+K_S^0)$ Γ_{98}/Γ_{44}

VALUE (units 10 ⁻²)	CL%	DOCUMENT ID	TECN	COMMENT
<2.3	90	MENDEZ	10 CLEO	e^+e^- at 4170 MeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

<4.1	90	ADAMS	07A CLEO	See MENDEZ 10
------	----	-------	----------	---------------

$\Gamma(2\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{99}/Γ

VALUE (%)	DOCUMENT ID	TECN	COMMENT
1.090±0.014 OUR FIT			
1.091±0.014 OUR AVERAGE			

1.09 ±0.01 ±0.01	ABLIKIM	24BD BES3	e^+e^- , 4.128–4.226 GeV
1.11 ±0.04 ±0.04	ONYISI	13 CLEO	e^+e^- at 4.17 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.11 ± 0.07 ± 0.04 ¹ ALEXANDER 08 CLEO See ONYISI 13

¹ ALEXANDER 08 uses single- and double-tagged events in an overall fit.

$\Gamma(2\pi^+\pi^-)/\Gamma(K^+K^-\pi^+)$ Γ_{99}/Γ_{52}

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.200 ± 0.004 OUR FIT				Error includes scale factor of 1.2.
0.199 ± 0.004 ± 0.009	≈ 10.5k	AUBERT	09O BABR	$e^+e^- \approx 10.6$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.265 ± 0.041 ± 0.031 98 FRABETTI 97D E687 γ Be ≈ 200 GeV

$\Gamma(\rho^0\pi^+)/\Gamma(2\pi^+\pi^-)$ Γ_{100}/Γ_{99}

<u>VALUE (units 10⁻²)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.04 ± 0.15 OUR AVERAGE			
1.038 ± 0.054 ± 0.097 ± 0.11	¹ AAIJ	23AN LHCB	Dalitz fit, 0.7M events
0.9 ± 0.4 ± 0.5	ABLIKIM	22BI BES3	Dalitz fit, 11.1k events
1.8 ± 0.5 ± 1.0	AUBERT	09O BABR	Dalitz fit, ≈ 10.5k evts

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen LINK 04 FOCS Dalitz fit, 1475 ± 50 evts
 5.8 ± 2.3 ± 3.7 AITALA 01A E791 Dalitz fit, 848 evts

¹ The last error reflects the uncertainty on the amplitude model.

$\Gamma(\omega\pi^+, \omega \rightarrow \pi^+\pi^-)/\Gamma(2\pi^+\pi^-)$ Γ_{101}/Γ_{99}

<u>VALUE (units 10⁻³)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
3.60 ± 0.16 ± 0.34 ± 0.16	¹ AAIJ	23AN LHCB	Dalitz fit, 0.7M events

¹ The last error reflects the uncertainty on the amplitude model.

$\Gamma(\pi^+(\pi^+\pi^-)_{S\text{-wave}})/\Gamma(2\pi^+\pi^-)$ Γ_{102}/Γ_{99}

This is the “fit fraction” from the Dalitz-plot analysis. See also KLEMPT 08, which uses 568 $D_s^+ \rightarrow 3\pi$ decays (over 280 background events) from FNAL E791 to study various parametrizations of the decay amplitudes. The emphasis there is more on S-wave $\pi\pi$ decay products — 20 different solutions are given — than on D_s^+ fit fractions.

<u>VALUE (units 10⁻²)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
84.7 ± 0.6 OUR AVERAGE			
84.97 ± 0.14 ± 0.30 ± 0.63	^{1,2} AAIJ	23AN LHCB	Dalitz fit, 0.7M events
84.2 ± 0.8 ± 1.2	² ABLIKIM	22BI BES3	Dalitz fit, 11.1k events
83.0 ± 0.9 ± 1.9	² AUBERT	09O BABR	Dalitz fit, ≈ 10.5k evts
87.04 ± 5.60 ± 4.38	³ LINK	04 FOCS	Dalitz fit, 1475 ± 50 evts

¹ The last error reflects the uncertainty due to the amplitude model.

² AAIJ 23AN, ABLIKIM 22BI, and AUBERT 09O give the amplitude and phase of the $\pi^+\pi^-$ S-wave in bins of $\pi^+\pi^-$ invariant-mass. (50 bins for AAIJ 23AN, 29 for ABLIKIM 22BI and AUBERT 09O.)

³ LINK 04 borrows a K-matrix parametrization from ANISOVICH 03 of the full $\pi-\pi$ S-wave isoscalar scattering amplitude to describe the $\pi^+\pi^-$ S-wave component of the $\pi^+\pi^+\pi^-$ state. The fit fraction given above is a sum over five f_0 mesons, the $f_0(980)$, $f_0(1300)$, $f_0(1200-1600)$, $f_0(1500)$, and $f_0(1750)$. See LINK 04 for details and discussion.

$\Gamma(f_0(980)\pi^+, f_0 \rightarrow \pi^+\pi^-)/\Gamma(2\pi^+\pi^-)$ Γ_{103}/Γ_{99}

This is the “fit fraction” from the Dalitz-plot analysis. See above for the full $\pi^+(\pi^+\pi^-)_{S\text{-wave}}$ fit fraction.

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.565 \pm 0.043 \pm 0.047$	AITALA	01A	E791 Dalitz fit, 848 evts
$1.074 \pm 0.140 \pm 0.043$	FRABETTI	97D	E687 γ Be \approx 200 GeV

$\Gamma(f_0(1370)\pi^+, f_0 \rightarrow \pi^+\pi^-)/\Gamma(2\pi^+\pi^-)$ Γ_{104}/Γ_{99}

This is the “fit fraction” from the Dalitz-plot analysis. See above for the full $\pi^+(\pi^+\pi^-)_{S\text{-wave}}$ fit fraction.

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.324 \pm 0.077 \pm 0.017$	AITALA	01A	E791 Dalitz fit, 848 evts
-----------------------------	--------	-----	---------------------------

$\Gamma(f_0(1500)\pi^+, f_0 \rightarrow \pi^+\pi^-)/\Gamma(2\pi^+\pi^-)$ Γ_{105}/Γ_{99}

This is the “fit fraction” from the Dalitz-plot analysis. See above for the full $\pi^+(\pi^+\pi^-)_{S\text{-wave}}$ fit fraction.

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.274 \pm 0.114 \pm 0.019$	¹ FRABETTI	97D	E687 γ Be \approx 200 GeV
-----------------------------	-----------------------	-----	------------------------------------

¹FRABETTI 97D calls this mode $S(1475)\pi^+$, but finds the mass and width of this $S(1475)$ to be in excellent agreement with those of the $f_0(1500)$.

$\Gamma(f_2(1270)\pi^+, f_2 \rightarrow \pi^+\pi^-)/\Gamma(2\pi^+\pi^-)$ Γ_{106}/Γ_{99}

This is the “fit fraction” from the Dalitz-plot analysis.

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

13.0 \pm 0.9 OUR AVERAGE Error includes scale factor of 1.9. See the ideogram below.


$13.69 \pm 0.14 \pm 0.22 \pm 0.49$	¹ AAIJ	23AN	LHCB Dalitz fit, 0.7M events
$10.5 \pm 0.8 \pm 1.1$	ABLIKIM	22BI	BES3 Dalitz fit, 11.1k events
$10.1 \pm 1.5 \pm 1.1$	AUBERT	09O	BABR Dalitz fit, \approx 10.5k evts
$9.74 \pm 4.49 \pm 2.94$	LINK	04	FOCS Dalitz fit, 1475 ± 50 evts

• • • We do not use the following data for averages, fits, limits, etc. • • •

$19.7 \pm 3.3 \pm 0.6$	AITALA	01A	E791 Dalitz fit, 848 evts
$12.3 \pm 5.6 \pm 1.8$	FRABETTI	97D	E687 γ Be \approx 200 GeV

¹The last error reflects the uncertainty on the amplitude model.

WEIGHTED AVERAGE
 13.0 ± 0.9 (Error scaled by 1.9)



$\Gamma(f'_2(1525)^0 \pi^+, f'_2 \rightarrow \pi^+ \pi^-) / \Gamma(2\pi^+ \pi^-)$ **$\Gamma_{107} / \Gamma_{99}$**

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$5.28 \pm 0.70 \pm 1.50 \pm 0.87$	¹ AAIJ	23AN LHCB	Dalitz fit, 0.7M events

¹ The last error reflects the uncertainty on the amplitude model.

$\Gamma(\rho(1450)^0 \pi^+, \rho^0 \rightarrow \pi^+ \pi^-) / \Gamma(2\pi^+ \pi^-)$ **$\Gamma_{108} / \Gamma_{99}$**

This is the "fit fraction" from the Dalitz-plot analysis.

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
1.7 ± 0.6 OUR AVERAGE			
$3.86 \pm 0.15 \pm 0.14 \pm 2.0$	¹ AAIJ	23AN LHCB	Dalitz fit, 0.7M events
$1.3 \pm 0.4 \pm 0.5$	ABLIKIM	22BI BES3	Dalitz fit, 11.1k events
$2.3 \pm 0.8 \pm 1.7$	AUBERT	09O BABR	Dalitz fit, ≈ 10.5 k evts
$6.56 \pm 3.43 \pm 4.40$	LINK	04 FOCS	Dalitz fit, 1475 ± 50 evts
$4.4 \pm 2.1 \pm 0.2$	AITALA	01A E791	Dalitz fit, 848 evts

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ The last error reflects the uncertainty on the amplitude model.

$\Gamma(\rho(1700)^0 \pi^+, \rho^0 \rightarrow \pi^+ \pi^-) / \Gamma(2\pi^+ \pi^-)$ **$\Gamma_{109} / \Gamma_{99}$**

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
$3.65 \pm 0.50 \pm 0.45 \pm 3.40$	¹ AAIJ	23AN LHCB	Dalitz fit, 0.7M events

¹ The last error reflects the uncertainty on the amplitude model.

$\Gamma(\pi^+ 2\pi^0) / \Gamma_{\text{total}}$ **Γ_{110} / Γ**

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.52 ± 0.05 OUR AVERAGE				Error includes scale factor of 1.1.
$0.50 \pm 0.04 \pm 0.02$	590	ABLIKIM	22Z BES3	$e^+ e^-$ at 4.178–4.226 GeV

0.65±0.13±0.03 72 ± 16 NAIK 09A CLEO e⁺e⁻ at 4170 MeV

$\Gamma(f_0(980)\pi^+, f_0 \rightarrow \pi^0\pi^0)/\Gamma(\pi^+2\pi^0)$ $\Gamma_{111}/\Gamma_{110}$

This is a "fit fraction" from an amplitude analysis.

VALUE (units 10 ⁻²)	DOCUMENT ID	TECN	COMMENT
55.4±6.8±7.3	ABLIKIM	22Z	BES3 Dalitz plot fit, 440 evts

$\Gamma(f_0(1370)\pi^+, f_0 \rightarrow \pi^0\pi^0)/\Gamma(\pi^+2\pi^0)$ $\Gamma_{112}/\Gamma_{110}$

This is a "fit fraction" from an amplitude analysis.

VALUE (units 10 ⁻²)	DOCUMENT ID	TECN	COMMENT
25.5±5.1±9.3	ABLIKIM	22Z	BES3 Dalitz plot fit, 440 evts

$\Gamma(f_2(1270)\pi^+, f_2 \rightarrow \pi^0\pi^0)/\Gamma(\pi^+2\pi^0)$ $\Gamma_{113}/\Gamma_{110}$

This is a "fit fraction" from an amplitude analysis.

VALUE (units 10 ⁻²)	DOCUMENT ID	TECN	COMMENT
9.7±2.9±6.0	ABLIKIM	22Z	BES3 Dalitz plot fit, 440 evts

$\Gamma(2\pi^+\pi^-\pi^0)/\Gamma(\phi\pi^+)$ Γ_{114}/Γ_{53}

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
•••				We do not use the following data for averages, fits, limits, etc. •••
<3.3	90	ANJOS	89E E691	Photoproduction

$\Gamma(\omega\pi^+)/\Gamma_{total}$ Γ_{115}/Γ

Unseen decay modes of the ω are included.

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.193±0.030 OUR FIT				
0.181±0.032 OUR AVERAGE				
0.177±0.032±0.013	65 ± 12	ABLIKIM	19AH BES3	e ⁺ e ⁻ at 4.178 GeV
0.21 ±0.09 ±0.01	6 ± 2.4	GE	09A CLEO	e ⁺ e ⁻ at 4170 MeV

$\Gamma(\omega\pi^+)/\Gamma(\eta\pi^+)$ $\Gamma_{115}/\Gamma_{116}$

Unseen decay modes of the resonances are included.

VALUE	DOCUMENT ID	TECN	COMMENT
0.114±0.018 OUR FIT			
0.16 ±0.04 ±0.03	BALEST	97	CLE2 e ⁺ e ⁻ ≈ $\Upsilon(4S)$

$\Gamma(\eta\pi^+)/\Gamma_{total}$ Γ_{116}/Γ

Unseen decay modes of the η are included.

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
1.686±0.027 OUR FIT				
1.689±0.027 OUR AVERAGE				
1.69 ±0.02 ±0.02		ABLIKIM	24BD BES3	e ⁺ e ⁻ , 4.128–4.226 GeV
1.67 ±0.08 ±0.06		ONYISI	13 CLEO	e ⁺ e ⁻ at 4.17 GeV
•••				We do not use the following data for averages, fits, limits, etc. •••
1.56 ±0.09 ±0.02	0.4k	^{1,2} ABLIKIM	25A BES3	e ⁺ e ⁻ , 4.128–4.226 GeV
1.82 ±0.14 ±0.07	0.8k	ZUPANC	13 BELL	e ⁺ e ⁻ at $\Upsilon(4S), \Upsilon(5S)$
1.58 ±0.11 ±0.18		³ ALEXANDER	08 CLEO	See ONYISI 13

¹ Subsumed by ABLIKIM 24BD which uses both $\eta \rightarrow \pi^+\pi^-\pi^0$ and $\eta \rightarrow 2\gamma$ modes.
² ABLIKIM 25A reports $[\Gamma(D_s^+ \rightarrow \eta\pi^+)/\Gamma_{total}] \times [B(\eta \rightarrow \pi^+\pi^-\pi^0)] = (3.58 \pm 0.21 \pm 0.06) \times 10^{-3}$ which we divide by our best (shown rounded) value $B(\eta \rightarrow \pi^+\pi^-\pi^0) = (23.02 \pm 0.25) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best (shown rounded) value.
³ ALEXANDER 08 uses single- and double-tagged events in an overall fit.

$\Gamma(\eta\pi^+)/\Gamma(K^+K_S^0)$ Γ_{116}/Γ_{44}

Unseen decay modes of the η are included.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

1.124±0.021 OUR FIT

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.236±0.043±0.063	2587 ± 89	MENDEZ	10	CLEO	See ONYISI 13
-------------------	-----------	--------	----	------	---------------

$\Gamma(\eta\pi^+)/\Gamma(K^+K^-\pi^+)$ Γ_{116}/Γ_{52}

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

31.94±0.33±0.49	19.5k	ABLIKIM	20R	BES3	e^+e^- , 4178 ~ 4226 MeV
------------------------	-------	---------	-----	------	----------------------------

$\Gamma(\eta\pi^+)/\Gamma(\phi\pi^+)$ Γ_{116}/Γ_{53}

Unseen decay modes of the resonances are included.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.48±0.03±0.04	920	JESSOP	98	CLE2	$e^+e^- \approx \Upsilon(4S)$
----------------	-----	--------	----	------	-------------------------------

0.54±0.09±0.06	165	ALEXANDER	92	CLE2	See JESSOP 98
----------------	-----	-----------	----	------	---------------

$\Gamma(\eta\pi^+)/\Gamma(\phi\pi^+, \phi \rightarrow K^+K^-)$ Γ_{116}/Γ_{54}

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

84.80±0.47±2.64	22k	GUAN	21	BELL	$e^+e^- \approx \Upsilon(4,5S)$
------------------------	-----	------	----	------	---------------------------------

$\Gamma((2\pi^+\pi^-\pi^0)_{\text{non-}\eta})/\Gamma_{\text{total}}$ Γ_{117}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

2.04±0.08±0.05	2.5k	ABLIKIM	25A	BES3	e^+e^- at 4.128–4.226 GeV
-----------------------	------	---------	-----	------	-----------------------------

$\Gamma(f_0(1370)\rho^+, f_0\rho^+ \rightarrow 2\pi^+\pi^-\pi^0)/\Gamma((2\pi^+\pi^-\pi^0)_{\text{non-}\eta})$ $\Gamma_{118}/\Gamma_{117}$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

24.9±3.8±2.1	2.5k	¹ ABLIKIM	25A	BES3	e^+e^- at 4.128–4.226 GeV
---------------------	------	----------------------	-----	------	-----------------------------

¹ Amplitude analysis with 11 components.

$\Gamma(f_0(980)^0\rho^+, f_0^0\rho^+ \rightarrow 2\pi^+\pi^-\pi^0)/\Gamma((2\pi^+\pi^-\pi^0)_{\text{non-}\eta})$ $\Gamma_{119}/\Gamma_{117}$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

12.6±2.1±1.0	2.5k	¹ ABLIKIM	25A	BES3	e^+e^- at 4.128–4.226 GeV
---------------------	------	----------------------	-----	------	-----------------------------

¹ Amplitude analysis with 11 components.

$\Gamma(f_2(1270)^0\rho^+, f_2(1270)^0\rho^+ \rightarrow 2\pi^+\pi^-\pi^0)/\Gamma((2\pi^+\pi^-\pi^0)_{\text{non-}\eta})$ $\Gamma_{120}/\Gamma_{117}$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

9.5±1.7±0.6	2.5k	¹ ABLIKIM	25A	BES3	e^+e^- at 4.128–4.226 GeV
--------------------	------	----------------------	-----	------	-----------------------------

¹ Amplitude analysis with 11 components.

$\Gamma((\rho^+\rho^0)_{S\text{-wave}} \rightarrow 2\pi^+\pi^-\pi^0)/\Gamma((2\pi^+\pi^-\pi^0)_{\text{non-}\eta})$ $\Gamma_{121}/\Gamma_{117}$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

3.5±1.2±0.6	2.5k	¹ ABLIKIM	25A	BES3	e^+e^- at 4.128–4.226 GeV
--------------------	------	----------------------	-----	------	-----------------------------

¹ Amplitude analysis with 11 components.

$$\Gamma((\rho(1450)^+ \rho^0)_{S\text{-wave}} \rightarrow 2\pi^+ \pi^- \pi^0) / \Gamma((2\pi^+ \pi^- \pi^0)_{\text{non-}\eta}) \quad \Gamma_{122} / \Gamma_{117}$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
$4.6 \pm 1.3 \pm 0.8$	2.5k	¹ ABLIKIM	25A	BES3 $e^+ e^-$ at 4.128–4.226 GeV

¹ Amplitude analysis with 11 components.

$$\Gamma((\rho^+ \rho(1450)^0)_{P\text{-wave}} \rightarrow 2\pi^+ \pi^- \pi^0) / \Gamma((2\pi^+ \pi^- \pi^0)_{\text{non-}\eta}) \quad \Gamma_{123} / \Gamma_{117}$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
$8.6 \pm 1.3 \pm 0.4$	2.5k	¹ ABLIKIM	25A	BES3 $e^+ e^-$ at 4.128–4.226 GeV

¹ Amplitude analysis with 11 components.

$$\Gamma(\phi \pi^+, \phi \rightarrow \rho \pi) / \Gamma((2\pi^+ \pi^- \pi^0)_{\text{non-}\eta}) \quad \Gamma_{124} / \Gamma_{117}$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
$24.9 \pm 1.2 \pm 0.4$	2.5k	¹ ABLIKIM	25A	BES3 $e^+ e^-$ at 4.128–4.226 GeV

¹ Amplitude analysis with 11 components.

$$\Gamma(\omega \pi^+, \omega \rightarrow \rho \pi) / \Gamma((2\pi^+ \pi^- \pi^0)_{\text{non-}\eta}) \quad \Gamma_{125} / \Gamma_{117}$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
$6.9 \pm 0.8 \pm 0.3$	2.5k	¹ ABLIKIM	25A	BES3 $e^+ e^-$ at 4.128–4.226 GeV

¹ Amplitude analysis with 11 components.

$$\Gamma(a_1(1260)^+ \pi^0, a_1^+ \rightarrow (\rho^0 \pi^+)_{S\text{-wave}}) / \Gamma((2\pi^+ \pi^- \pi^0)_{\text{non-}\eta}) \quad \Gamma_{126} / \Gamma_{117}$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
$12.5 \pm 1.6 \pm 1.0$	2.5k	¹ ABLIKIM	25A	BES3 $e^+ e^-$ at 4.128–4.226 GeV

¹ Amplitude analysis with 11 components.

$$\Gamma(a_1(1260)^0 \pi^+, a_1^0 \rightarrow (\rho \pi)_{S\text{-wave}}) / \Gamma((2\pi^+ \pi^- \pi^0)_{\text{non-}\eta}) \quad \Gamma_{127} / \Gamma_{117}$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
$6.3 \pm 1.9 \pm 1.2$	2.5k	¹ ABLIKIM	25A	BES3 $e^+ e^-$ at 4.128–4.226 GeV

¹ Amplitude analysis with 11 components.

$$\Gamma(\pi(1300)^0 \pi^+, \pi^0 \rightarrow (\rho \pi)_{P\text{-wave}}) / \Gamma((2\pi^+ \pi^- \pi^0)_{\text{non-}\eta}) \quad \Gamma_{128} / \Gamma_{117}$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
$11.7 \pm 2.3 \pm 2.2$	2.5k	¹ ABLIKIM	25A	BES3 $e^+ e^-$ at 4.128–4.226 GeV

¹ Amplitude analysis with 11 components.

$$\Gamma(3\pi^+ 2\pi^-) / \Gamma(K^+ K^- \pi^+) \quad \Gamma_{129} / \Gamma_{52}$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.146 ± 0.014 OUR AVERAGE				
$0.145 \pm 0.011 \pm 0.010$	671	LINK	03D	FOCS γ A, $\bar{E}_\gamma \approx 180$ GeV
$0.158 \pm 0.042 \pm 0.031$	37	FRABETTI	97C	E687 γ Be, $\bar{E}_\gamma \approx 200$ GeV

$$\Gamma(2\pi^+ \pi^- 2\pi^0) / \Gamma_{\text{total}} \quad \Gamma_{130} / \Gamma$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
$4.41 \pm 0.15 \pm 0.13$	2k	ABLIKIM	25F	BES3 $e^+ e^-$ at 4.128–4.226 GeV

$$\Gamma(\omega\rho^+, \omega \rightarrow \pi^+\pi^-\pi^0)/\Gamma(2\pi^+\pi^-2\pi^0) \quad \Gamma_{131}/\Gamma_{130}$$

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$19.98 \pm 1.40^{+0.92}_{-1.20}$	2k	¹ ABLIKIM	25F BES3	e^+e^- at 4.128–4.226 GeV

¹ Amplitude analysis of 2.0k $D_S^+ \rightarrow 2\pi^+\pi^-2\pi^0$ events, with 14 components. The value quoted here includes an $\omega\rho^+$ S -, P -, and D -wave, where the D -wave component dominates.

$$\Gamma(\phi\rho^+, \phi \rightarrow \pi^+\pi^-\pi^0)/\Gamma(2\pi^+\pi^-2\pi^0) \quad \Gamma_{132}/\Gamma_{130}$$

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$13.86 \pm 1.03^{+0.47}_{-0.35}$	2k	¹ ABLIKIM	25F BES3	e^+e^- at 4.128–4.226 GeV

¹ Amplitude analysis of 2.0k $D_S^+ \rightarrow 2\pi^+\pi^-2\pi^0$ events, with 14 components. The value quoted here includes a $\phi\rho^+$ S - and P -wave components.

$$\Gamma(\rho(1450)^+\pi^0, \rho(1450)^+ \rightarrow \omega\pi^+, \omega \rightarrow \pi^+\pi^-\pi^0)/\Gamma(2\pi^+\pi^-2\pi^0) \quad \Gamma_{133}/\Gamma_{130}$$

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$7.84 \pm 0.83^{+0.49}_{-0.58}$	2k	¹ ABLIKIM	25F BES3	e^+e^- at 4.128–4.226 GeV

¹ Amplitude analysis of 2.0k $D_S^+ \rightarrow 2\pi^+\pi^-2\pi^0$ events, with 14 components.

$$\Gamma(a_1(1260)^0\rho^+, a_1(1260)^0 \rightarrow \rho^+\pi^-)/\Gamma(2\pi^+\pi^-2\pi^0) \quad \Gamma_{134}/\Gamma_{130}$$

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$11.43 \pm 0.67 \pm 0.35$	2k	¹ ABLIKIM	25F BES3	e^+e^- at 4.128–4.226 GeV

¹ Amplitude analysis of 2.0k $D_S^+ \rightarrow 2\pi^+\pi^-2\pi^0$ events, with 14 components. The value quoted here includes an $a_1\rho$ S - and P -wave components.

$$\Gamma(a_1(1260)^0\rho^+, a_1(1260)^0 \rightarrow \rho^-\pi^+)/\Gamma(2\pi^+\pi^-2\pi^0) \quad \Gamma_{135}/\Gamma_{130}$$

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$7.39 \pm 0.44 \pm 0.26$	2k	¹ ABLIKIM	25F BES3	e^+e^- at 4.128–4.226 GeV

¹ Amplitude analysis of 2.0k $D_S^+ \rightarrow 2\pi^+\pi^-2\pi^0$ events, with 14 components. The value quoted here includes an $a_1\rho$ S - and P -wave components.

$$\Gamma(a_1(1260)^+\rho^0, a_1(1260)^+ \rightarrow \rho^+\pi^0)/\Gamma(2\pi^+\pi^-2\pi^0) \quad \Gamma_{136}/\Gamma_{130}$$

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$16.53 \pm 1.37 \pm 1.52$	2k	¹ ABLIKIM	25F BES3	e^+e^- at 4.128–4.226 GeV

¹ Amplitude analysis of 2.0k $D_S^+ \rightarrow 2\pi^+\pi^-2\pi^0$ events, with 14 components. The value quoted here includes an $a_1\rho$ S - and P -wave components.

$$\Gamma(b_1(1235)^+\pi^0, b_1(1235)^+ \rightarrow \omega\pi^+, \omega \rightarrow \pi^+\pi^-\pi^0)/\Gamma(2\pi^+\pi^-2\pi^0) \quad \Gamma_{137}/\Gamma_{130}$$

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$10.79 \pm 0.98 \pm 0.68$	2k	¹ ABLIKIM	25F BES3	e^+e^- at 4.128–4.226 GeV

¹ Amplitude analysis of 2.0k $D_S^+ \rightarrow 2\pi^+\pi^-2\pi^0$ events, with 14 components.

$\Gamma(b_1(1235)^0 \pi^+, b_1(1235)^0 \rightarrow \omega \pi^0, \omega \rightarrow \pi^+ \pi^- \pi^0) / \Gamma(2\pi^+ \pi^- 2\pi^0)$ $\Gamma_{138} / \Gamma_{130}$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
$14.60 \pm 1.20^{+0.52}_{-0.49}$	2k	¹ ABLIKIM	25F BES3	$e^+ e^-$ at 4.128–4.226 GeV

¹ Amplitude analysis of 2.0k $D_s^+ \rightarrow 2\pi^+ \pi^- 2\pi^0$ events, with 14 components.

$\Gamma(\eta \rho^+) / \Gamma_{\text{total}}$ Γ_{139} / Γ
 Unseen decay modes of the η are included.

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$8.9 \pm 0.6 \pm 0.5$	328 ± 22	NAIK	09A CLEO	$\eta \rightarrow 2\gamma$

$\Gamma(\eta \rho^+) / \Gamma(\phi \pi^+)$ $\Gamma_{139} / \Gamma_{53}$
 Unseen decay modes of the resonances are included.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$2.98 \pm 0.20 \pm 0.39$ 447 JESSOP 98 CLE2 $e^+ e^- \approx \mathcal{T}(4S)$

$2.86 \pm 0.38^{+0.36}_{-0.38}$ 217 AVERY 92 CLE2 See JESSOP 98

$\Gamma(\eta \rho^+) / \Gamma(\eta \pi^+ \pi^0)$ $\Gamma_{139} / \Gamma_{140}$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
$78.3 \pm 5.0 \pm 2.1$	1.2k	ABLIKIM	19BE BES3	$\eta \pi^+ \pi^0$ amplitude analysis

$\Gamma(\eta \pi^+ \pi^0) / \Gamma_{\text{total}}$ Γ_{140} / Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
9.10 ± 0.17 OUR AVERAGE				

$9.10 \pm 0.09 \pm 0.15$ ABLIKIM 24BD BES3 $e^+ e^-$, 4.128–4.226 GeV

$9.2 \pm 0.4 \pm 1.1$ ONYISI 13 CLEO $e^+ e^-$ at 4.17 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$9.50 \pm 0.28 \pm 0.41$ 2.6k ¹ ABLIKIM 19BE BES3 $e^+ e^-$ at 4.178 GeV

¹ Superseded by ABLIKIM 24BD.

$\Gamma(\eta(\pi^+ \pi^0)_{P\text{-wave}}) / \Gamma(\eta \pi^+ \pi^0)$ $\Gamma_{141} / \Gamma_{140}$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
$5.4 \pm 2.1 \pm 2.5$	1.2k	ABLIKIM	19BE BES3	$\eta \pi^+ \pi^0$ amplitude analysis

$\Gamma(a_0(980)^+ \pi^0 \pi^+, a_0(980)^+ \rightarrow \eta \pi^+ \pi^0) / \Gamma(\eta \pi^+ \pi^0)$ $\Gamma_{142} / \Gamma_{140}$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
23.2 ± 2.3 ± 3.3	1.2k	¹ ABLIKIM	19BE BES3	$\eta \pi^+ \pi^0$ amplitude analysis

¹ Coherent sum of $D_s^+ \rightarrow a_0^+ \pi^0 \rightarrow \eta \pi^+ \pi^0$ and $D_s^+ \rightarrow a_0^0 \pi^+ \rightarrow \eta \pi^+ \pi^0$. ABLIKIM 19BE find $a_0(980)^0 - f(980)$ mixing effects negligibly small in this $D_s^+ \rightarrow \eta \pi^+ \pi^0$ Dalitz plot analysis.

$\Gamma(\omega \pi^+ \pi^0) / \Gamma_{\text{total}}$ Γ_{143} / Γ
 Unseen decay modes of the ω are included.

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$2.78 \pm 0.65 \pm 0.25$	34 ± 7.9	GE	09A CLEO	$e^+ e^-$ at 4170 MeV

$$\Gamma(2\pi^+\pi^-\eta)/\Gamma_{\text{total}} \qquad \Gamma_{144}/\Gamma$$

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

3.08±0.06±0.05		ABLIKIM	24BD BES3	e^+e^- , 4.128–4.226 GeV
-----------------------	--	---------	-----------	----------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.12±0.13±0.09	2.1k	¹ ABLIKIM	21AR BES3	e^+e^- , 4.178–4.226 GeV
----------------	------	----------------------	-----------	----------------------------

¹ Superseded by ABLIKIM 24BD.

$$\Gamma(a_1(1260)^+\eta, a_1^+ \rightarrow \rho(770)^0\pi^+, \rho^0 \rightarrow \pi^+\pi^-)/\Gamma(2\pi^+\pi^-\eta) \qquad \Gamma_{145}/\Gamma_{144}$$

<u>VALUE (units 10^{-2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	--------------------	-------------	----------------

55.4±3.9±2.0	¹ ABLIKIM	21AR BES3	e^+e^- at 4.178–4.226 GeV
---------------------	----------------------	-----------	-----------------------------

¹ $D_s^+ \rightarrow 2\pi^+\pi^-\eta$ amplitude analysis with 11 components.

$$\Gamma(a_1(1260)^+\eta, a_1^+ \rightarrow f_0(500)\pi^+, f_0 \rightarrow \pi^+\pi^-)/\Gamma(2\pi^+\pi^-\eta) \qquad \Gamma_{146}/\Gamma_{144}$$

<u>VALUE (units 10^{-2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	--------------------	-------------	----------------

8.1±1.9±2.1	¹ ABLIKIM	21AR BES3	e^+e^- at 4.178–4.226 GeV
--------------------	----------------------	-----------	-----------------------------

¹ $D_s^+ \rightarrow 2\pi^+\pi^-\eta$ amplitude analysis with 11 components.

$$\Gamma(a_0(980)^+\rho(770)^0, a_0^+ \rightarrow \eta\pi^+)/\Gamma(2\pi^+\pi^-\eta) \qquad \Gamma_{147}/\Gamma_{144}$$

<u>VALUE (units 10^{-2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	--------------------	-------------	----------------

6.7±2.5±1.5	¹ ABLIKIM	21AR BES3	e^+e^- at 4.178–4.226 GeV
--------------------	----------------------	-----------	-----------------------------

¹ $D_s^+ \rightarrow 2\pi^+\pi^-\eta$ amplitude analysis with 11 components.

$$\Gamma(\eta(1405)\pi^+, \eta \rightarrow a_0(980)^-\pi^+, a_0^- \rightarrow \eta\pi^-)/\Gamma(2\pi^+\pi^-\eta) \qquad \Gamma_{148}/\Gamma_{144}$$

<u>VALUE (units 10^{-2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	--------------------	-------------	----------------

0.7±0.2±0.1	¹ ABLIKIM	21AR BES3	e^+e^- at 4.178–4.226 GeV
--------------------	----------------------	-----------	-----------------------------

¹ $D_s^+ \rightarrow 2\pi^+\pi^-\eta$ amplitude analysis with 11 components.

$$\Gamma(\eta(1405)\pi^+, \eta \rightarrow a_0(980)^+\pi^-, a_0^+ \rightarrow \eta\pi^+)/\Gamma(2\pi^+\pi^-\eta) \qquad \Gamma_{149}/\Gamma_{144}$$

<u>VALUE (units 10^{-2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	--------------------	-------------	----------------

0.7±0.2±0.1	¹ ABLIKIM	21AR BES3	e^+e^- at 4.178–4.226 GeV
--------------------	----------------------	-----------	-----------------------------

¹ $D_s^+ \rightarrow 2\pi^+\pi^-\eta$ amplitude analysis with 11 components.

$$\Gamma(f_1(1420)\pi^+, f_1 \rightarrow a_0(980)^-\pi^+, a_0^- \rightarrow \eta\pi^-)/\Gamma(2\pi^+\pi^-\eta) \qquad \Gamma_{150}/\Gamma_{144}$$

<u>VALUE (units 10^{-2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	--------------------	-------------	----------------

1.9±0.5±0.3	¹ ABLIKIM	21AR BES3	e^+e^- at 4.178–4.226 GeV
--------------------	----------------------	-----------	-----------------------------

¹ $D_s^+ \rightarrow 2\pi^+\pi^-\eta$ amplitude analysis with 11 components.

$$\Gamma(f_1(1420)\pi^+, f_1 \rightarrow a_0(980)^+\pi^-, a_0^+ \rightarrow \eta\pi^+)/\Gamma(2\pi^+\pi^-\eta) \qquad \Gamma_{151}/\Gamma_{144}$$

<u>VALUE (units 10^{-2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	--------------------	-------------	----------------

1.7±0.5±0.3	¹ ABLIKIM	21AR BES3	e^+e^- at 4.178–4.226 GeV
--------------------	----------------------	-----------	-----------------------------

¹ $D_s^+ \rightarrow 2\pi^+\pi^-\eta$ amplitude analysis with 11 components.

$\Gamma(3\pi^+ 2\pi^- \pi^0)/\Gamma_{\text{total}}$ Γ_{152}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.049^{+0.033}_{-0.030}	BARLAG	92C	ACCM π^- 230 GeV

$\Gamma(\omega 2\pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{153}/Γ

Unseen decay modes of the ω are included.

<u>VALUE (%)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.58\pm0.45\pm0.09	29 \pm 8.2	GE	09A	CLEO $e^+ e^-$ at 4170 MeV

$\Gamma(\eta'(958)\pi^+)/\Gamma_{\text{total}}$ Γ_{154}/Γ

Unseen decay modes of the $\eta'(958)$ are included.

<u>VALUE (%)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
3.95\pm0.08 OUR AVERAGE			

3.95 \pm 0.04 \pm 0.07 ABLIKIM 24BD BES3 $e^+ e^-$, 4.128–4.226 GeV

3.94 \pm 0.15 \pm 0.20 ONYISI 13 CLEO $e^+ e^-$ at 4.17 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.77 \pm 0.25 \pm 0.30 ¹ALEXANDER 08 CLEO See ONYISI 13

¹ALEXANDER 08 uses single- and double-tagged events in an overall fit.

$\Gamma(\eta'(958)\pi^+)/\Gamma(K^+ K_S^0)$ Γ_{154}/Γ_{44}

Unseen decay modes of the $\eta'(958)$ are included.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.654\pm0.088\pm0.139	1436 \pm 47	MENDEZ	10	CLEO See ONYISI 13

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.654 \pm 0.088 \pm 0.139 1436 \pm 47 MENDEZ 10 CLEO See ONYISI 13

$\Gamma(\eta'(958)\pi^+)/\Gamma(K^+ K^- \pi^+)$ Γ_{154}/Γ_{52}

<u>VALUE (units 10⁻²)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
69.4\pm0.8\pm3.8	9.9k	ABLIKIM	20R	BES3 $e^+ e^-$, 4178 \sim 4226 MeV

$\Gamma(\eta'(958)\pi^+)/\Gamma(\phi\pi^+)$ Γ_{154}/Γ_{53}

Unseen decay modes of the resonances are included.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.03\pm0.06\pm0.07	537	JESSOP	98	CLE2 $e^+ e^- \approx \gamma(4S)$
1.20 \pm 0.15 \pm 0.11	281	ALEXANDER	92	CLE2 See JESSOP 98
2.5 \pm 1.0 ^{+1.5} _{-0.4}	22	ALVAREZ	91	NA14 Photoproduction
2.5 \pm 0.5 \pm 0.3	215	ALBRECHT	90D	ARG $e^+ e^- \approx 10.4$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.03 \pm 0.06 \pm 0.07 537 JESSOP 98 CLE2 $e^+ e^- \approx \gamma(4S)$

1.20 \pm 0.15 \pm 0.11 281 ALEXANDER 92 CLE2 See JESSOP 98

2.5 \pm 1.0 ^{+1.5}_{-0.4} 22 ALVAREZ 91 NA14 Photoproduction

2.5 \pm 0.5 \pm 0.3 215 ALBRECHT 90D ARG $e^+ e^- \approx 10.4$ GeV

$\Gamma(\omega\eta\pi^+)/\Gamma_{\text{total}}$ Γ_{156}/Γ

Unseen decay modes of the ω and η are included.

<u>VALUE (units 10⁻³)</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
5.4\pm1.2\pm0.4		78	ABLIKIM	23AL	BES3 $e^+ e^-$ at 4.128–4.226 GeV
<21.3	90		GE	09A	CLEO $e^+ e^-$ at 4170 MeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

<21.3 90 GE 09A CLEO $e^+ e^-$ at 4170 MeV

$\Gamma(\eta'(958)\rho^+)/\Gamma_{\text{total}}$ Γ_{157}/Γ

<u>VALUE (%)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
5.8\pm1.4\pm0.4	ABLIKIM	15Z	BES3 482 pb ⁻¹ , 4009 MeV

$\Gamma(\eta'(958)\rho^+)/\Gamma(\phi\pi^+)$ Γ_{157}/Γ_{53}

Unseen decay modes of the resonances are included.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$2.78 \pm 0.28 \pm 0.30$	137	¹ JESSOP	98 CLE2	$e^+e^- \approx \mathcal{I}(4S)$
$3.44 \pm 0.62^{+0.44}_{-0.46}$	68	AVERY	92 CLE2	See JESSOP 98

¹ This JESSOP 98 fraction, when combined with other η' fractions, greatly overshoots the inclusive η' fraction. See the measurement just above, which fits nicely.

 $\Gamma(\eta'(958)\rho^+)/\Gamma(\eta'(958)\pi^+\pi^0)$ $\Gamma_{157}/\Gamma_{158}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
≈ 1	395	¹ ABLIKIM	22AA BES3	e^+e^- at 4.178–4.226 GeV

¹ Result of an amplitude analysis of $D_S^+ \rightarrow \pi^+\pi^0\eta'$ which found that $D_S^+ \rightarrow \rho^+\eta'$ is the dominant decay mode, with other contributions negligible. No uncertainty is assigned to this 100% fit fraction; however, the fit fractions of non-resonant contributions are shown to be below 1%.

 $\Gamma(\eta'(958)\pi^+\pi^0)/\Gamma_{\text{total}}$ Γ_{158}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
6.14 ± 0.18 OUR AVERAGE				
$6.17 \pm 0.12 \pm 0.14$		ABLIKIM	24BD BES3	e^+e^- , 4.128–4.226 GeV
$5.6 \pm 0.5 \pm 0.6$		ONYISI	13 CLEO	e^+e^- at 4.17 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •				
$6.15 \pm 0.25 \pm 0.18$	837	¹ ABLIKIM	22AA BES3	e^+e^- , 4.178–4.226 GeV

¹ An amplitude analysis in the same publication finds that $D_S^+ \rightarrow \rho^+\eta'$ is the only statistically significant contribution to this decay. Superseded by ABLIKIM 24BD.

 $\Gamma(\eta'(958)\pi^+\pi^0 \text{ nonresonant})/\Gamma_{\text{total}}$ Γ_{159}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 5.1 \times 10^{-2}$	90	ABLIKIM	15Z BES3	482 pb^{-1} , 4009 MeV

Modes with one or three K's

 $\Gamma(K^+\pi^0)/\Gamma(K^+K_S^0)$ Γ_{160}/Γ_{44}

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
4.2 ± 1.4 ± 0.2	202 ± 70	MENDEZ	10 CLEO	e^+e^- at 4170 MeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$5.5 \pm 1.3 \pm 0.7$	141 ± 34	ADAMS	07A CLEO	See MENDEZ 10

 $\Gamma(K^+\pi^0)/\Gamma(K^+K^-\pi^+)$ Γ_{160}/Γ_{52}

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
13.73 ± 0.90 ± 0.33	2.3k	ABLIKIM	20R BES3	e^+e^- , 4178 ~ 4226 MeV

 $\Gamma(K^+\pi^0)/\Gamma(\phi\pi^+, \phi \rightarrow K^+K^-)$ Γ_{160}/Γ_{54}

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
3.28 ± 0.23 ± 0.13	12k	GUAN	21 BELL	$e^+e^- \approx \mathcal{I}(4,5S)$

$$\Gamma(K_S^0 \pi^+)/\Gamma(K^+ K_S^0) \quad \Gamma_{161}/\Gamma_{44}$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
8.12±0.28 OUR AVERAGE				
8.5 ±0.7 ±0.2	393 ± 33	MENDEZ	10	CLEO $e^+ e^-$ at 4170 MeV
8.03±0.24±0.19	17.6k±481	WON	09	BELL $e^+ e^-$ at $\Upsilon(4S)$
10.4 ±2.4 ±1.4	113 ± 26	LINK	08	FOCS $\gamma A, \bar{E}_\gamma \approx 180$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
8.2 ±0.9 ±0.2	206 ± 22	ADAMS	07A	CLEO See MENDEZ 10

$$\Gamma(K_S^0 \pi^+)/\Gamma(K^+ K^- \pi^+) \quad \Gamma_{161}/\Gamma_{52}$$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
20.35±0.62±0.42	2.7k	ABLIKIM	20R	BES3 $e^+ e^-$, 4178 ~ 4226 MeV

$$\Gamma(K^+ \eta)/\Gamma(K^+ K_S^0) \quad \Gamma_{162}/\Gamma_{44}$$

Unseen decay modes of the η are included.

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
11.8±2.2±0.6	222 ± 41	MENDEZ	10	CLEO $e^+ e^-$ at 4170 MeV

$$\Gamma(K^+ \eta)/\Gamma(K^+ K^- \pi^+) \quad \Gamma_{162}/\Gamma_{52}$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
2.97±0.18±0.06	1.8k	ABLIKIM	20R	BES3 $e^+ e^-$, 4178 ~ 4226 MeV

$$\Gamma(K^+ \eta)/\Gamma(\phi \pi^+, \phi \rightarrow K^+ K^-) \quad \Gamma_{162}/\Gamma_{54}$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
7.81±0.22±0.24	14k	GUAN	21	BELL $e^+ e^- \approx \Upsilon(4, 5S)$

$$\Gamma(K^+ \eta)/\Gamma(\eta \pi^+) \quad \Gamma_{162}/\Gamma_{116}$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
8.9±1.5±0.4	113 ± 18	ADAMS	07A	CLEO See MENDEZ 10

$$\Gamma(K^+ \omega)/\Gamma_{\text{total}} \quad \Gamma_{163}/\Gamma$$

Unseen decay modes of the ω are included.

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
9.9±1.5 OUR FIT					
8.7±2.4±0.8		29	¹ ABLIKIM	19AH	BES3 $e^+ e^-$ at 4.178 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<24		90	GE	09A	CLEO $e^+ e^-$ at 4170 MeV

¹Evidence for mode at 4.4σ .

$$\Gamma(K^+ \omega)/\Gamma(K^+ \pi^+ \pi^- \pi^0) \quad \Gamma_{163}/\Gamma_{183}$$

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
10.3±1.5 OUR FIT			
10.9±1.8±0.1	¹ ABLIKIM	22BL	BES3 PWA, 550 $D_s^\pm \rightarrow K^\pm \pi^\pm \pi^\mp \pi^0$

¹ ABLIKIM 22BL reports $[\Gamma(D_s^+ \rightarrow K^+ \omega)/\Gamma(D_s^+ \rightarrow K^+ \pi^+ \pi^- \pi^0)] \times [B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0)] = (9.7 \pm 1.5 \pm 0.6) \times 10^{-2}$ which we divide by our best (shown rounded) value $B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best (shown rounded) value.

$\Gamma(K^+ \eta'(958))/\Gamma(K^+ K_S^0)$ Γ_{164}/Γ_{44}

Unseen decay modes of the $\eta'(958)$ are included.

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
11.8±3.6±0.7	56 ± 17	MENDEZ	10	CLEO $e^+ e^-$ at 4170 MeV

$\Gamma(K^+ \eta'(958))/\Gamma(K^+ K^- \pi^+)$ Γ_{164}/Γ_{52}

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
4.91±0.31±0.31	675	ABLIKIM	20R BES3	$e^+ e^-$, 4178 ~ 4226 MeV

$\Gamma(K^+ \eta'(958))/\Gamma(\eta'(958)\pi^+)$ $\Gamma_{164}/\Gamma_{154}$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.2±1.3±0.3	28 ± 9	ADAMS	07A CLEO	See MENDEZ 10
-------------	--------	-------	----------	---------------

$\Gamma(K^+ \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{165}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

6.23±0.10 OUR FIT

6.22±0.10 OUR AVERAGE

6.20±0.09±0.06		ABLIKIM	24BD BES3	$e^+ e^-$, 4.128–4.226 GeV
6.54±0.33±0.25		ONYISI	13 CLEO	$e^+ e^-$ at 4.17 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

6.11±0.18±0.11	1.3k	¹ ABLIKIM	22AC BES3	$e^+ e^-$, 4.178–4.226 GeV
6.9 ±0.5 ±0.3		² ALEXANDER	08 CLEO	See ONYISI 13

¹ Superseded by ABLIKIM 24BD.

² ALEXANDER 08 uses single- and double-tagged events in an overall fit.

$\Gamma(K^+ \pi^+ \pi^-)/\Gamma(K^+ K^- \pi^+)$ Γ_{165}/Γ_{52}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.1142±0.0025 OUR FIT Error includes scale factor of 1.1.

0.127 ±0.007 ±0.014	567 ± 31	LINK	04F FOCS	$\gamma A, \bar{E}_\gamma \approx 180$ GeV
----------------------------	----------	------	----------	--

$\Gamma(K^+ \rho^0)/\Gamma(K^+ \pi^+ \pi^-)$ $\Gamma_{166}/\Gamma_{165}$

This is the “fit fraction” from the Dalitz-plot analysis.

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.35 ±0.04 OUR AVERAGE

0.321±0.037±0.037	ABLIKIM	22AC BES3	Dalitz plot fit, 1.3k evts
0.388±0.053±0.026	LINK	04F FOCS	Dalitz plot fit, 567 evts

$\Gamma(K^+ \rho(1450)^0, \rho^0 \rightarrow \pi^+ \pi^-)/\Gamma(K^+ \pi^+ \pi^-)$ $\Gamma_{167}/\Gamma_{165}$

This is the “fit fraction” from the Dalitz-plot analysis.

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.117±0.028 OUR AVERAGE

0.131±0.031±0.029	ABLIKIM	22AC BES3	Dalitz plot fit, 1.3k evts
0.106±0.035±0.010	LINK	04F FOCS	Dalitz plot fit, 567 evts

$\Gamma(K^+ f_0(500), f_0 \rightarrow \pi^+ \pi^-)/\Gamma(K^+ \pi^+ \pi^-)$ $\Gamma_{168}/\Gamma_{165}$

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

7.2±2.1±4.4	ABLIKIM	22AC BES3	Dalitz plot fit, 1.3k evts
--------------------	---------	-----------	----------------------------

$\Gamma(K^+ f_0(980), f_0 \rightarrow \pi^+ \pi^-) / \Gamma(K^+ \pi^+ \pi^-)$ $\Gamma_{169} / \Gamma_{165}$

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
4.5 ± 1.3 ± 1.2	ABLIKIM	22AC BES3	Dalitz plot fit, 1.3k evts

 $\Gamma(K^+ f_0(1370), f_0 \rightarrow \pi^+ \pi^-) / \Gamma(K^+ \pi^+ \pi^-)$ $\Gamma_{170} / \Gamma_{165}$

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
19.9 ± 2.9 ± 9.3	ABLIKIM	22AC BES3	Dalitz plot fit, 1.3k evts

 $\Gamma(K^*(892)^0 \pi^+, K^{*0} \rightarrow K^+ \pi^-) / \Gamma(K^+ \pi^+ \pi^-)$ $\Gamma_{171} / \Gamma_{165}$

This is the "fit fraction" from the Dalitz-plot analysis.

VALUE	DOCUMENT ID	TECN	COMMENT
0.27 ± 0.04 OUR AVERAGE	Error includes scale factor of 2.0.		
0.302 ± 0.018 ± 0.020	ABLIKIM	22AC BES3	Dalitz plot fit, 1.3k evts
0.2164 ± 0.0321 ± 0.0114	LINK	04F FOCS	Dalitz plot fit, 567 evts

 $\Gamma(K^*(1410)^0 \pi^+, K^{*0} \rightarrow K^+ \pi^-) / \Gamma(K^+ \pi^+ \pi^-)$ $\Gamma_{172} / \Gamma_{165}$

This is the "fit fraction" from the Dalitz-plot analysis.

VALUE	DOCUMENT ID	TECN	COMMENT
0.10 ± 0.07 OUR AVERAGE	Error includes scale factor of 2.7.		
0.045 ± 0.021 ± 0.025	ABLIKIM	22AC BES3	Dalitz plot fit, 1.3k evts
0.1882 ± 0.0403 ± 0.0122	LINK	04F FOCS	Dalitz plot fit, 567 evts

 $\Gamma(K^*(1430)^0 \pi^+, K^{*0} \rightarrow K^+ \pi^-) / \Gamma(K^+ \pi^+ \pi^-)$ $\Gamma_{173} / \Gamma_{165}$

This is the "fit fraction" from the Dalitz-plot analysis.

VALUE	DOCUMENT ID	TECN	COMMENT
0.15 ± 0.05 OUR AVERAGE	Error includes scale factor of 1.7.		
0.185 ± 0.025 ± 0.026	ABLIKIM	22AC BES3	Dalitz plot fit, 1.3k evts
0.0765 ± 0.0500 ± 0.0170	LINK	04F FOCS	Dalitz plot fit, 567 evts

 $\Gamma(K^+ \pi^+ \pi^- \text{ nonresonant}) / \Gamma(K^+ \pi^+ \pi^-)$ $\Gamma_{174} / \Gamma_{165}$

This is the "fit fraction" from the Dalitz-plot analysis.

VALUE	DOCUMENT ID	TECN	COMMENT
0.1588 ± 0.0492 ± 0.0153	LINK	04F FOCS	Dalitz fit, 567 evts

 $\Gamma(K_S^0 \pi^+ \pi^0) / \Gamma_{\text{total}}$ Γ_{176} / Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
5.09 ± 0.22 OUR AVERAGE				

5.1 ± 0.2 ± 0.1 ABLIKIM 24BD BES3 $e^+ e^-$, 4.128–4.226 GeV5.0 ± 0.9 ± 0.2 44 ¹ NAIK 09A CLEO $e^+ e^-$ at 4170 MeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

5.43 ± 0.30 ± 0.15 666 ² ABLIKIM 21AB BES3 $e^+ e^-$, 4.178–4.226 GeV¹ NAIK 09A reports $B(D_S^+ \rightarrow K^0 \pi^+ \pi^-) = (1.00 \pm 0.18 \pm 0.04) \times 10^{-2}$ which we have divided by 2.² ABLIKIM 21AB uses an amplitude analysis with 5 resonant modes plus one background component. Superseded by ABLIKIM 24BD. $\Gamma(K_S^0 \rho(770)^+, \rho^+ \rightarrow \pi^+ \pi^0) / \Gamma(K_S^0 \pi^+ \pi^0)$ $\Gamma_{177} / \Gamma_{176}$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
50.2 ± 7.2 ± 3.9	666	¹ ABLIKIM	21AB BES3	$e^+ e^-$ at 4.178–4.226 GeV

¹ ABLIKIM 21AB uses an amplitude analysis with 5 resonant modes plus one background component.

$$\Gamma(K_S^0 \rho(1450)^+, \rho^+ \rightarrow \pi^+ \pi^0) / \Gamma(K_S^0 \pi^+ \pi^0) \quad \Gamma_{178} / \Gamma_{176}$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
20.4 ± 4.3 ± 4.4	666	¹ ABLIKIM	21AB BES3	$e^+ e^-$ at 4.178–4.226 GeV

¹ ABLIKIM 21AB uses an amplitude analysis with 5 resonant modes plus one background component.

$$\Gamma(K^*(892)^0 \pi^+, K^{*0} \rightarrow K_S^0 \pi^0) / \Gamma(K_S^0 \pi^+ \pi^0) \quad \Gamma_{179} / \Gamma_{176}$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
8.4 ± 2.2 ± 0.9	666	¹ ABLIKIM	21AB BES3	$e^+ e^-$ at 4.178–4.226 GeV

¹ ABLIKIM 21AB uses an amplitude analysis with 5 resonant modes plus one background component.

$$\Gamma(K^*(892)^+ \pi^0, K^{*+} \rightarrow K_S^0 \pi^+) / \Gamma(K_S^0 \pi^+ \pi^0) \quad \Gamma_{180} / \Gamma_{176}$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
4.6 ± 1.4 ± 0.4	666	¹ ABLIKIM	21AB BES3	$e^+ e^-$ at 4.178–4.226 GeV

¹ ABLIKIM 21AB uses an amplitude analysis with 5 resonant modes plus one background component.

$$\Gamma(K^*(1410)^0 \pi^+, K^{*0} \rightarrow K_S^0 \pi^0) / \Gamma(K_S^0 \pi^+ \pi^0) \quad \Gamma_{181} / \Gamma_{176}$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
3.3 ± 1.6 ± 0.5	666	¹ ABLIKIM	21AB BES3	$e^+ e^-$ at 4.178–4.226 GeV

¹ ABLIKIM 21AB uses an amplitude analysis with 5 resonant modes plus one background component.

$$\Gamma(K_S^0 2\pi^+ \pi^-) / \Gamma(K_S^0 K^- 2\pi^+) \quad \Gamma_{182} / \Gamma_{82}$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.18 ± 0.04 ± 0.05	179 ± 36	LINK	08	FOCS $\gamma A, \bar{E}_{\gamma} \approx 180$ GeV

$$\Gamma(K^+ \pi^+ \pi^- \pi^0) / \Gamma(K^+ K^- \pi^+ \pi^0) \quad \Gamma_{183} / \Gamma_{74}$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
17.13 ± 0.62 ± 0.51	26k	LI	23G BELL	$e^+ e^-$ at/near $\Upsilon(nS)$, $n=1, \dots, 5$

$$\Gamma(K^+ \pi^+ \pi^- \pi^0) / \Gamma_{\text{total}} \quad \Gamma_{183} / \Gamma$$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
9.7 ± 0.6 OUR FIT				
9.75 ± 0.54 ± 0.17	776	ABLIKIM	22BL BES3	$e^+ e^-$ at 4.178–4.226 GeV

$$\Gamma(K^*(892)^0 \rho^+, K^{*0} \rightarrow K^+ \pi^-) / \Gamma(K^+ \pi^+ \pi^- \pi^0) \quad \Gamma_{184} / \Gamma_{183}$$

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
40.5 ± 2.8 ± 1.5	ABLIKIM	22BL BES3	PWA, 550 $D_S^\pm \rightarrow K^\pm \pi^\pm \pi^\mp \pi^0$

$$\Gamma(K^*(892)^+ \rho^0, K^{*+} \rightarrow K^+ \pi^0) / \Gamma(K^+ \pi^+ \pi^- \pi^0) \quad \Gamma_{185} / \Gamma_{183}$$

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
4.3 ± 1.1 ± 0.6	ABLIKIM	22BL BES3	PWA, 550 $D_S^\pm \rightarrow K^\pm \pi^\pm \pi^\mp \pi^0$

$$\Gamma(K_1(1270)^0 \pi^+, K_1^0 \rightarrow K^+ \rho^-) / \Gamma(K^+ \pi^+ \pi^- \pi^0) \quad \Gamma_{186} / \Gamma_{183}$$

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
$4.0 \pm 1.2 \pm 0.6$	ABLIKIM	22BL BES3	PWA, 550 $D_S^\pm \rightarrow K^\pm \pi^\pm \pi^\mp \pi^0$

$$\Gamma(K_1(1400)^0 \pi^+, K_1^0 \rightarrow K^*(890)^+ \pi^-, K^{*+} \rightarrow K^+ \pi^0) / \Gamma(K^+ \pi^+ \pi^- \pi^0) \quad \Gamma_{187} / \Gamma_{183}$$

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
$5.6 \pm 0.9 \pm 0.2$	ABLIKIM	22BL BES3	PWA, 550 $D_S^\pm \rightarrow K^\pm \pi^\pm \pi^\mp \pi^0$

$$\Gamma(K_1(1400)^0 \pi^+, K_1^0 \rightarrow K^*(890)^0 \pi^0, K^{*0} \rightarrow K^+ \pi^-) / \Gamma(K^+ \pi^+ \pi^- \pi^0) \quad \Gamma_{188} / \Gamma_{183}$$

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
$6.1 \pm 0.9 \pm 0.2$	ABLIKIM	22BL BES3	PWA, 550 $D_S^\pm \rightarrow K^\pm \pi^\pm \pi^\mp \pi^0$

$$\Gamma(K^+ a_1(1260)^0, a_1 \rightarrow \rho^+ \pi^-) / \Gamma(K^+ \pi^+ \pi^- \pi^0) \quad \Gamma_{189} / \Gamma_{183}$$

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
$1.9 \pm 0.7 \pm 0.9$	ABLIKIM	22BL BES3	PWA, 550 $D_S^\pm \rightarrow K^\pm \pi^\pm \pi^\mp \pi^0$

$$\Gamma(K^+ a_1(1260)^0, a_1 \rightarrow \rho^- \pi^+) / \Gamma(K^+ \pi^+ \pi^- \pi^0) \quad \Gamma_{190} / \Gamma_{183}$$

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
$1.9 \pm 0.7 \pm 0.9$	ABLIKIM	22BL BES3	PWA, 550 $D_S^\pm \rightarrow K^\pm \pi^\pm \pi^\mp \pi^0$

$$\Gamma(K^+ \pi^+ \pi^- \pi^0 \text{ nonresonant}) / \Gamma(K^+ \pi^+ \pi^- \pi^0) \quad \Gamma_{191} / \Gamma_{183}$$

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
$9.5 \pm 2.2 \pm 0.9$	ABLIKIM	22BL BES3	PWA, 550 $D_S^\pm \rightarrow K^\pm \pi^\pm \pi^\mp \pi^0$

$$\Gamma((K^+ \pi^0)_{\rho\text{-wave}} \rho^0) / \Gamma(K^+ \pi^+ \pi^- \pi^0) \quad \Gamma_{192} / \Gamma_{183}$$

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
$10.4 \pm 2.0 \pm 0.6$	ABLIKIM	22BL BES3	PWA, 550 $D_S^\pm \rightarrow K^\pm \pi^\pm \pi^\mp \pi^0$

$$\Gamma(K^+ \omega \pi^0) / \Gamma_{\text{total}} \quad \Gamma_{193} / \Gamma$$

Unseen decay modes of the ω are included.

VALUE (%)	CL%	DOCUMENT ID	TECN	COMMENT
<0.82	90	GE	09A	CLEO $e^+ e^-$ at 4170 MeV

$$\Gamma(K^+ \omega \pi^+ \pi^-) / \Gamma_{\text{total}} \quad \Gamma_{194} / \Gamma$$

Unseen decay modes of the ω are included.

VALUE (%)	CL%	DOCUMENT ID	TECN	COMMENT
<0.54	90	GE	09A	CLEO $e^+ e^-$ at 4170 MeV

$$\Gamma(K^+ \omega \eta) / \Gamma_{\text{total}} \quad \Gamma_{195} / \Gamma$$

Unseen decay modes of the ω and η are included.

VALUE (%)	CL%	DOCUMENT ID	TECN	COMMENT
<0.79	90	GE	09A	CLEO $e^+ e^-$ at 4170 MeV

$\Gamma(2K^+K^-)/\Gamma(K^+K^-\pi^+)$ Γ_{196}/Γ_{52}

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
4.0 ± 0.3 ± 0.2	748 ± 60	DEL-AMO-SA..11G	BABR	$e^+e^- \approx \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
8.95 ± 2.12 $^{+2.24}_{-2.31}$	31	LINK	02I	FOCS $\gamma A, \approx 180$ GeV

$\Gamma(\phi K^+, \phi \rightarrow K^+K^-)/\Gamma(2K^+K^-)$ $\Gamma_{197}/\Gamma_{196}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.41 ± 0.08 ± 0.03	DEL-AMO-SA..11G	BABR	$e^+e^- \approx \Upsilon(4S)$

———— Radiative decays ————

$\Gamma(\rho(770)^+\gamma)/\Gamma_{\text{total}}$ Γ_{198}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<6.1 × 10⁻⁴	90	¹ ABLIKIM	24CT BES3	e^+e^- at 4.128–4.226 GeV
¹ ABLIKIM 24CT reports an absolute branching fraction $(2.2 \pm 0.9 \pm 0.2) \times 10^{-4}$ at 2.5 σ significance.				

———— Doubly Cabibbo-suppressed modes ————

$\Gamma(2K^+\pi^-)/\Gamma_{\text{total}}$ Γ_{199}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.293 ± 0.027 OUR FIT	Error includes scale factor of 1.1.			
1.24 $^{+0.28}_{-0.26}$ ± 0.06	33 $^{+8}_{-7}$	ABLIKIM	24AV BES3	e^+e^- at 4.128–4.266 GeV

$\Gamma(2K^+\pi^-)/\Gamma(K^+K^-\pi^+)$ Γ_{199}/Γ_{52}

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.371 ± 0.035 OUR FIT				
2.371 ± 0.034 OUR AVERAGE				
2.372 ± 0.024 ± 0.025	67k	AAIJ	19G LHCb	pp at 8 TeV
2.3 ± 0.3 ± 0.2	356 ± 52	DEL-AMO-SA..11G	BABR	$e^+e^- \approx \Upsilon(4S)$
2.29 ± 0.28 ± 0.12	281 ± 34	KO	09 BELL	e^+e^- at $\Upsilon(4S)$
5.2 ± 1.7 ± 1.1	27 ± 9	LINK	05K FOCS	<0.78%, CL = 90%

$\Gamma(K^+K^*(892)^0, K^{*0} \rightarrow K^+\pi^-)/\Gamma(2K^+\pi^-)$ $\Gamma_{200}/\Gamma_{199}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.47 ± 0.22 ± 0.15	DEL-AMO-SA..11G	BABR	$e^+e^- \approx \Upsilon(4S)$

$\Gamma(2K^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{201}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.7 × 10⁻⁴	90	ABLIKIM	24AV BES3	e^+e^- at 4.128–4.266 GeV

———— Baryon-antibaryon mode ————

$\Gamma(p\bar{n})/\Gamma_{\text{total}}$ Γ_{202}/Γ

This is the only baryonic mode allowed kinematically.

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.22 ± 0.11 OUR AVERAGE				
1.21 ± 0.10 ± 0.05	193 ± 17	ABLIKIM	190BES3	e^+e^- , $E_{\text{cm}} = 4178$ MeV
1.30 ± 0.36 $^{+0.12}_{-0.16}$	13.0 ± 3.6	ATHAR	08 CLEO	e^+e^- , $E_{\text{cm}} \approx 4170$ MeV

$\Gamma(\rho\bar{\rho}e^+\nu_e)/\Gamma_{\text{total}}$					Γ_{203}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<2.0 \times 10^{-4}$	90	ABLIKIM	19BD BES3	e^+e^- at 4178 MeV	

————— Rare or forbidden modes —————

$\Gamma(\pi^+e^+e^-)/\Gamma_{\text{total}}$					Γ_{204}/Γ
This mode is not a useful test for a $\Delta C=1$ weak neutral current because both quarks must change flavor in this decay.					

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$< 5.5 \times 10^{-6}$	90	AAIJ	21T LHCb	$1.6 \text{ fb}^{-1} pp$	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
$<13 \times 10^{-6}$	90	LEES	11G BABR	$e^+e^- \approx \Upsilon(4S)$	
$< 2.2 \times 10^{-5}$	90	¹ RUBIN	10 CLEO	e^+e^- at 4170 MeV	
$<27 \times 10^{-5}$	90	AITALA	99G E791	$\pi^- N$ 500 GeV	

¹This RUBIN 10 limit is for the e^+e^- mass in the continuum away from the $\phi(1020)$. See the next data block.

$\Gamma(\pi^+\phi, \phi \rightarrow e^+e^-)/\Gamma_{\text{total}}$					Γ_{205}/Γ
This is <i>not</i> a test for the $\Delta C = 1$ weak neutral current, but leads to the $\pi^+e^+e^-$ final state.					

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT	
$1.17^{+0.23}_{-0.21} \pm 0.03$	38	ABLIKIM	24CF BES3	e^+e^- at 4.128–4.226 GeV	

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

$0.6^{+0.8}_{-0.4} \pm 0.1$	3	RUBIN	10 CLEO	e^+e^- at 4.170 GeV	
-----------------------------	---	-------	---------	-----------------------	--

$\Gamma(\pi^+\pi^0e^+e^-)/\Gamma_{\text{total}}$					Γ_{206}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<7.0 \times 10^{-5}$	90	ABLIKIM	24CF BES3	e^+e^- at 4.128–4.226 GeV	

$\Gamma(\rho^+\phi, \phi \rightarrow e^+e^-)/\Gamma_{\text{total}}$					Γ_{207}/Γ
VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT	
$2.44^{+0.67}_{-0.62} \pm 0.16$	38	ABLIKIM	24CF BES3	e^+e^- at 4.128–4.226 GeV	

$\Gamma(\pi^+\mu^+\mu^-)/\Gamma_{\text{total}}$					Γ_{208}/Γ
This mode is not a useful test for a $\Delta C=1$ weak neutral current because both quarks must change flavor in this decay.					

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.8 \times 10^{-7}$	90	AAIJ	21T LHCb	$1.6 \text{ fb}^{-1} pp$	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
$<4.1 \times 10^{-7}$	90	AAIJ	13AF LHCb	pp at 7 TeV	
$<4.3 \times 10^{-5}$	90	LEES	11G BABR	$e^+e^- \approx \Upsilon(4S)$	
$<2.6 \times 10^{-5}$	90	LINK	03F FOCS	$\gamma A, \bar{E}_\gamma \approx 180 \text{ GeV}$	
$<1.4 \times 10^{-4}$	90	AITALA	99G E791	$\pi^- N$ 500 GeV	
$<4.3 \times 10^{-4}$	90	KODAMA	95 E653	π^- emulsion 600 GeV	

$\Gamma(K^+ e^+ e^-)/\Gamma_{\text{total}}$ Γ_{209}/Γ

A test for the $\Delta C=1$ weak neutral current. Allowed by higher-order electroweak interactions.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<3.7 \times 10^{-6}$	90	LEES	11G BABR	$e^+ e^- \approx \Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$<4.9 \times 10^{-6}$	90	AAIJ	21T LHCB	$1.6 \text{ fb}^{-1} pp$
$<5.2 \times 10^{-5}$	90	RUBIN	10 CLEO	$e^+ e^-$ at 4170 MeV
$<1.6 \times 10^{-3}$	90	AITALA	99G E791	$\pi^- N$ 500 GeV

$\Gamma(K^+ \pi^0 e^+ e^-)/\Gamma_{\text{total}}$ Γ_{210}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<7.1 \times 10^{-5}$	90	ABLIKIM	24CF BES3	$e^+ e^-$ at 4.128–4.226 GeV

$\Gamma(K_S^0 \pi^+ e^+ e^-)/\Gamma_{\text{total}}$ Γ_{211}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<8.1 \times 10^{-5}$	90	ABLIKIM	24CF BES3	$e^+ e^-$ at 4.128–4.226 GeV

$\Gamma(K^+ \mu^+ \mu^-)/\Gamma_{\text{total}}$ Γ_{212}/Γ

A test for the $\Delta C=1$ weak neutral current. Allowed by higher-order electroweak interactions.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<1.4 \times 10^{-7}$	90	AAIJ	21T LHCB	$1.6 \text{ fb}^{-1} pp$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$<21 \times 10^{-6}$	90	LEES	11G BABR	$e^+ e^- \approx \Upsilon(4S)$
$<3.6 \times 10^{-5}$	90	LINK	03F FOCS	$\gamma A, \bar{E}_\gamma \approx 180 \text{ GeV}$
$<1.4 \times 10^{-4}$	90	AITALA	99G E791	$\pi^- N$ 500 GeV
$<5.9 \times 10^{-4}$	90	KODAMA	95 E653	π^- emulsion 600 GeV

$\Gamma(K^*(892)^+ \mu^+ \mu^-)/\Gamma_{\text{total}}$ Γ_{213}/Γ

A test for the $\Delta C=1$ weak neutral current. Allowed by higher-order electroweak interactions.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<1.4 \times 10^{-3}$	90	KODAMA	95 E653	π^- emulsion 600 GeV

$\Gamma(\pi^+ e^+ \mu^-)/\Gamma_{\text{total}}$ Γ_{214}/Γ

A test of lepton-family-number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<1.1 \times 10^{-6}$	90	AAIJ	21T LHCB	$1.6 \text{ fb}^{-1} pp$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$<12 \times 10^{-6}$	90	LEES	11G BABR	$e^+ e^- \approx \Upsilon(4S)$

$\Gamma(\pi^+ e^- \mu^+)/\Gamma_{\text{total}}$ Γ_{215}/Γ

A test of lepton-family-number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<9.4 \times 10^{-7}$	90	AAIJ	21T LHCB	$1.6 \text{ fb}^{-1} pp$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$<20 \times 10^{-6}$	90	LEES	11G BABR	$e^+ e^- \approx \Upsilon(4S)$

$\Gamma(K^+ e^+ \mu^-)/\Gamma_{\text{total}}$ Γ_{216}/Γ

A test of lepton-family-number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 7.9 \times 10^{-7}$	90	AAIJ	21T LHCb	$1.6 \text{ fb}^{-1} pp$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$< 14 \times 10^{-6}$	90	LEES	11G BABR	$e^+ e^- \approx \Upsilon(4S)$

 $\Gamma(K^+ e^- \mu^+)/\Gamma_{\text{total}}$ Γ_{217}/Γ

A test of lepton-family-number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 5.6 \times 10^{-7}$	90	AAIJ	21T LHCb	$1.6 \text{ fb}^{-1} pp$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$< 9.7 \times 10^{-6}$	90	LEES	11G BABR	$e^+ e^- \approx \Upsilon(4S)$

 $\Gamma(\pi^- 2e^+)/\Gamma_{\text{total}}$ Γ_{218}/Γ

A test of lepton-number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 1.4 \times 10^{-6}$	90	AAIJ	21T LHCb	$1.6 \text{ fb}^{-1} pp$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$< 4.1 \times 10^{-6}$	90	LEES	11G BABR	$e^+ e^- \approx \Upsilon(4S)$
$< 1.8 \times 10^{-5}$	90	RUBIN	10 CLEO	$e^+ e^-$ at 4170 MeV
$< 69 \times 10^{-5}$	90	AITALA	99G E791	$\pi^- N$ 500 GeV

 $\Gamma(\pi^- 2\mu^+)/\Gamma_{\text{total}}$ Γ_{219}/Γ

A test of lepton-number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 8.6 \times 10^{-8}$	90	AAIJ	21T LHCb	$1.6 \text{ fb}^{-1} pp$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$< 1.2 \times 10^{-7}$	90	AAIJ	13AF LHCb	pp at 7 TeV
$< 1.4 \times 10^{-5}$	90	LEES	11G BABR	$e^+ e^- \approx \Upsilon(4S)$
$< 2.9 \times 10^{-5}$	90	LINK	03F FOCS	$\gamma A, \bar{E}_\gamma \approx 180 \text{ GeV}$
$< 8.2 \times 10^{-5}$	90	AITALA	99G E791	$\pi^- N$ 500 GeV
$< 4.3 \times 10^{-4}$	90	KODAMA	95 E653	π^- emulsion 600 GeV

 $\Gamma(\pi^- e^+ \mu^+)/\Gamma_{\text{total}}$ Γ_{220}/Γ

A test of lepton-number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 6.3 \times 10^{-7}$	90	AAIJ	21T LHCb	$1.6 \text{ fb}^{-1} pp$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$< 8.4 \times 10^{-6}$	90	LEES	11G BABR	$e^+ e^- \approx \Upsilon(4S)$
$< 7.3 \times 10^{-4}$	90	AITALA	99G E791	$\pi^- N$ 500 GeV

 $\Gamma(K^- 2e^+)/\Gamma_{\text{total}}$ Γ_{221}/Γ

A test of lepton-number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 7.7 \times 10^{-7}$	90	AAIJ	21T LHCb	$1.6 \text{ fb}^{-1} pp$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$< 5.2 \times 10^{-6}$	90	LEES	11G BABR	$e^+ e^- \approx \Upsilon(4S)$
$< 1.7 \times 10^{-5}$	90	RUBIN	10 CLEO	$e^+ e^-$ at 4170 MeV
$< 63 \times 10^{-5}$	90	AITALA	99G E791	$\pi^- N$ 500 GeV

$\Gamma(\pi^- \pi^0 e^+ e^+)/\Gamma_{\text{total}}$					Γ_{222}/Γ
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$<2.9 \times 10^{-5}$	90	ABLIKIM	25BC BES3	$e^+ e^-$ at 4.128–4.266 GeV	
$\Gamma(K^- \pi^0 e^+ e^+)/\Gamma_{\text{total}}$					Γ_{223}/Γ
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$<3.4 \times 10^{-5}$	90	ABLIKIM	25BC BES3	$e^+ e^-$ at 4.128–4.266 GeV	
$\Gamma(K_S^0 \pi^- e^+ e^+)/\Gamma_{\text{total}}$					Γ_{224}/Γ
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$<1.3 \times 10^{-5}$	90	ABLIKIM	25BC BES3	$e^+ e^-$ at 4.128–4.266 GeV	
$\Gamma(K_S^0 K^- e^+ e^+)/\Gamma_{\text{total}}$					Γ_{225}/Γ
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$<2.9 \times 10^{-5}$	90	ABLIKIM	25BC BES3	$e^+ e^-$ at 4.128–4.266 GeV	
$\Gamma(\phi \pi^- e^+ e^+)/\Gamma_{\text{total}}$					Γ_{226}/Γ
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$<6.9 \times 10^{-5}$	90	ABLIKIM	25BC BES3	$e^+ e^-$ at 4.128–4.266 GeV	
$\Gamma(\phi K^- e^+ e^+)/\Gamma_{\text{total}}$					Γ_{227}/Γ
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$<9.9 \times 10^{-5}$	90	ABLIKIM	25BC BES3	$e^+ e^-$ at 4.128–4.266 GeV	
$\Gamma(K^- 2\mu^+)/\Gamma_{\text{total}}$					Γ_{228}/Γ
A test of lepton-number conservation.					
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$<2.6 \times 10^{-8}$	90	AAIJ	21T LHCB	$1.6 \text{ fb}^{-1} pp$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<1.3 \times 10^{-5}$	90	LEES	11G BABR	$e^+ e^- \approx \Upsilon(4S)$	
$<1.3 \times 10^{-5}$	90	LINK	03F FOCS	$\gamma A, \bar{E}_\gamma \approx 180 \text{ GeV}$	
$<1.8 \times 10^{-4}$	90	AITALA	99G E791	$\pi^- N 500 \text{ GeV}$	
$<5.9 \times 10^{-4}$	90	KODAMA	95 E653	π^- emulsion 600 GeV	
$\Gamma(K^- e^+ \mu^+)/\Gamma_{\text{total}}$					Γ_{229}/Γ
A test of lepton-number conservation.					
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$<2.6 \times 10^{-7}$	90	AAIJ	21T LHCB	$1.6 \text{ fb}^{-1} pp$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<6.1 \times 10^{-6}$	90	LEES	11G BABR	$e^+ e^- \approx \Upsilon(4S)$	
$<6.8 \times 10^{-4}$	90	AITALA	99G E791	$\pi^- N 500 \text{ GeV}$	
$\Gamma(K^*(892)^- 2\mu^+)/\Gamma_{\text{total}}$					Γ_{230}/Γ
A test of lepton-number conservation.					
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$<1.4 \times 10^{-3}$	90	KODAMA	95 E653	π^- emulsion 600 GeV	

D_s^\pm Amplitude analyses

$D_s^+ \rightarrow K^+ K^- \pi^+$ partial wave analyses

Amplitude analyses of D_s^+ decays to the $K^+ K^- \pi^+$ final state, fitting simultaneously different partial wave components.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	18.6k	¹ ABLIKIM	21AE BES3	$e^+ e^-$ at 4.178 GeV
seen	96k	¹ DEL-AMO-SA...11G	BABR	$e^+ e^-$ at $\Upsilon(4S)$
seen	12k	¹ MITCHELL	09A CLEO	$e^+ e^-$ at 4.17 GeV
seen	701	² FRABETTI	95B E687	

¹ Amplitude analysis with 6 components.

² Amplitude analysis with 5 components.

$D_s^+ \rightarrow K^+ K_S \pi^0$ partial wave analyses

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
	990	¹ ABLIKIM	22AH BES3	$e^+ e^-$ at 4.178-4.226 GeV

¹ Amplitude analysis with 5 components.

$D_s^+ \rightarrow 2\pi^+ \pi^-$ partial wave analyses

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
	0.7M	¹ AAIJ	23AN LHCb	Dalitz fit, 0.7M events
	11.1k	² ABLIKIM	22BI BES3	Dalitz fit
	10.5k	² AUBERT	09O BABR	Dalitz fit
	1.5k	³ LINK	04 FOCS	Dalitz fit
	848	⁴ AITALA	01A E791	Dalitz fit

¹ Amplitude analysis with 7 components, one of which is a model-independent $\pi^+ \pi^-$ S-wave parametrisation as complex numbers in 50 $\pi^+ \pi^-$ mass bins.

² Amplitude analysis with 4 components, one of which is a model-independent $\pi^+ \pi^-$ S-wave parametrisation as complex numbers in 29 $\pi^+ \pi^-$ mass bins.

³ Amplitude analysis with 5 components.

⁴ Amplitude analysis with 6 components.

$D_s^+ \rightarrow 2\pi^+ \pi^- \pi^0$ partial wave analyses

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
	2.5k	¹ ABLIKIM	25A BES3	$e^+ e^-$ at 4.128-4.226 GeV

¹ Amplitude analysis with 11 components contributing to the $2\pi^+ \pi^- \pi^0_{non-\eta}$ final state.

$D_s^+ \rightarrow 2\pi^+ \pi^- \eta$ partial wave analyses

Amplitude analyses of D_s^+ decays to the $\pi^+ \pi^+ \pi^- \eta$ final state, fitting simultaneously different partial wave components.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	2.1k	¹ ABLIKIM	21AR BES3	$e^+ e^-$ at 4.178-4.226 GeV

¹ Amplitude analysis with 11 components.

$D_s^+ \rightarrow \pi^+ \pi^0 \eta'$ partial wave analyses.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
	395	¹ ABLIKIM	22AA BES3	$e^+ e^-$ at 4.178-4.226 GeV

¹ The only significant contribution found in this analysis is $D_s^+ \rightarrow \rho^+ \eta'$.

$D_s^+ \rightarrow \pi^+ 2\pi^0$ partial wave analyses.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
	440	ABLIKIM	22Z BES3	$e^+ e^-$ at 4.178–4.226 GeV

$D_s^+ \rightarrow 2\pi^+ \pi^- 2\pi^0$ partial wave analyses

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
	2k	¹ ABLIKIM	25F BES3	$e^+ e^-$ at 4.128–4.226 GeV

¹ Amplitude analysis with 14 components.

$D_s^+ \rightarrow K^+ \pi^+ \pi^-$ partial wave analyses

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
	1.3k	¹ ABLIKIM	22AC BES3	$e^+ e^-$ at 4.178–4.226 GeV

¹ ABLIKIM 22AC uses an amplitude analysis with 8 components.

$D_s^+ \rightarrow K_S^0 \pi^+ \pi^0$ partial wave analyses

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
	666	¹ ABLIKIM	21AB BES3	$e^+ e^-$ at 4.178–4.226 GeV

¹ ABLIKIM 21AB uses an amplitude analysis with 5 resonant modes plus one background component.

$D_s^+ \rightarrow K^+ \pi^+ \pi^- \pi^0$ partial wave analyses

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
seen	550	¹ ABLIKIM	22BL BES3	$e^+ e^-$ at 4.178–4.226 GeV

¹ Amplitude analysis with 11 components.

$D_s^+ \rightarrow 2K_S^0 \pi^+$ partial wave analyses

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
	400	¹ ABLIKIM	22F BES3	$e^+ e^-$ at 4.178–4.226 GeV

¹ Amplitude analysis with 2 components.

$D_s^+ \rightarrow (KS)^0 K^- 2\pi^+$ partial wave analyses

Amplitude analyses of D_s^+ decays to the $(KS)^0 K^- 2\pi^+$ final state, fitting simultaneously different partial wave components.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
seen	1.3k	¹ ABLIKIM	21K BES3	$e^+ e^-$ at 4.178–4.226 GeV

¹ Amplitude analysis with 13 components.

$D_s^+ \rightarrow K^- K^+ \pi^+ \pi^0$ partial wave analyses

Amplitude analyses of D_s^+ decays to the $K^- K^+ \pi^+ \pi^0$ final state, fitting simultaneously different partial wave components.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
seen	3k	¹ ABLIKIM	21U BES3	$e^+ e^-$ at 4.178–4.226 GeV

¹ ABLIKIM 21U uses an amplitude analysis with 9 components.

$D_s^+ \rightarrow K^- K^+ 2\pi^+ \pi^-$ partial wave analyses

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
	309	ABLIKIM	22AB BES3	$e^+ e^-$ at 4.178–4.226 GeV

$D_s^+ \rightarrow K_S^0 K_L^0 \pi^+$ partial wave analyses

VALUE	EVTS	DOCUMENT ID
	2.35k	¹ ABLIKIM 25BR

¹ ABLIKIM 25BR amplitude analysis with 4 resonant modes plus one background component measures the asymmetry of the branching fractions of $D_s^+ \rightarrow K_S^0 K^*(892)^+$ and $D_s^+ \rightarrow K_L^0 K^*(892)^+$.

$D_s^+ - D_s^-$ CP-VIOLATING DECAY-RATE ASYMMETRIES

This is the difference between D_s^+ and D_s^- partial widths for the decay to state f , divided by the sum of the widths:

$$A_{CP}(f) = [\Gamma(D_s^+ \rightarrow f) - \Gamma(D_s^- \rightarrow \bar{f})] / [\Gamma(D_s^+ \rightarrow f) + \Gamma(D_s^- \rightarrow \bar{f})].$$

$A_{CP}(\mu^\pm \nu)$ in $D_s^+ \rightarrow \mu^+ \nu$, $D_s^- \rightarrow \mu^- \bar{\nu}_\mu$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-0.2 ± 2.5 OUR AVERAGE				
-1.2 ± 2.5 ± 1.0	2.2k	ABLIKIM	21BE BES3	$e^+ e^-$ at 4.178, 4.226 GeV
4.8 ± 6.1		ALEXANDER	09 CLEO	$e^+ e^-$ at 4170 MeV

$A_{CP}(\tau^\pm \nu)$ in $D_s^+ \rightarrow \tau^+ \nu_\tau$, $D_s^- \rightarrow \tau^- \bar{\nu}_\tau$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
2.9 ± 4.8 ± 1.0	950	¹ ABLIKIM	21BE BES3	$e^+ e^-$ at 4.178, 4.226 GeV

¹ ABLIKIM 21BE also reports that when constrained by the Standard Model ratio of $\Gamma(D_s^+ \rightarrow \tau^+ \nu_\tau) / \Gamma(D_s^+ \rightarrow \mu^+ \nu_\mu) = 9.75$, the result is $(-0.1 \pm 1.9 \pm 1.0)\%$.

$A_{CP}(K^\pm K_S^0)$ in $D_s^\pm \rightarrow K^\pm K_S^0$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.07 ± 0.24 OUR AVERAGE				
0.29 ± 0.50 ± 0.21		ABLIKIM	24BD BES3	$e^+ e^-$, 4.128–4.226 GeV
-0.05 ± 0.23 ± 0.24	288k	¹ LEES	13E BABR	$e^+ e^-$ at $\Upsilon(4S)$
0.12 ± 0.36 ± 0.22		KO	10 BELL	$e^+ e^- \approx \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.6 ± 2.8 ± 0.6	1.8k	² ABLIKIM	19AMBES3	$e^+ e^-$ at 4178 MeV
2.6 ± 1.5 ± 0.6		ONYISI	13 CLEO	$e^+ e^-$ at 4.17 GeV
4.7 ± 1.8 ± 0.9	4.0k	MENDEZ	10 CLEO	See ONYISI 13
4.9 ± 2.1 ± 0.9		ALEXANDER	08 CLEO	See MENDEZ 10

¹ LEES 13E finds that after subtracting the contribution due to $K^0 - \bar{K}^0$ mixing, the CP asymmetry is $(+0.28 \pm 0.23 \pm 0.24)\%$.

² Superseded by ABLIKIM 24BD.

$A_{CP}(K^\pm K_L^0)$ in $D_s^\pm \rightarrow K^\pm K_L^0$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
-1.1 ± 2.6 ± 0.6	2.3k	ABLIKIM	19AMBES3	$e^+ e^-$ at 4178 MeV

$A_{CP}(K^+ K^- \pi^\pm)$ in $D_s^\pm \rightarrow K^+ K^- \pi^\pm$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
0.35 ± 0.34 OUR AVERAGE			
0.48 ± 0.26 ± 0.24	ABLIKIM	24BD BES3	$e^+ e^-$, 4.128–4.226 GeV

−0.5 ±0.8 ±0.4 ONYISI 13 CLEO e^+e^- at 4.17 GeV

••• We do not use the following data for averages, fits, limits, etc. •••

0.3 ±1.1 ±0.8 ALEXANDER 08 CLEO See ONYISI 13

$A_{CP}(\phi\pi^\pm)$ in $D_s^\pm \rightarrow \phi\pi^\pm$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
−0.38±0.26±0.08	ABAZOV	14B	D0 $p\bar{p}$ at 1.96 TeV

$A_{CP}(K^\pm K_S^0\pi^0)$ in $D_s^\pm \rightarrow K^\pm K_S^0\pi^0$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
−0.9 ±1.9 OUR AVERAGE			
−0.85±1.97±0.46	ABLIKIM	24BD	BES3 e^+e^- , 4.128–4.226 GeV
−1.6 ±6.0 ±1.1	ONYISI	13	CLEO e^+e^- at 4.17 GeV

$A_{CP}(2K_S^0\pi^\pm)$ in $D_s^\pm \rightarrow 2K_S^0\pi^\pm$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
1.3 ±1.6 OUR AVERAGE			
1.14±1.58±0.44	ABLIKIM	24BD	BES3 e^+e^- , 4.128–4.226 GeV
3.1 ±5.2 ±0.6	ONYISI	13	CLEO e^+e^- at 4.17 GeV

$A_{CP}(K^+K^-\pi^\pm\pi^0)$ in $D_s^\pm \rightarrow K^+K^-\pi^\pm\pi^0$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
−0.6 ±0.9 OUR AVERAGE			
−0.66±0.91±0.33	ABLIKIM	24BD	BES3 e^+e^- , 4.128–4.226 GeV
0.0 ±2.7 ±1.2	ONYISI	13	CLEO e^+e^- at 4.17 GeV
••• We do not use the following data for averages, fits, limits, etc. •••			
−5.9 ±4.2 ±1.2	ALEXANDER	08	CLEO See ONYISI 13

$A_{CP}(K^\pm K_S^0\pi^+\pi^-)$ in $D_s^\pm \rightarrow K^\pm K_S^0\pi^+\pi^-$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
0.7 ±2.9 OUR AVERAGE			Error includes scale factor of 1.3.
2.00±2.37±0.70	ABLIKIM	24BD	BES3 e^+e^- , 4.128–4.226 GeV
−5.7 ±5.3 ±0.9	ONYISI	13	CLEO e^+e^- at 4.17 GeV

$A_{CP}(K_S^0 K^\mp 2\pi^\pm)$ in $D_s^\pm \rightarrow K_S^0 K^\mp 2\pi^\pm$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
0.7 ±1.8 OUR AVERAGE			Error includes scale factor of 1.3.
−0.24±1.05±1.07	ABLIKIM	24BD	BES3 e^+e^- , 4.128–4.226 GeV
4.1 ±2.7 ±0.9	ONYISI	13	CLEO e^+e^- at 4.17 GeV
••• We do not use the following data for averages, fits, limits, etc. •••			
−0.7 ±3.6 ±1.1	ALEXANDER	08	CLEO See ONYISI 13

$A_{CP}(\pi^+\pi^-\pi^\pm)$ in $D_s^\pm \rightarrow \pi^+\pi^-\pi^\pm$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
−0.9 ±1.1 OUR AVERAGE			
−0.88±1.17±0.38	ABLIKIM	24BD	BES3 e^+e^- , 4.128–4.226 GeV
−0.7 ±3.0 ±0.6	ONYISI	13	CLEO e^+e^- at 4.17 GeV
••• We do not use the following data for averages, fits, limits, etc. •••			
2.0 ±4.6 ±0.7	ALEXANDER	08	CLEO See ONYISI 13

$A_{CP}(\pi^\pm \eta)$ in $D_s^\pm \rightarrow \pi^\pm \eta$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.24 ± 0.29 OUR AVERAGE				
−0.44 ± 0.89 ± 0.19		ABLIKIM	24BD BES3	$e^+ e^-$, 4.128–4.226 GeV
0.32 ± 0.51 ± 0.12	136k	AAIJ	23E LHCb	6 fb^{-1} , pp at 13 TeV, $\eta \rightarrow \gamma \pi \pi$
0.8 ± 0.7 ± 0.5	38k	AAIJ	21U LHCb	pp at 13 TeV
0.2 ± 0.3 ± 0.3	22k	GUAN	21 BELL	$e^+ e^- \approx \Upsilon(4,5S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.1 ± 3.0 ± 0.8		ONYISI	13 CLEO	$e^+ e^-$ at 4.17 GeV
−4.6 ± 2.9 ± 0.3	2.5k	MENDEZ	10 CLEO	See ONYISI 13
−8.2 ± 5.2 ± 0.8		ALEXANDER	08 CLEO	See MENDEZ 10

 $A_{CP}(\pi^\pm \pi^+ \pi^- \eta)$ in $D_s^\pm \rightarrow \pi^\pm \pi^+ \pi^- \eta$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
2.42 ± 2.85 ± 0.78	ABLIKIM	24BD BES3	$e^+ e^-$, 4.128–4.226 GeV

 $A_{CP}(\pi^\pm \eta')$ in $D_s^\pm \rightarrow \pi^\pm \eta'$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
−0.08 ± 0.17 OUR AVERAGE Error includes scale factor of 1.2.				
−0.59 ± 0.76 ± 0.20		ABLIKIM	24BD BES3	$e^+ e^-$, 4.128–4.226 GeV
0.01 ± 0.12 ± 0.08	1M	AAIJ	23E LHCb	6 fb^{-1} , pp at 13 TeV, $\eta \rightarrow \gamma \pi \pi$
−0.82 ± 0.36 ± 0.35	152k	AAIJ	17AF LHCb	pp at 7, 8 TeV
−2.2 ± 2.2 ± 0.6		ONYISI	13 CLEO	$e^+ e^-$ at 4.17 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
−6.1 ± 3.0 ± 0.3	1.4k	MENDEZ	10 CLEO	See ONYISI 13
−5.5 ± 3.7 ± 1.2		ALEXANDER	08 CLEO	See MENDEZ 10

 $A_{CP}(\eta \pi^\pm \pi^0)$ in $D_s^\pm \rightarrow \eta \pi^\pm \pi^0$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
0.9 ± 1.5 OUR AVERAGE			
1.05 ± 1.45 ± 0.62	ABLIKIM	24BD BES3	$e^+ e^-$, 4.128–4.226 GeV
−0.5 ± 3.9 ± 2.0	ONYISI	13 CLEO	$e^+ e^-$ at 4.17 GeV

 $A_{CP}(\eta' \pi^\pm \pi^0)$ in $D_s^\pm \rightarrow \eta' \pi^\pm \pi^0$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
−1.5 ± 2.5 OUR AVERAGE			
−1.60 ± 2.57 ± 0.64	ABLIKIM	24BD BES3	$e^+ e^-$, 4.128–4.226 GeV
−0.4 ± 7.4 ± 1.9	ONYISI	13 CLEO	$e^+ e^-$ at 4.17 GeV

 $A_{CP}(K^\pm \pi^0)$ in $D_s^\pm \rightarrow K^\pm \pi^0$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
2 ± 4 OUR AVERAGE Error includes scale factor of 1.2.				
−0.8 ± 3.9 ± 1.2	2.8k	AAIJ	21U LHCb	pp at 7, 8, 13 TeV
6.4 ± 4.4 ± 1.1	12k	GUAN	21 BELL	$e^+ e^- \approx \Upsilon(4,5S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
−26.6 ± 23.8 ± 0.9	202	MENDEZ	10 CLEO	$e^+ e^-$ at 4170 MeV
2 ± 29		ADAMS	07A CLEO	See MENDEZ 10

$A_{CP}(\bar{K}^0/K^0\pi^\pm)$ in $D_s^+ \rightarrow \bar{K}^0\pi^+$, $D_s^- \rightarrow K^0\pi^-$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

0.4 ± 0.5 OUR AVERAGE

0.38 ± 0.46 ± 0.17	121k	¹ AAIJ	14BD LHCb	pp at 7, 8 TeV
0.3 ± 2.0 ± 0.3	14k	LEES	13E BABR	e^+e^- at $\Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.61 ± 0.83 ± 0.14	26k	AAIJ	13W LHCb	See AAIJ 14BD
--------------------	-----	------	----------	---------------

¹AAIJ 14BD reports its result as $A_{CP}(D_s^\pm \rightarrow K_S^0 K^\pm)$ with CP -violation effects in the $K^0 - \bar{K}^0$ system subtracted. It also measures $A_{CP}(D^\pm \rightarrow \bar{K}^0/K^0 K^\pm) + A_{CP}(D_s^\pm \rightarrow \bar{K}^0/K^0 \pi^\pm) = (0.41 \pm 0.49 \pm 0.26)\%$.

$A_{CP}(K_S^0\pi^\pm)$ in $D_s^\pm \rightarrow K_S^0\pi^\pm$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

0.20 ± 0.18 OUR AVERAGE

0.16 ± 0.17 ± 0.05	721k	AAIJ	19T LHCb	pp at 7, 8, 13 TeV
0.6 ± 2.0 ± 0.3	14k	LEES	13E BABR	e^+e^- at $\Upsilon(4S)$
5.45 ± 2.50 ± 0.33		KO	10 BELL	$e^+e^- \approx \Upsilon(4S)$
16.3 ± 7.3 ± 0.3	0.4k	MENDEZ	10 CLEO	e^+e^- at 4170 MeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

27 ± 11		ADAMS	07A CLEO	See MENDEZ 10
---------	--	-------	----------	---------------

$A_{CP}(K^\pm\pi^+\pi^-)$ in $D_s^\pm \rightarrow K^\pm\pi^+\pi^-$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

2.2 ± 1.9 OUR AVERAGE

1.81 ± 2.01 ± 0.45		ABLIKIM	24BD BES3	e^+e^- , 4.128–4.226 GeV
4.5 ± 4.8 ± 0.6		ONYISI	13 CLEO	e^+e^- at 4.17 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.3 ± 3.0 ± 1.3	1.3k	¹ ABLIKIM	22AC BES3	e^+e^- , 4.178–4.226 GeV
11.2 ± 7.0 ± 0.9		ALEXANDER	08 CLEO	See ONYISI 13

¹Superseded by ABLIKIM 24BD.

$A_{CP}(K_S^0\pi^+\pi^0)$ in $D_s^\pm \rightarrow K_S^0\pi^\pm\pi^0$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

−2.17 ± 4.65 ± 1.10		ABLIKIM	24BD BES3	e^+e^- , 4.128–4.226 GeV
----------------------------	--	---------	-----------	----------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.7 ± 5.5 ± 0.9	666	¹ ABLIKIM	21AB BES3	e^+e^- , 4.178–4.226 GeV
-----------------	-----	----------------------	-----------	----------------------------

¹ABLIKIM 21AB uses an amplitude analysis with 5 resonant modes plus one background component. Superseded by ABLIKIM 24BD.

$A_{CP}(K^\pm\pi^+\pi^-\pi^0)$ in $D_s^\pm \rightarrow K^\pm\pi^+\pi^-\pi^0$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

6.6 ± 5.4 ± 0.7	776	ABLIKIM	22BL BES3	e^+e^- at 4.178–4.226 GeV
------------------------	-----	---------	-----------	-----------------------------

$A_{CP}(K^\pm\eta)$ in $D_s^\pm \rightarrow K^\pm\eta$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

1.8 ± 1.9 OUR AVERAGE

0.9 ± 3.7 ± 1.1	2.5k	AAIJ	21U LHCb	pp at 13 TeV
-----------------	------	------	----------	----------------

$2.1 \pm 2.1 \pm 0.4$	14k	GUAN	21	BELL	$e^+e^- \approx \Upsilon(4,5S)$
••• We do not use the following data for averages, fits, limits, etc. •••					
$9.3 \pm 15.2 \pm 0.9$	222	MENDEZ	10	CLEO	e^+e^- at 4170 MeV
-20 ± 18		ADAMS	07A	CLEO	See MENDEZ 10

$A_{CP}(K^\pm \eta'(958))$ in $D_s^\pm \rightarrow K^\pm \eta'(958)$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$6.0 \pm 18.9 \pm 0.9$	56 ± 17	MENDEZ	10	CLEO e^+e^- at 4170 MeV
••• We do not use the following data for averages, fits, limits, etc. •••				
-17 ± 37		ADAMS	07A	CLEO See MENDEZ 10

$D_s^\pm \chi^2$ TESTS OF CP-VIOLATION (CPV)

We list model-independent searches for local CP violation in phase-space distributions of multi-body decays.

Most of these searches divide phase space (Dalitz plot for 3-body decays, five-dimensional equivalent for 4-body decays) into bins, and perform a χ^2 test comparing normalised yields N_i, \bar{N}_i in CP -conjugate bin pairs i : $\chi^2 = \sum_i (N_i - \alpha \bar{N}_i) / \sigma(N_i - \alpha \bar{N}_i)$. The factor $\alpha = (\sum_i N_i) / (\sum_i \bar{N}_i)$ removes the dependence on phase-space-integrated rate asymmetries. The result is used to obtain the probability (p-value) to obtain the measured χ^2 or larger under the assumption of CP conservation [AUBERT 08AO, BEDIAGA 09]. Alternative methods obtain p-values from other test variables based on unbinned analyses [WILLIAMS 11, AAIJ 14C]. Results can be combined using Fisher's method [MOSTELLER 48].

Local CPV in $D_s^\pm \rightarrow K^+ K^- K^\pm$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.133	970k	AAIJ	23L	LHCB χ^2

CP VIOLATING ASYMMETRIES OF P-ODD (T-ODD) MOMENTS

$A_{Tviol}(K_S^0 K^\pm \pi^+ \pi^-)$ in $D_s^\pm \rightarrow K_S^0 K^\pm \pi^+ \pi^-$

$C_T \equiv \vec{p}_{K^+} \cdot (\vec{p}_{\pi^+} \times \vec{p}_{\pi^-})$ is a parity-odd correlation of the K^+ , π^+ , and π^- momenta for the D_s^+ . $\bar{C}_T \equiv \vec{p}_{K^-} \cdot (\vec{p}_{\pi^-} \times \vec{p}_{\pi^+})$ is the corresponding quantity for the D_s^- . Then

$A_T \equiv [\Gamma(C_T > 0) - \Gamma(C_T < 0)] / [\Gamma(C_T > 0) + \Gamma(C_T < 0)]$, and

$\bar{A}_T \equiv [\Gamma(-\bar{C}_T > 0) - \Gamma(-\bar{C}_T < 0)] / [\Gamma(-\bar{C}_T > 0) + \Gamma(-\bar{C}_T < 0)]$, and

$A_{Tviol} \equiv \frac{1}{2}(A_T - \bar{A}_T)$. C_T and \bar{C}_T are commonly referred to as T -odd moments, because they are odd under T reversal. However, the T -conjugate process $K_S^0 K^\pm \pi^+ \pi^- \rightarrow D_s^\pm$ is not accessible, while the P -conjugate process is.

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
-8 ± 6	OUR AVERAGE			
$-4.6 \pm 6.3 \pm 3.8$	70k	MOON	23	BELL 980 fb $^{-1}$ at $\sim \Upsilon(4S)$
$-13.6 \pm 7.7 \pm 3.4$	29.8k	LEES	11E	BABR $e^+e^- \approx \Upsilon(4S)$
••• We do not use the following data for averages, fits, limits, etc. •••				
$-36 \pm 67 \pm 23$	508	LINK	05E	FOCS $\gamma A, \bar{E}_\gamma \approx 180$ GeV

$A_{T^{viol}}(K_S^0 K^\mp \pi^\pm \pi^\pm)$ in $D_S^\pm \rightarrow K_S^0 K^\mp \pi^\pm \pi^\pm$

$C_T \equiv \vec{p}_{K_S^0} \cdot (\vec{p}_{K^-} \times \vec{p}_{\pi^+})$ is a parity-odd correlation of the momenta of the K_S^0 , K^- , and highest-momenta π^+ for the D_S^+ . $\bar{C}_T \equiv \vec{p}_{K_S^0} \cdot (\vec{p}_{K^+} \times \vec{p}_{\pi^-})$ is the corresponding quantity for the D_S^- . Then

$$A_T \equiv [\Gamma(C_T > 0) - \Gamma(C_T < 0)] / [\Gamma(C_T > 0) + \Gamma(C_T < 0)], \text{ and}$$

$$\bar{A}_T \equiv [\Gamma(-\bar{C}_T > 0) - \Gamma(-\bar{C}_T < 0)] / [\Gamma(-\bar{C}_T > 0) + \Gamma(-\bar{C}_T < 0)], \text{ and}$$

$A_{T^{viol}} \equiv \frac{1}{2}(A_T - \bar{A}_T)$. C_T and \bar{C}_T are commonly referred to as T -odd moments, because they are odd under T reversal. However, the T -conjugate process $K_S^0 K^\pm \pi^+ \pi^- \rightarrow D_S^\pm$ is not accessible, while the P -conjugate process is.

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$0.2 \pm 2.4 \pm 0.8$	303k	¹ AGGARWAL	25 BEL2	$e^+ e^- \approx \Upsilon(4S)$

¹ AGGARWAL 25 measures $C_T \equiv \vec{p}_{K_S^0} \cdot (\vec{p}_{K^-} \times \vec{p}_{\pi^+})$ and other T -odd observables, all compatible with T -conservation, using Belle and Belle II data.

D_S^+ Semileptonic Form Factors and Decay Constants

$r_2 \equiv A_2(0)/A_1(0)$ in $D_S^+ \rightarrow \phi \ell^+ \nu_\ell$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.83 ± 0.08 OUR AVERAGE		Error includes scale factor of 1.8.		See the ideogram below.
$0.71 \pm 0.14 \pm 0.02$		¹ ABLIKIM	23BZ BES3	$D_S^+ \rightarrow K^+ K^- \mu^+ \nu_\mu$
$0.816 \pm 0.036 \pm 0.030$	25k	² AUBERT	08AN BABR	$\phi e^+ \nu_e$
$0.713 \pm 0.202 \pm 0.284$	793	LINK	04C FOCS	$\phi \mu^+ \nu_\mu$
$1.57 \pm 0.25 \pm 0.19$	271	AITALA	99D E791	$\phi e^+ \nu_e, \phi \mu^+ \nu_\mu$
$1.4 \pm 0.5 \pm 0.3$	308	AVERY	94B CLE2	$\phi e^+ \nu_e$
$1.1 \pm 0.8 \pm 0.1$	90	FRABETTI	94F E687	$\phi \mu^+ \nu_\mu$
$2.1 \begin{smallmatrix} +0.6 \\ -0.5 \end{smallmatrix} \pm 0.2$	19	KODAMA	93 E653	$\phi \mu^+ \nu_\mu$

¹ Partial wave analysis of 939 $D_S^+ \rightarrow K^+ K^- \mu^+ \nu_\mu$ events.

² To compare with previous measurements, this AUBERT 08AN value is from a fit that fixes the pole masses at $m_A = 2.5 \text{ GeV}/c^2$ and $m_V = 2.1 \text{ GeV}/c^2$. A simultaneous fit to r_2, r_V, r_0 (a significant s -wave contribution) and m_A , gives $r_2 = 0.763 \pm 0.071 \pm 0.065$.

$f_+(0) |V_{cs}|$ in $D_s^+ \rightarrow \eta e^+ \nu_e$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.449 ± 0.009 OUR AVERAGE

0.430 ± 0.021 ± 0.016	716	¹ ABLIKIM	24CI BES3	$e^+ e^-$ at 4237–4699 MeV
-----------------------	-----	----------------------	-----------	----------------------------

0.4519 ± 0.0071 ± 0.0065	4k	ABLIKIM	23BO BES3	$e^+ e^-$ at 4128–4226 MeV
--------------------------	----	---------	-----------	----------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.4455 ± 0.0053 ± 0.0044	1.8k	² ABLIKIM	19S BES3	$e^+ e^-$ at 4178 MeV
--------------------------	------	----------------------	----------	-----------------------

¹ Using a two parameter fit in the z expansion .

² Superseded by ABLIKIM 23BO

$f_+(0) |V_{cs}|$ in $D_s^+ \rightarrow \eta \mu^+ \nu_\mu$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.452 ± 0.010 ± 0.007	3.1k	ABLIKIM	24AQ BES3	$e^+ e^-$, 4.128–4.266 GeV
------------------------------	------	---------	-----------	-----------------------------

$r_1 \equiv a_1/a_0$ in $D_s^+ \rightarrow \eta \mu^+ \nu_\mu$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

−2.9 ± 0.6 ± 0.2	3.1k	ABLIKIM	24AQ BES3	$e^+ e^-$, 4.128–4.266 GeV
-------------------------	------	---------	-----------	-----------------------------

$\langle A_{FB}^\eta \rangle$ in $D_s^+ \rightarrow \eta \mu^+ \nu_\mu$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

−5.9 ± 3.1 ± 0.5	3.1k	ABLIKIM	24AQ BES3	$e^+ e^-$, 4.128–4.266 GeV
-------------------------	------	---------	-----------	-----------------------------

$f_+(0) |V_{cs}|$ in $D_s^+ \rightarrow \eta' e^+ \nu_e$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.527 ± 0.024 OUR AVERAGE

0.542 ± 0.062 ± 0.023	134	¹ ABLIKIM	24CI BES3	$e^+ e^-$ at 4237–4699 MeV
-----------------------	-----	----------------------	-----------	----------------------------

0.525 ± 0.024 ± 0.009	675	ABLIKIM	23BO BES3	$e^+ e^-$ at 4128–4226 MeV
-----------------------	-----	---------	-----------	----------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.477 ± 0.049 ± 0.011	261	² ABLIKIM	19S BES3	$e^+ e^-$ at 4178 MeV
-----------------------	-----	----------------------	----------	-----------------------

¹ Using a two parameter fit in the z expansion.

² Superseded by ABLIKIM 23BO

$r_1 \equiv a_1/a_0$ in $D_s^+ \rightarrow \eta' \mu^+ \nu_\mu$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

−10.8 ± 5.3 ± 1.4	390	ABLIKIM	24AQ BES3	$e^+ e^-$, 4.128–4.266 GeV
--------------------------	-----	---------	-----------	-----------------------------

$\langle A_{FB}^{\eta'} \rangle$ in $D_s^+ \rightarrow \eta' \mu^+ \nu_\mu$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

−6.4 ± 7.9 ± 0.6	390	ABLIKIM	24AQ BES3	$e^+ e^-$, 4.128–4.266 GeV
-------------------------	-----	---------	-----------	-----------------------------

$f_+(0) |V_{cs}|$ in $D_s^+ \rightarrow \eta' \mu^+ \nu_\mu$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.504 ± 0.037 ± 0.012	390	ABLIKIM	24AQ BES3	$e^+ e^-$, 4.128–4.266 GeV
------------------------------	-----	---------	-----------	-----------------------------

$f_+(0) |V_{cs}|$ in $D_s^+ \rightarrow f_0(980) e^+ \nu_e$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.504 ± 0.017 ± 0.035	0.4k	¹ ABLIKIM	24AT BES3	$e^+ e^-$ at 4128–4226 MeV
------------------------------	------	----------------------	-----------	----------------------------

¹From an analysis of $D_s^+ \rightarrow \pi^+ \pi^- e^+ \mu_e$ decays. ABLIKIM 24AT uses a simple pole parametrization of the hadronic form factor and the Flatte' formula for describing the $f_0(980)$ decay.

$f_+(0)|V_{cd}|$ in $D_s^+ \rightarrow K^0 e^+ \nu_e$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.145±0.010 OUR AVERAGE				
0.143±0.011±0.003	228	¹ ABLIKIM	24CH BES3	$e^+ e^-$ at 4128–4226 MeV
0.152±0.022±0.005	51	¹ ABLIKIM	24CI BES3	$e^+ e^-$ at 4237–4699 MeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.162±0.019±0.003	117	^{1,2} ABLIKIM	19D BES3	$K_S^0 e^+ \nu_e$

¹Using a two parameter fit in the z expansion.

²Superseded by ABLIKIM 24CH.

$r_V \equiv V(0)/A_1(0)$ in $D_s^+ \rightarrow K^*(892)^0 e^+ \nu_e$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
1.67±0.34±0.16	155	ABLIKIM	19D BES3	$e^+ e^-$ at 4178 MeV

$r_2 \equiv A_2(0)/A_1(0)$ in $D_s^+ \rightarrow K^*(892)^0 e^+ \nu_e$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.77±0.28±0.07	155	ABLIKIM	19D BES3	$e^+ e^-$ at 4178 MeV

$f_{D_s^+} |V_{cs}|$ in $D_s^+ \rightarrow \mu^+ \nu_\mu$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
242.7±3.0 OUR AVERAGE				
246.5±5.9±3.6±0.5	0.5k	¹ ABLIKIM	24CG BES3	$e^+ e^-$ at 4.237–4.699 GeV
241.8±2.5±2.2	2.5k	ABLIKIM	23BR BES3	$e^+ e^-$ at 4.128–4.226 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
243.1±3.0±3.6±1.0	2.2K	^{2,3} ABLIKIM	21BE BES3	$e^+ e^-$ at 4.178, 4.226 GeV
246.2±3.6±3.5	1.1k	³ ABLIKIM	19E BES3	$e^+ e^-$ at 4.178 GeV

¹The third uncertainty is due to the uncertainty of the D_s^+ lifetime.

²The third uncertainty is dominated by the uncertainty of the D_s^+ lifetime.

³Superseded by ABLIKIM 23BR.

$f_{D_s^+} |V_{cs}|$ in $D_s^+ \rightarrow \tau^+ \nu_\tau$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
247.5±2.3 OUR AVERAGE				
252.7±3.6±4.5±0.6	2.8k	¹ ABLIKIM	24CG BES3	$e^+ e^-$ at 4.237–4.699 GeV
248.3±3.9±3.1±1.0	2.4k	² ABLIKIM	23BP BES3	$e^+ e^-$ at 4.128–4.226 GeV
246.7±3.9±3.6	2.3k	³ ABLIKIM	23BX BES3	$e^+ e^-$ at 4.128–4.226 GeV
251.6±5.9±4.9	1.7k	⁴ ABLIKIM	21AF BES3	$e^+ e^-$ at 4.178, 4.226 GeV
244.4±2.3±2.9	4.9k	⁵ ABLIKIM	21AZ BES3	$e^+ e^-$ at 4.178, 4.226 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
243.0±5.8±4.0±1.0	950	^{6,7} ABLIKIM	21BE BES3	$e^+ e^-$ at 4.178, 4.226 GeV

¹ABLIKIM 24CG uses $\tau^+ \rightarrow (e^+ \nu_e, \mu^+ \nu_\mu, \pi^+, \rho^+) \bar{\nu}_\tau$ decays. The third uncertainty is due to the uncertainty of the D_s^+ lifetime.

- ² ABLIKIM 23BP uses $\tau^+ \rightarrow \pi^+ \bar{\nu}_\tau$ decays. The third uncertainty is due to the input parameters, mainly the D_s^+ lifetime.
- ³ ABLIKIM 23BX uses $\tau^+ \rightarrow \mu^+ \nu_\mu \bar{\nu}_\tau$ decays.
- ⁴ ABLIKIM 21AF uses $\tau^+ \rightarrow \pi^+ \pi^0 \bar{\nu}_\tau$ decays.
- ⁵ ABLIKIM 21AZ uses $\tau^+ \rightarrow e^+ \nu_e \bar{\nu}_\tau$ decays.
- ⁶ ABLIKIM 21BE uses $\tau^+ \rightarrow \pi^+ \bar{\nu}_\tau$ decays. When constrained by the Standard Model ratio of $\Gamma(D_s^+ \rightarrow \tau^+ \nu_\tau)/\Gamma(D_s^+ \rightarrow \mu^+ \nu_\mu) = 9.75$, the result is $243.2 \pm 2.3 \pm 3.3 \pm 1.0$.
- ⁷ The third uncertainty is dominated by the uncertainty of the D_s^+ lifetime. Superseded by ABLIKIM 23BP.

D_s^\pm REFERENCES

ABLIKIM	25A	PRL 134 011904	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	25BC	JHEP 2501 109	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	25BR	PRL 135 161902	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	25F	PRL 134 201902	M. Ablikim <i>et al.</i>	(BESIII Collab.)
AGGARWAL	25	JHEP 2504 036	L. Aggarwal <i>et al.</i>	(BELLE and BELLE II Collab.)
ABLIKIM	24AQ	PRL 132 091802	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	24AT	PRL 132 141901	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	24AV	PR D109 032011	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	24BD	JHEP 2405 335	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	24CF	PRL 133 121801	M. Ablikim	(BESIII Collab.)
ABLIKIM	24CG	PR D110 052002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	24CH	PR D110 052012	M. Ablikim	(BESIII Collab.)
ABLIKIM	24CI	PR D110 072017	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	24CT	JHEP 2411 119	M. Ablikim <i>et al.</i>	(BESIII Collab.)
AAIJ	23AN	JHEP 2307 204	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	23E	JHEP 2304 081	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	23L	JHEP 2307 067	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	23AL	PR D107 052010	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	23AV	PR D108 032001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	23BO	PR D108 092003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	23BP	PR D108 092014	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	23BR	PR D108 112001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	23BS	PR D108 112002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	23BX	JHEP 2309 124	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	23BZ	JHEP 2312 072	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ADACHI	23G	PRL 131 171803	I. Adachi <i>et al.</i>	(BELLE II Collab.)
LI	23G	PR D107 033003	L.K. Li <i>et al.</i>	(BELLE Collab.)
MOON	23	PR D108 L111102	H.K. Moon <i>et al.</i>	(BELLE Collab.)
ABLIKIM	22AA	JHEP 2204 058	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22AB	JHEP 2207 051	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22AC	JHEP 2208 196	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22AH	PRL 129 182001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22BH	PR D106 112004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22BI	PR D106 112006	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22BL	JHEP 2209 242	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22F	PR D105 L051103	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22J	PR D105 L031101	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22Z	JHEP 2201 052	M. Ablikim <i>et al.</i>	(BESIII Collab.)
AAIJ	21T	JHEP 2106 044	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	21U	JHEP 2106 019	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	21AB	JHEP 2106 181	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21AC	PR D104 012003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21AE	PR D104 012016	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21AF	PR D104 032001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21AR	PR D104 L071101	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21AZ	PRL 127 171801	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21BE	PR D104 052009	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21K	PR D103 092006	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21U	PR D104 032011	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21Y	PR D103 092004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
GUAN	21	PR D103 112005	Y. Guan <i>et al.</i>	(BELLE Collab.)
LEES	21A	PR D104 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)

ABLIKIM	20R	JHEP 2008 146	M. Ablikim <i>et al.</i>	(BESIII Collab.)
AAIJ	19G	JHEP 1903 176	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	19T	PRL 122 191803	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	19AD	PR D99 072002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19AH	PR D99 091101	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19AM	PR D99 112005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19BD	PR D100 112008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19BE	PRL 123 112001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19D	PRL 122 061801	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19E	PRL 122 071802	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19O	PR D99 031101	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19S	PRL 122 121801	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	18A	PR D97 012006	M. Ablikim <i>et al.</i>	(BESIII Collab.)
AAIJ	17AF	PL B771 21	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	17AN	PRL 119 101801	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	16O	PR D94 072004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	16T	PR D94 112003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	15Z	PL B750 466	M. Ablikim <i>et al.</i>	(BESIII Collab.)
HIETALA	15	PR D92 012009	J. Hietala <i>et al.</i>	(MINN, LUTH, OXF)
LEES	15D	PR D91 019901 (errat.)	J.P. Lees <i>et al.</i>	(BABAR Collab.)
AAIJ	14BD	JHEP 1410 025	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	14C	PL B728 585	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABAZOV	14B	PRL 112 111804	V.M. Abazov <i>et al.</i>	(D0 Collab.)
AAIJ	13AF	PL B724 203	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	13V	JHEP 1306 065	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	13W	JHEP 1306 112	R. Aaij <i>et al.</i>	(LHCb Collab.)
LEES	13E	PR D87 052012	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ONYISI	13	PR D88 032009	P.U.E. Onyisi <i>et al.</i>	(CLEO Collab.)
ZUPANC	13	JHEP 1309 139	A. Zupanc <i>et al.</i>	(BELLE Collab.)
DEL-AMO-SA...	11G	PR D83 052001	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
LEES	11E	PR D84 031103	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	11G	PR D84 072006	J.P. Lees <i>et al.</i>	(BABAR Collab.)
MARTIN	11	PR D84 012005	L. Martin <i>et al.</i>	(CLEO Collab.)
WILLIAMS	11	PR D84 054015	M. Williams	(LOIC)
ASNER	10	PR D81 052007	D.M. Asner <i>et al.</i>	(CLEO Collab.)
DEL-AMO-SA...	10J	PR D82 091103	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
Also		PR D91 019901 (errat.)	J.P. Lees <i>et al.</i>	(BABAR Collab.)
KO	10	PRL 104 181602	B.R. Ko <i>et al.</i>	(BELLE Collab.)
MENDEZ	10	PR D81 052013	H. Mendez <i>et al.</i>	(CLEO Collab.)
RUBIN	10	PR D82 092007	P. Rubin <i>et al.</i>	(CLEO Collab.)
ALEXANDER	09	PR D79 052001	J.P. Alexander <i>et al.</i>	(CLEO Collab.)
AUBERT	09O	PR D79 032003	B. Aubert <i>et al.</i>	(BABAR Collab.)
BEDIAGA	09	PR D80 096006	I. Bediaga <i>et al.</i>	(CBPF, NDAM)
DOBBS	09	PR D79 112008	S. Dobbs <i>et al.</i>	(CLEO Collab.)
ECKLUND	09	PR D80 052009	K.M. Ecklund <i>et al.</i>	(CLEO Collab.)
GE	09A	PR D80 051102	J.Y. Ge <i>et al.</i>	(CLEO Collab.)
KO	09	PRL 102 221802	B.R. Ko <i>et al.</i>	(BELLE Collab.)
MITCHELL	09A	PR D79 072008	R.E. Mitchell <i>et al.</i>	(CLEO Collab.)
NAIK	09A	PR D80 112004	P. Naik <i>et al.</i>	(CLEO Collab.)
ONYISI	09	PR D79 052002	P.U.E. Onyisi <i>et al.</i>	(CLEO Collab.)
WON	09	PR D80 111101	E. Won <i>et al.</i>	(BELLE Collab.)
YELTON	09	PR D80 052007	J. Yelton <i>et al.</i>	(CLEO Collab.)
ALEXANDER	08	PRL 100 161804	J.P. Alexander <i>et al.</i>	(CLEO Collab.)
ATHAR	08	PRL 100 181802	S.B. Athar <i>et al.</i>	(CLEO Collab.)
AUBERT	08AN	PR D78 051101	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08AO	PR D78 051102	B. Aubert <i>et al.</i>	(BABAR Collab.)
ECKLUND	08	PRL 100 161801	K.M. Ecklund <i>et al.</i>	(CLEO Collab.)
KLEMPPT	08	EPJ C55 39	E. Klempt, M. Matveev, A.V. Sarantsev	(BONN+)
LINK	08	PL B660 147	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
WIDHALM	08	PRL 100 241801	L. Widhalm <i>et al.</i>	(BELLE Collab.)
ADAMS	07A	PRL 99 191805	G.S. Adams <i>et al.</i>	(CLEO Collab.)
AUBERT	07V	PRL 98 141801	B. Aubert <i>et al.</i>	(BABAR Collab.)
PEDLAR	07A	PR D76 072002	T.K. Pedlar <i>et al.</i>	(CLEO Collab.)
Also		PRL 99 071802	M. Artuso <i>et al.</i>	(CLEO Collab.)
AUBERT	06N	PR D74 031103	B. Aubert <i>et al.</i>	(BABAR Collab.)
HUANG	06B	PR D74 112005	G.S. Huang <i>et al.</i>	(CLEO Collab.)
PDG	06	JP G33 1	W.-M. Yao <i>et al.</i>	(PDG Collab.)
AUBERT	05V	PR D71 091104	B. Aubert <i>et al.</i>	(BABAR Collab.)
LINK	05E	PL B622 239	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	05J	PRL 95 052003	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	05K	PL B624 166	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)

LINK	04	PL B585 200	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	04C	PL B586 183	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	04D	PL B586 191	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	04F	PL B601 10	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
ACOSTA	03D	PR D68 072004	D. Acosta <i>et al.</i>	(FNAL CDF-II Collab.)
ANISOVICH	03	EPJ A16 229	V.V. Anisovich <i>et al.</i>	
LINK	03D	PL B561 225	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	03F	PL B572 21	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
AUBERT	02G	PR D65 091104	B. Aubert <i>et al.</i>	(BABAR Collab.)
HEISTER	02I	PL B528 1	A. Heister <i>et al.</i>	(ALEPH Collab.)
LINK	02I	PL B541 227	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	02J	PL B541 243	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
ABBIENDI	01L	PL B516 236	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
AITALA	01A	PRL 86 765	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
IORI	01	PL B523 22	M. Iori <i>et al.</i>	(FNAL SELEX Collab.)
LINK	01C	PRL 87 162001	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
ALEXANDROV	00	PL B478 31	Y. Alexandrov <i>et al.</i>	(CERN BEATRICE Collab.)
AITALA	99	PL B445 449	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
AITALA	99D	PL B450 294	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
AITALA	99G	PL B462 401	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
BONVICINI	99	PRL 82 4586	G. Bonvicini <i>et al.</i>	(CLEO Collab.)
CHADHA	98	PR D58 032002	M. Chada <i>et al.</i>	(CLEO Collab.)
JESSOP	98	PR D58 052002	C.P. Jessop <i>et al.</i>	(CLEO Collab.)
ACCIARRI	97F	PL B396 327	M. Acciarri <i>et al.</i>	(L3 Collab.)
BALEST	97	PRL 79 1436	R. Balest <i>et al.</i>	(CLEO Collab.)
FRABETTI	97C	PL B401 131	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
FRABETTI	97D	PL B407 79	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
ARTUSO	96	PL B378 364	M. Artuso <i>et al.</i>	(CLEO Collab.)
BAI	95C	PR D52 3781	J.Z. Bai <i>et al.</i>	(BES Collab.)
BRANDENB...	95	PRL 75 3804	G.W. Brandenburg <i>et al.</i>	(CLEO Collab.)
FRABETTI	95B	PL B351 591	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
KODAMA	95	PL B345 85	K. Kodama <i>et al.</i>	(FNAL E653 Collab.)
ACOSTA	94	PR D49 5690	D. Acosta <i>et al.</i>	(CLEO Collab.)
VERY	94B	PL B337 405	P. Avery <i>et al.</i>	(CLEO Collab.)
BROWN	94	PR D50 1884	D. Brown <i>et al.</i>	(CLEO Collab.)
BUTLER	94	PL B324 255	F. Butler <i>et al.</i>	(CLEO Collab.)
FRABETTI	94F	PL B328 187	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
FRABETTI	93F	PRL 71 827	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
FRABETTI	93G	PL B313 253	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
KODAMA	93	PL B309 483	K. Kodama <i>et al.</i>	(FNAL E653 Collab.)
ALBRECHT	92B	ZPHY C53 361	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALEXANDER	92	PRL 68 1275	J. Alexander <i>et al.</i>	(CLEO Collab.)
VERY	92	PRL 68 1279	P. Avery <i>et al.</i>	(CLEO Collab.)
BARLAG	92C	ZPHY C55 383	S. Barlag <i>et al.</i>	(ACCMOR Collab.)
Also		ZPHY C48 29	S. Barlag <i>et al.</i>	(ACCMOR Collab.)
FRABETTI	92	PL B281 167	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
ALBRECHT	91	PL B255 634	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALVAREZ	91	PL B255 639	M.P. Alvarez <i>et al.</i>	(CERN NA14/2 Collab.)
ALBRECHT	90D	PL B245 315	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALEXANDER	90B	PRL 65 1531	J. Alexander <i>et al.</i>	(CLEO Collab.)
BARLAG	90C	ZPHY C46 563	S. Barlag <i>et al.</i>	(ACCMOR Collab.)
FRABETTI	90	PL B251 639	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
ANJOS	89E	PL B223 267	J.C. Anjos <i>et al.</i>	(FNAL E691 Collab.)
CHEN	89	PL B226 192	W.Y. Chen <i>et al.</i>	(CLEO Collab.)
ALBRECHT	88	PL B207 349	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ANJOS	88	PRL 60 897	J.C. Anjos <i>et al.</i>	(FNAL E691 Collab.)
RAAB	88	PR D37 2391	J.R. Raab <i>et al.</i>	(FNAL E691 Collab.)
BECKER	87B	PL B184 277	H. Becker <i>et al.</i>	(NA11 and NA32 Collabs.)
BLAYLOCK	87	PRL 58 2171	G.T. Blaylock <i>et al.</i>	(Mark III Collab.)
USHIDA	86	PRL 56 1767	N. Ushida <i>et al.</i>	(FNAL E531 Collab.)
ALBRECHT	85D	PL 153B 343	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
DERRICK	85B	PRL 54 2568	M. Derrick <i>et al.</i>	(HRS Collab.)
AIHARA	84D	PRL 53 2465	H. Aihara <i>et al.</i>	(TPC Collab.)
ALTHOFF	84	PL 136B 130	M. Althoff <i>et al.</i>	(TASSO Collab.)
BAILEY	84	PL 139B 320	R. Bailey <i>et al.</i>	(ACCMOR Collab.)
CHEN	83C	PRL 51 634	A. Chen <i>et al.</i>	(CLEO Collab.)
MOSTELLER	48	AST 2 No.5 30	R.A. Fisher, F. Mosteller	

————— **OTHER RELATED PAPERS** —————

RICHMAN 95 RMP 67 893 J.D. Richman, P.R. Burchat (UCSB, STAN)
