

$\phi(1020)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

 $\phi(1020)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1019.460 ±0.016	OUR AVERAGE			
1019.444 ±0.013 ±0.060		DRUZHININ 25	SND	$e^+e^- \rightarrow K_L^0 K_S^0$
1019.463 ±0.061	2.3M	¹ KOZYREV 18	CMD3	$e^+e^- \rightarrow K^+ K^-, K_S^0 K_L^0$
1019.462 ±0.042 ±0.056	28k	² LEES 14H	BABR	$e^+e^- \rightarrow K_S^0 K_L^0 \gamma$
1019.51 ±0.02 ±0.05		³ LEES 13Q	BABR	$e^+e^- \rightarrow K^+ K^- \gamma$
1019.30 ±0.02 ±0.10	105k	AKHMETSHIN 06	CMD2	$0.98-1.06 e^+e^- \rightarrow \pi^+ \pi^- \pi^0$
1019.52 ±0.05 ±0.05	17.4k	AKHMETSHIN 05	CMD2	$0.60-1.38 e^+e^- \rightarrow \eta \gamma$
1019.483 ±0.011 ±0.025	272k	⁴ AKHMETSHIN 04	CMD2	$e^+e^- \rightarrow K_L^0 K_S^0$
1019.42 ±0.05	1900k	⁵ ACHASOV 01E	SND	$e^+e^- \rightarrow K^+ K^-, K_S^0 K_L^0, \pi^+ \pi^- \pi^0$
1019.40 ±0.04 ±0.05	23k	AKHMETSHIN 01B	CMD2	$e^+e^- \rightarrow \eta \gamma$
1019.36 ±0.12		⁶ ACHASOV 00B	SND	$e^+e^- \rightarrow \eta \gamma$
1019.38 ±0.07 ±0.08	2200	⁷ AKHMETSHIN 99F	CMD2	$e^+e^- \rightarrow \pi^+ \pi^- \geq 2\gamma$
1019.51 ±0.07 ±0.10	11169	AKHMETSHIN 98	CMD2	$e^+e^- \rightarrow \pi^+ \pi^- \pi^0$
1019.5 ±0.4		BARBERIS 98	OMEG	$450 pp \rightarrow pp 2K^+ 2K^-$
1019.42 ±0.06	55600	AKHMETSHIN 95	CMD2	$e^+e^- \rightarrow \text{hadrons}$
1019.7 ±0.3	2012	DAVENPORT 86	MPSF	$400 pA \rightarrow 4K X$
1019.7 ±0.1 ±0.1	5079	ALBRECHT 85D	ARG	$10 e^+e^- \rightarrow K^+ K^- X$
1019.3 ±0.1	1500	ARENTOON 82	AEMS	$11.8 \text{ polar. } pp \rightarrow KK$
1019.67 ±0.17	25080	⁸ PELLINEN 82	RVUE	
1019.52 ±0.13	3681	BUKIN 78C	OLYA	$e^+e^- \rightarrow \text{hadrons}$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1019.443 ±0.010 ±0.060	1.28M	^{9,10} ACHASOV 24	SND	$e^+e^- \rightarrow K_L^0 K_S^0$
1019.0330 ±0.0084		¹¹ DUBNICKA 24	RVUE	$e^+e^- \rightarrow K^+ K^-, K_S^0 K_L^0$
1019.761 ±0.130	34M	¹² IGNATOV 24	CMD3	$e^+e^- \rightarrow \pi^+ \pi^-$
1018.4 ±0.5 ±0.1		¹³ ALBRECHT 20	CBAR	$0.9 \bar{p}p \rightarrow K^+ K^- \pi^0$
1019.21 ±0.04 ±0.03		¹⁴ HOID 20	RVUE	$e^+e^- \rightarrow \pi^0 \gamma$
1019.54 ±0.10 ±0.51		¹⁵ AAIJ 19H	LHCB	$pp \rightarrow D^\pm X$
1019.20 ±0.02 ±0.01		¹⁶ HOFERICHT... 19	RVUE	$e^+e^- \rightarrow \pi^+ \pi^- \pi^0$
1019.469 ±0.061	1.7M	KOZYREV 18	CMD3	$e^+e^- \rightarrow K^+ K^-$
1019.457 ±0.061	610k	KOZYREV 16	CMD3	$e^+e^- \rightarrow K_S^0 K_L^0$
1019.48 ±0.01		LEES 13F	BABR	$D^+ \rightarrow K^+ K^- \pi^+$

1019.441	± 0.008	± 0.080	542k	17	AKHMETSHIN 08	CMD2	1.02	$e^+e^- \rightarrow K^+K^-$
1019.63	± 0.07		12540	18	AUBERT,B 05J	BABR	$D^0 \rightarrow \bar{K}^0 K^+ K^-$	
1019.8	± 0.7				ARMSTRONG 86	OMEG	85 $\pi^+ / p p \rightarrow \pi^+ / p 4K p$	
1020.1	± 0.11		5526	18	ATKINSON 86	OMEG	20–70 γp	
1019.7	± 1.0				BEBEK 86	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$	
1019.411	± 0.008		642k	19	DIJKSTRA 86	SPEC	100–200 $\pi^\pm, \bar{p}, p, K^\pm$, on Be	
1020.9	± 0.2			18	FRAME 86	OMEG	13 $K^+ p \rightarrow \phi K^+ p$	
1021.0	± 0.2			18	ARMSTRONG 83B	OMEG	18.5 $K^- p \rightarrow K^- K^+ \Lambda$	
1020.0	± 0.5			18	ARMSTRONG 83B	OMEG	18.5 $K^- p \rightarrow K^- K^+ \Lambda$	
1019.7	± 0.3			18	BARATE 83	GOLI	190 $\pi^- \text{Be} \rightarrow 2\mu X$	
1019.8	± 0.2	± 0.5	766		IVANOV 81	OLYA	1–1.4 $e^+e^- \rightarrow K^+K^-$	
1019.4	± 0.5		337		COOPER 78B	HBC	0.7–0.8 $\bar{p} p \rightarrow K_S^0 K_L^0 \pi^+ \pi^-$	
1020	± 1		383	18	BALDI 77	CNTR	10 $\pi^- p \rightarrow \pi^- \phi p$	
1018.9	± 0.6		800		COHEN 77	ASPK	6 $\pi^\pm N \rightarrow K^+ K^- N$	
1019.7	± 0.5		454		KALBFLEISCH 76	HBC	2.18 $K^- p \rightarrow \Lambda K \bar{K}$	
1019.4	± 0.8		984		BESCH 74	CNTR	2 $\gamma p \rightarrow p K^+ K^-$	
1020.3	± 0.4		100		BALLAM 73	HBC	2.8–9.3 γp	
1019.4	± 0.7				BINNIE 73B	CNTR	$\pi^- p \rightarrow \phi n$	
1019.6	± 0.5		120	20	AGUILAR-...	HBC	3.9,4.6 $K^- p \rightarrow \Lambda K^+ K^-$	
1019.9	± 0.5		100	20	AGUILAR-...	HBC	3.9,4.6 $K^- p \rightarrow K^- p K^+ K^-$	
1020.4	± 0.5		131		COLLEY 72	HBC	10 $K^+ p \rightarrow K^+ p \phi$	
1019.9	± 0.3		410		STOTTLE... 71	HBC	2.9 $K^- p \rightarrow \Sigma / \Lambda K \bar{K}$	

¹ Average of KOZYREV 16 and KOZYREV 18 values taking into account the correlated uncertainties. Supersedes individual KOZYREV 16 and KOZYREV 18 results.

² Using a vector meson dominance model with contribution from $\phi(1020)$ and higher mass excitations of $\rho(770)$, $\omega(782)$, and $\phi(1020)$.

³ Using a phenomenological model based on KUHN 90 with a sum of Breit-Wigner resonances for $\rho(770)$, $\omega(782)$, $\phi(1020)$ and their higher mass excitations.

⁴ Update of AKHMETSHIN 99D

⁵ From the combined fit assuming that the total $\phi(1020)$ production cross section is saturated by those of K^+K^- , $K_S K_L$, $\pi^+ \pi^- \pi^0$, and $\eta\gamma$ decays modes and using ACHASOV 00B for the $\eta\gamma$ decay mode.

⁶ Using a total width of 4.43 ± 0.05 MeV. Systematic uncertainty included.

⁷ Using a total width of 4.43 ± 0.05 MeV.

⁸ PELLINEN 82 review includes AKERLOF 77, DAUM 81, BALDI 77, AYRES 74, DE-GROOT 74.

⁹ From a fit of the cross section in the energy region $1.000 < \sqrt{s} < 1.100$ GeV using a vector meson dominance model with contribution from $\phi(1020)$, $\rho(770)$, $\omega(782)$ and higher vector resonances.

¹⁰ Superseded by DRUZHININ 25.

¹¹ From the fit to data of CMD3, BABAR and BES3.

- ¹² From a fit of the pion form factor in the energy range $0.32 < \sqrt{s} < 1.2$ GeV using the GOUNARIS 68 parametrization with the complex phase of the $\rho - \omega$ interference leaving $\rho(1450)$, $\rho(1700)$ resonances as free parameters of the fit. Systematic errors not estimated.
- ¹³ Width fixed at 4.2 MeV.
- ¹⁴ The values were extracted from a dispersively improved Breit-Wigner parameterization and do not include vacuum polarization. Inclusion of vacuum polarization gives 1019.457 ± 0.020 MeV.
- ¹⁵ From the $D^\pm \rightarrow K^\pm K^+ K^-$ Dalitz plot fit with the Triple-M amplitude in the multi-meson model of AOUDE 18.
- ¹⁶ The values were extracted from a dispersively improved Breit-Wigner parameterization and do not include vacuum polarization.
- ¹⁷ Strongly correlated with AKHMETSHIN 04.
- ¹⁸ Systematic errors not evaluated.
- ¹⁹ Weighted and scaled average of 12 measurements of DIJKSTRA 86.
- ²⁰ Mass errors enlarged by us to Γ/\sqrt{N} ; see the note with the $K^*(892)$ mass.

$\phi(1020)$ WIDTH

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
4.249±0.013 OUR AVERAGE		Error includes scale factor of 1.1.		
4.234±0.22 ±0.12		DRUZHININ 25	SND	$e^+e^- \rightarrow K_L^0 K_S^0$
4.245±0.013	2.3M	¹ KOZYREV 18	CMD3	$e^+e^- \rightarrow K^+ K^-, K_S^0 K_L^0$
4.205±0.103±0.067	28k	² LEES 14H	BABR	$e^+e^- \rightarrow K_S^0 K_L^0 \gamma$
4.29 ±0.04 ±0.07		³ LEES 13Q	BABR	$e^+e^- \rightarrow K^+ K^- \gamma$
4.30 ±0.06 ±0.17	105k	AKHMETSHIN 06	CMD2	0.98–1.06 $e^+e^- \rightarrow \pi^+ \pi^- \pi^0$
4.280±0.033±0.025	272k	⁴ AKHMETSHIN 04	CMD2	$e^+e^- \rightarrow K_L^0 K_S^0$
4.21 ±0.04	1900k	⁵ ACHASOV 01E	SND	$e^+e^- \rightarrow K^+ K^-, K_S K_L, \pi^+ \pi^- \pi^0$
4.44 ±0.09	55600	AKHMETSHIN 95	CMD2	$e^+e^- \rightarrow$ hadrons
4.5 ±0.7	1500	ARENTON 82	AEMS	11.8 polar. $pp \rightarrow KK$
4.2 ±0.6	766	⁶ IVANOV 81	OLYA	1–1.4 $e^+e^- \rightarrow K^+ K^-$
4.3 ±0.6		⁶ CORDIER 80	DM1	$e^+e^- \rightarrow \pi^+ \pi^- \pi^0$
4.36 ±0.29	3681	⁶ BUKIN 78C	OLYA	$e^+e^- \rightarrow$ hadrons
4.4 ±0.6	984	⁶ BESCH 74	CNTR	$2 \gamma p \rightarrow p K^+ K^-$
4.67 ±0.72	681	⁶ BALAKIN 71	OSPK	$e^+e^- \rightarrow$ hadrons
4.09 ±0.29		BIZOT 70	OSPK	$e^+e^- \rightarrow$ hadrons
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
4.212±0.20 ±0.13	1.28M	^{7,8} ACHASOV 24	SND	$e^+e^- \rightarrow K_L^0 K_S^0$
4.151±0.018		⁹ DUBNICKA 24	RVUE	$e^+e^- \rightarrow K^+ K^-, K_S^0 K_L^0$
4.681±0.277	34M	¹⁰ IGNATOV 24	CMD3	$e^+e^- \rightarrow \pi^+ \pi^-$
4.07 ±0.13 ±0.01		¹¹ HOID 20	RVUE	$e^+e^- \rightarrow \pi^0 \gamma$
4.23 ±0.04 ±0.02		¹² HOFERICHT... 19	RVUE	$e^+e^- \rightarrow \pi^+ \pi^- \pi^0$
4.249±0.015	1.7M	KOZYREV 18	CMD3	$e^+e^- \rightarrow K^+ K^-$
4.240±0.017	610k	KOZYREV 16	CMD3	$e^+e^- \rightarrow K_S^0 K_L^0$
4.37 ±0.02		LEES 13F	BABR	$D^+ \rightarrow K^+ K^- \pi^+$

4.24 ±0.02 ±0.03	542k	¹³ AKHMETSHIN 08	CMD2	1.02 e ⁺ e ⁻ → K ⁺ K ⁻
4.28 ±0.13	12540	¹⁴ AUBERT,B 05J	BABR	D ⁰ → $\overline{K}^0 K^+ K^-$
4.45 ±0.06	271k	DIJKSTRA 86	SPEC	100 π ⁻ Be
3.6 ±0.8	337	⁶ COOPER 78B	HBC	0.7–0.8 $\overline{p} p \rightarrow K_S^0 K_L^0 \pi^+ \pi^-$
4.5 ±0.50	1300	^{6,14} AKERLOF 77	SPEC	400 ρA → K ⁺ K ⁻ X
4.5 ±0.8	500	^{6,14} AYRES 74	ASPK	3–6 π ⁻ p → K ⁺ K ⁻ n, K ⁻ p → K ⁺ K ⁻ Λ/Σ ⁰
3.81 ±0.37		COSME 74B	OSPK	e ⁺ e ⁻ → K _L ⁰ K _S ⁰
3.8 ±0.7	454	⁶ BORENSTEIN 72	HBC	2.18 K ⁻ p → K \overline{K} n

¹ Average of KOZYREV 16 and KOZYREV 18 values taking into account the correlated uncertainties. Supersedes individual KOZYREV 16 and KOZYREV 18 results.

² Using a vector meson dominance model with contribution from φ(1020) and higher mass excitations of ρ(770), ω(782), and φ(1020).

³ Using a phenomenological model based on KUHN 90 with a sum of Breit-Wigner resonances for ρ(770), ω(782), φ(1020) and their higher mass excitations.

⁴ Update of AKHMETSHIN 99D

⁵ From the combined fit assuming that the total φ(1020) production cross section is saturated by those of K⁺ K⁻, K_S K_L, π⁺ π⁻ π⁰, and ηγ decays modes and using ACHASOV 00B for the ηγ decay mode.

⁶ Width errors enlarged by us to 4Γ/√N; see the note with the K*(892) mass.

⁷ From a fit of the cross section in the energy region 1.000 < √s < 1.100 GeV using a vector meson dominance model with contribution from φ(1020), ρ(770), ω(782) and higher vector resonances.

⁸ Superseded by DRUZHININ 25.

⁹ From the fit to data of CMD3, BABAR and BES3.

¹⁰ From a fit of the pion form factor in the energy range 0.32 < √s < 1.2 GeV using the GOUNARIS 68 parametrization with the complex phase of the ρ – ω interference leaving ρ(1450), ρ(1700) resonances as free parameters of the fit. Systematic errors not estimated.

¹¹ The values were extracted from a dispersively improved Breit-Wigner parameterization.

¹² The values were extracted from a dispersively improved Breit-Wigner parameterization and do not include vacuum polarization.

¹³ Strongly correlated with AKHMETSHIN 04.

¹⁴ Systematic errors not evaluated.

φ(1020) DECAY MODES

Mode	Fraction (Γ _i /Γ)	Scale factor/ Confidence level
Γ ₁ K ⁺ K ⁻	(50.1 ±0.6) %	S=1.6
Γ ₂ K _L ⁰ K _S ⁰	(33.6 ±0.4) %	S=1.3
Γ ₃ ρπ + π ⁺ π ⁻ π ⁰	(14.7 ±0.4) %	S=1.5
Γ ₄ π ⁺ π ⁻ π ⁰		
Γ ₅ ηγ	(1.306±0.024) %	S=1.2
Γ ₆ π ⁰ γ	(1.33 ±0.05) × 10 ⁻³	
Γ ₇ ℓ ⁺ ℓ ⁻	—	
Γ ₈ e ⁺ e ⁻	(2.963±0.034) × 10 ⁻⁴	S=1.4
Γ ₉ μ ⁺ μ ⁻	(2.86 ±0.22) × 10 ⁻⁴	S=1.2

Γ_{10}	$\eta e^+ e^-$	$(1.08 \pm 0.04) \times 10^{-4}$	
Γ_{11}	$\pi^+ \pi^-$	$(9.5 \pm 1.9) \times 10^{-5}$	S=2.0
Γ_{12}	$\omega \pi^0$	$(4.7 \pm 0.5) \times 10^{-5}$	
Γ_{13}	$\omega \gamma$	< 5	% CL=84%
Γ_{14}	$\rho \gamma$	< 1.2	$\times 10^{-5}$ CL=90%
Γ_{15}	$\pi^+ \pi^- \gamma$	$(4.1 \pm 1.3) \times 10^{-5}$	
Γ_{16}	$f_0(980) \gamma$	$(3.22 \pm 0.19) \times 10^{-4}$	S=1.1
Γ_{17}	$\pi^0 \pi^0 \gamma$	$(1.13 \pm 0.06) \times 10^{-4}$	
Γ_{18}	$\pi^+ \pi^- \pi^+ \pi^-$	$(3.9 \begin{smallmatrix} +2.8 \\ -2.2 \end{smallmatrix}) \times 10^{-6}$	
Γ_{19}	$\pi^+ \pi^+ \pi^- \pi^- \pi^0$	< 4.6	$\times 10^{-6}$ CL=90%
Γ_{20}	$\pi^0 e^+ e^-$	$(1.33 \begin{smallmatrix} +0.07 \\ -0.10 \end{smallmatrix}) \times 10^{-5}$	
Γ_{21}	$\pi^0 \eta \gamma$	$(7.27 \pm 0.30) \times 10^{-5}$	S=1.5
Γ_{22}	$a_0(980) \gamma$	$(7.6 \pm 0.6) \times 10^{-5}$	
Γ_{23}	$K^0 \bar{K}^0 \gamma$	< 1.9	$\times 10^{-8}$ CL=90%
Γ_{24}	$\eta'(958) \gamma$	$(6.23 \pm 0.21) \times 10^{-5}$	
Γ_{25}	$\eta \pi^0 \pi^0 \gamma$	< 2	$\times 10^{-5}$ CL=90%
Γ_{26}	$\mu^+ \mu^- \gamma$	$(1.4 \pm 0.5) \times 10^{-5}$	
Γ_{27}	$\rho \gamma \gamma$	< 1.2	$\times 10^{-4}$ CL=90%
Γ_{28}	$\eta \pi^+ \pi^-$	< 1.8	$\times 10^{-5}$ CL=90%
Γ_{29}	$\eta \mu^+ \mu^-$	< 9.4	$\times 10^{-6}$ CL=90%
Γ_{30}	$\eta U \rightarrow \eta e^+ e^-$	< 1	$\times 10^{-6}$ CL=90%
Γ_{31}	invisible	< 1.7	$\times 10^{-4}$ CL=90%

Lepton Family number (*LF*) violating modes

Γ_{32}	$e^\pm \mu^\mp$	<i>LF</i>	< 2	$\times 10^{-6}$	CL=90%
Γ_{33}	$\pi^+ \pi^+ e^- e^-$	<i>L</i>	< 1.3	$\times 10^{-5}$	CL=90%

CONSTRAINED FIT INFORMATION

An overall fit to 30 branching ratios uses 84 measurements and one constraint to determine 14 parameters. The overall fit has a $\chi^2 = 98.5$ for 71 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x ₂	-70									
x ₃	-67	-6								
x ₅	-25	28	0							
x ₆	-14	17	1	8						
x ₈	60	-77	-3	-37	-23					
x ₉	-6	7	0	3	2	-9				
x ₁₁	-4	4	0	2	1	-6	1			
x ₁₂	-5	6	0	3	2	-8	1	0		
x ₁₆	0	0	0	0	0	0	0	0	0	
x ₁₇	-12	15	0	18	4	-19	2	1	2	0
x ₁₈	-1	1	0	1	0	-2	0	0	0	0
x ₂₂	0	0	0	0	0	0	0	0	0	0
x ₂₄	-8	9	0	32	3	-12	1	1	1	0
	x ₁	x ₂	x ₃	x ₅	x ₆	x ₈	x ₉	x ₁₁	x ₁₂	x ₁₆

x ₁₈	0		
x ₂₂	0	0	
x ₂₄	6	0	0
	x ₁₇	x ₁₈	x ₂₂

$\phi(1020)$ PARTIAL WIDTHS

$\Gamma(\eta\gamma)$ Γ_5

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
58.9±0.5±2.4	ACHASOV	00	SND $e^+e^- \rightarrow \eta\gamma$

$\Gamma(\pi^0\gamma)$ Γ_6

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
5.40±0.16 ^{+0.43} _{-0.40}	ACHASOV	00	SND $e^+e^- \rightarrow \pi^0\gamma$

$\Gamma(\ell^+\ell^-)$ Γ_7

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
1.320±0.017±0.015	¹ AMBROSINO	05	KLOE 1.02 $e^+e^- \rightarrow \mu^+\mu^-$

¹Weighted average of Γ_{ee} and $\sqrt{\Gamma_{ee}\Gamma_{\mu\mu}}$ from AMBROSINO 05 assuming lepton universality.

$\Gamma(e^+e^-)$ Γ_8

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
1.27 ±0.04 OUR EVALUATION			
1.251±0.021 OUR AVERAGE			Error includes scale factor of 1.1.
1.235±0.006±0.022	¹ AKHMETSHIN	11	CMD2 1.02 $e^+e^- \rightarrow \phi$

$1.32 \pm 0.05 \pm 0.03$	² AMBROSINO 05	KLOE	$1.02 e^+ e^- \rightarrow e^+ e^-$
1.28 ± 0.05	AKHMETSHIN 95	CMD2	$1.02 e^+ e^- \rightarrow \phi$
¹ Combined analysis of the CMD-2 data on $\phi \rightarrow K^+ K^-, K_S^0 K_L^0, \pi^+ \pi^- \pi^0, \eta \gamma$ assuming that the sum of their branching fractions is 0.99741 ± 0.00007 .			
² From forward-backward asymmetry and using $\Gamma_{\text{total}} = 4.26 \pm 0.05$ MeV from the 2004 edition of this Review.			

$(\Gamma(e^+ e^-) \times \Gamma(\mu^+ \mu^-))^{1/2}$	$(\Gamma_8 \Gamma_9)^{1/2}$		
VALUE (keV)	DOCUMENT ID	TECN	COMMENT
$1.320 \pm 0.018 \pm 0.017$	AMBROSINO 05	KLOE	$1.02 e^+ e^- \rightarrow \mu^+ \mu^-$

$\phi(1020) \Gamma(i) \Gamma(e^+ e^-) / \Gamma(\text{total})$

$\Gamma(K^+ K^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$	$\Gamma_1 \Gamma_8 / \Gamma$			
VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT

$0.6340 \pm 0.0070 \pm 0.0039$	¹ LEES	13Q	BABR	$e^+ e^- \rightarrow K^+ K^- \gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				

$0.669 \pm 0.001 \pm 0.023$	1.7M	KOZYREV	18	CMD3	$e^+ e^- \rightarrow K^+ K^-$
¹ Using a phenomenological model based on KUHN 90 with a sum of Breit-Wigner resonances for $\rho(770)$, $\omega(782)$, $\phi(1020)$ and their higher mass excitations. The first error combines statistical and systematic uncertainties. The second one is due to the parametrization of the charged kaon form factor and mass calibration.					

$\Gamma(K_L^0 K_S^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$	$\Gamma_2 \Gamma_8 / \Gamma$			
VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT

$0.4200 \pm 0.0033 \pm 0.0123$	28k	¹ LEES	14H	BABR	$e^+ e^- \rightarrow K_S^0 K_L^0 \gamma$
¹ Using a vector meson dominance model with contribution from $\phi(1020)$ and higher mass excitations of $\rho(770)$, $\omega(782)$, and $\phi(1020)$.					

$[\Gamma(\rho\pi) + \Gamma(\pi^+ \pi^- \pi^0)] \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$	$\Gamma_3 \Gamma_8 / \Gamma$		
VALUE (eV)	DOCUMENT ID	TECN	COMMENT

$184.1 \pm 2.1 \pm 8.0$	¹ LEES	21B	BABR	$10.5 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 \gamma$
¹ From the cross section for $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$ with contributions from $\rho(770)$, $\omega(782)$, $\phi(1020)$, $\omega(1420)$, and $\omega(1650)$.				


$\phi(1020) \Gamma(i) \Gamma(e^+ e^-) / \Gamma^2(\text{total})$

$\Gamma(K^+ K^-) / \Gamma_{\text{total}} \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$	$\Gamma_1 / \Gamma \times \Gamma_8 / \Gamma$			
VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT

14.85 ± 0.30 OUR FIT	Error includes scale factor of 1.6.				
14.6 ± 0.5 OUR AVERAGE	Error includes scale factor of 1.8. See the ideogram below.				
15.789 ± 0.541	1.7M	KOZYREV	18	CMD3	$e^+ e^- \rightarrow K^+ K^-$
$14.27 \pm 0.05 \pm 0.31$	542k	AKHMETSHIN 08	CMD2	$1.02 e^+ e^- \rightarrow K^+ K^-$	
$13.93 \pm 0.14 \pm 0.99$	1000k	¹ ACHASOV	01E	SND	$e^+ e^- \rightarrow K^+ K^-, K_S^0 K_L^0, \pi^+ \pi^- \pi^0$

¹ From the combined fit assuming that the total $\phi(1020)$ production cross section is saturated by those of $K^+ K^-$, $K_S^0 K_L^0$, $\pi^+ \pi^- \pi^0$, and $\eta \gamma$ decays modes and using ACHASOV 00B for the $\eta \gamma$ decay mode.

WEIGHTED AVERAGE
 14.6 ± 0.5 (Error scaled by 1.8)



$\Gamma(K_L^0 K_S^0)/\Gamma_{\text{total}} \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_2/\Gamma \times \Gamma_8/\Gamma$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
9.96 ± 0.08				OUR FIT
9.93 ± 0.08				OUR AVERAGE
9.83 ± 0.04 ± 0.10		DRUZHININ 25	SND	$e^+ e^- \rightarrow K_L^0 K_S^0$
10.078 ± 0.223	610k	¹ KOZYREV 16	CMD3	$e^+ e^- \rightarrow K_S^0 K_L^0$
10.01 ± 0.04 ± 0.17	272k	² AKHMETSHIN 04	CMD2	$e^+ e^- \rightarrow K_L^0 K_S^0$
10.27 ± 0.07 ± 0.34	500k	³ ACHASOV 01E	SND	$e^+ e^- \rightarrow K^+ K^-, K_S^0 K_L^0, \pi^+ \pi^- \pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

9.85 ± 0.03 ± 0.10 1.28M ^{4,5} ACHASOV 24 SND $e^+ e^- \rightarrow K_L^0 K_S^0$

¹ KOZYREV 16 also reports $\Gamma(e^+ e^-) B(\phi \rightarrow K_S^0 K_L^0) = (0.428 \pm 0.001 \pm 0.009)$ keV.

² Update of AKHMETSHIN 99D

³ From the combined fit assuming that the total $\phi(1020)$ production cross section is saturated by those of $K^+ K^-$, $K_S^0 K_L^0$, $\pi^+ \pi^- \pi^0$, and $\eta \gamma$ decays modes and using ACHASOV 00B for the $\eta \gamma$ decay mode.

⁴ From a fit of the cross section in the energy region $1.000 < \sqrt{s} < 1.100$ GeV using a vector meson dominance model with contribution from $\phi(1020)$, $\rho(770)$, $\omega(782)$ and higher vector resonances.

⁵ Superseded by DRUZHININ 25.

$[\Gamma(\rho\pi) + \Gamma(\pi^+ \pi^- \pi^0)]/\Gamma_{\text{total}} \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_3/\Gamma \times \Gamma_8/\Gamma$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
4.35 ± 0.13				OUR FIT Error includes scale factor of 1.4.
4.51 ± 0.14				OUR AVERAGE
4.51 ± 0.16 ± 0.11	105k	AKHMETSHIN 06	CMD2	0.98–1.06 $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$

4.665 ± 0.042 ± 0.261	400k	¹ ACHASOV	01E	SND	$e^+e^- \rightarrow K^+K^-,$ $K_S^0 K_L^0, \pi^+\pi^-\pi^0$
4.35 ± 0.27 ± 0.08	11169	² AKHMETSHIN 98	CMD2		$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
4.38 ± 0.12		BENAYOUN	10	RVUE	0.4–1.05 e^+e^-
4.30 ± 0.08 ± 0.21		³ AUBERT,B	04N	BABR	10.6 $e^+e^- \rightarrow$ $\pi^+\pi^-\pi^0\gamma$

¹ From the combined fit assuming that the total $\phi(1020)$ production cross section is saturated by those of K^+K^- , $K_S^0 K_L^0$, $\pi^+\pi^-\pi^0$, and $\eta\gamma$ decays modes and using ACHASOV 00B for the $\eta\gamma$ decay mode.


² Recalculated by us from the cross section in the peak.

³ Superseded by LEES 21B.

$\Gamma(\eta\gamma)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_5/\Gamma \times \Gamma_8/\Gamma$

<u>VALUE (units 10⁻⁶)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
3.87 ± 0.07 OUR FIT	Error includes scale factor of 1.2.			
3.93 ± 0.09 OUR AVERAGE	Error includes scale factor of 1.3. See the ideogram below.			
4.050 ± 0.067 ± 0.118	33k	¹ ACHASOV	07B	SND 0.6–1.38 $e^+e^- \rightarrow \eta\gamma$
4.093 ^{+0.040} _{-0.043} ± 0.247	17.4k	² AKHMETSHIN	05	CMD2 0.60-1.38 $e^+e^- \rightarrow \eta\gamma$
3.850 ± 0.041 ± 0.159	23k	^{3,4} AKHMETSHIN	01B	CMD2 $e^+e^- \rightarrow \eta\gamma$
4.00 ± 0.04 ± 0.11		⁵ ACHASOV	00	SND $e^+e^- \rightarrow \eta\gamma$
3.53 ± 0.08 ± 0.17	2200	^{6,7} AKHMETSHIN	99F	CMD2 $e^+e^- \rightarrow \eta\gamma$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
4.19 ± 0.06		⁸ BENAYOUN	10	RVUE 0.4–1.05 e^+e^-

WEIGHTED AVERAGE
3.93 ± 0.09 (Error scaled by 1.3)



- ¹ From a combined fit of $\sigma(e^+e^- \rightarrow \eta\gamma)$ with $\eta \rightarrow 3\pi^0$ and $\eta \rightarrow \pi^+\pi^-\pi^0$, and fixing $B(\eta \rightarrow 3\pi^0) / B(\eta \rightarrow \pi^+\pi^-\pi^0) = 1.44 \pm 0.04$. Recalculated by us from the cross section at the peak. Supersedes ACHASOV 00D and ACHASOV 06A.
- ² From the $\eta \rightarrow 2\gamma$ decay and using $B(\eta \rightarrow \gamma\gamma) = 39.43 \pm 0.26\%$.
- ³ From the $\eta \rightarrow 3\pi^0$ decay and using $B(\eta \rightarrow 3\pi^0) = (32.24 \pm 0.29) \times 10^{-2}$.
- ⁴ The combined fit from 600 to 1380 MeV taking into account $\rho(770)$, $\omega(782)$, $\phi(1020)$, and $\rho(1450)$ (mass and width fixed at 1450 MeV and 310 MeV respectively).
- ⁵ From the $\eta \rightarrow 2\gamma$ decay and using $B(\eta \rightarrow 2\gamma) = (39.21 \pm 0.34) \times 10^{-2}$.
- ⁶ Recalculated by the authors from the cross section in the peak.
- ⁷ From the $\eta \rightarrow \pi^+\pi^-\pi^0$ decay and using $B(\eta \rightarrow \pi^+\pi^-\pi^0) = (23.1 \pm 0.5) \times 10^{-2}$.
- ⁸ A simultaneous fit of $e^+e^- \rightarrow \pi^+\pi^-$, $\pi^+\pi^-\pi^0$, $\pi^0\gamma$, $\eta\gamma$ data.

$\Gamma(\pi^0\gamma)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_6/\Gamma \times \Gamma_8/\Gamma$

VALUE (units 10^{-7})	EVTS	DOCUMENT ID	TECN	COMMENT
3.94 ± 0.16 OUR FIT				
3.95 ± 0.17 OUR AVERAGE				
4.04 ± 0.09 ± 0.19		¹ ACHASOV 16A	SND	0.60–1.38 $e^+e^- \rightarrow \pi^0\gamma$
3.75 ± 0.11 ± 0.29	18k	AKHMETSHIN 05	CMD2	0.60–1.38 $e^+e^- \rightarrow \pi^0\gamma$
• • •				We do not use the following data for averages, fits, limits, etc. • • •
4.29 ± 0.11		² BENAYOUN 10	RVUE	0.4–1.05 e^+e^-
3.67 ± 0.10 ^{+0.27} _{-0.25}		³ ACHASOV 00	SND	$e^+e^- \rightarrow \pi^0\gamma$


- ¹ From the VMD model with the interfering $\rho(770)$, $\omega(782)$, $\phi(1020)$ resonances, and an additional resonance describing the total contribution of the $\rho(1450)$ and $\omega(1420)$ states. Supersedes ACHASOV 00.
- ² A simultaneous fit of $e^+e^- \rightarrow \pi^+\pi^-$, $\pi^+\pi^-\pi^0$, $\pi^0\gamma$, $\eta\gamma$ data.
- ³ From the $\pi^0 \rightarrow 2\gamma$ decay and using $B(\pi^0 \rightarrow 2\gamma) = (98.798 \pm 0.032) \times 10^{-2}$.

$\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_9/\Gamma \times \Gamma_8/\Gamma$

VALUE (units 10^{-8})	DOCUMENT ID	TECN	COMMENT
8.5 ± 0.6 OUR FIT			Error includes scale factor of 1.2.
8.8 ± 0.9 OUR AVERAGE			Error includes scale factor of 1.5. See the ideogram below.
8.36 ± 0.59 ± 0.37	ACHASOV 01G	SND	$e^+e^- \rightarrow \mu^+\mu^-$
9.9 ± 1.4 ± 0.9	¹ ACHASOV 99C	SND	$e^+e^- \rightarrow \mu^+\mu^-$
14.4 ± 3.0	² VASSERMAN 81	OLYA	$e^+e^- \rightarrow \mu^+\mu^-$
8.6 ± 5.9	² AUGUSTIN 73	OSPK	$e^+e^- \rightarrow \mu^+\mu^-$

- ¹ Recalculated by the authors from the cross section in the peak.
- ² Recalculated by us from the cross section in the peak.

WEIGHTED AVERAGE
 8.8 ± 0.9 (Error scaled by 1.5)



Values above of weighted average, error, and scale factor are based upon the data in this ideogram only. They are not necessarily the same as our 'best' values, obtained from a least-squares constrained fit utilizing measurements of other (related) quantities as additional information.

χ^2

+	ACHASOV	01G	SND	0.5
+	ACHASOV	99C	SND	0.4
+	VASSERMAN	81	OLYA	3.4
+	AUGUSTIN	73	OSPK	4.3

(Confidence Level = 0.116)

$$\Gamma(\mu^+ \mu^-) / \Gamma_{\text{total}} \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$$

$$\Gamma_9 / \Gamma \times \Gamma_8 / \Gamma$$

$$\Gamma(\pi^+ \pi^-) / \Gamma_{\text{total}} \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$$

$$\Gamma_{11} / \Gamma \times \Gamma_8 / \Gamma$$

VALUE (units 10^{-8})	EVTS	DOCUMENT ID	TECN	COMMENT
2.8 ± 0.6	OUR FIT	Error includes scale factor of 2.0.		
2.8 ± 0.5	OUR AVERAGE	Error includes scale factor of 1.8. See the ideogram below.		
$3.51 \pm 0.33 \pm 0.24$	34M	¹ IGNATOV 24	CMD3	$e^+ e^- \rightarrow \pi^+ \pi^-$
$2.1 \pm 0.3 \pm 0.3$		² ACHASOV 00c	SND	$e^+ e^- \rightarrow \pi^+ \pi^-$
$1.95^{+1.15}_{-0.87}$		³ GOLUBEV 86	ND	$e^+ e^- \rightarrow \pi^+ \pi^-$
$6.01^{+3.19}_{-2.51}$		³ VASSERMAN 81	OLYA	$e^+ e^- \rightarrow \pi^+ \pi^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
3.31 ± 0.99		⁴ BENAYOUN 13	RVUE	$0.4-1.05 e^+ e^-$


¹ From a fit of the pion form factor in the energy range $0.32 < \sqrt{s} < 1.2$ GeV using the GOUNARIS 68 parametrization with the complex phase of the $\rho - \omega$ interference with ω and ϕ masses and widths constrained by the values and their errors from PDG 22, and leaving $\rho(1450)$, $\rho(1700)$ resonances as free parameters of the fit. A fit without the constraints from PDG 22 gives a value $\Gamma(e^+ e^-) \times \Gamma(\pi^+ \pi^-) / \Gamma_{\text{total}}^2 = (3.65 \pm 0.24) \times 10^{-8}$.

² Recalculated by the authors from the cross section in the peak.

³ Recalculated by us from the cross section in the peak.

⁴ A simultaneous fit to $e^+ e^- \rightarrow \pi^+ \pi^-, \pi^+ \pi^- \pi^0, \pi^0 \gamma, \eta \gamma, K \bar{K},$ and $\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$ data.

WEIGHTED AVERAGE
 2.8 ± 0.5 (Error scaled by 1.8)



$\Gamma(\omega\pi^0)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{12}/\Gamma \times \Gamma_8/\Gamma$

VALUE (units 10^{-8})	DOCUMENT ID	TECN	COMMENT
1.40 ± 0.15 OUR FIT			
1.37 ± 0.17 ± 0.01	^{1,2} AMBROSINO 08G	KLOE	$e^+e^- \rightarrow \pi^+\pi^-2\pi^0, 2\pi^0\gamma$

¹ Recalculated by the authors from the cross section at the peak.
² AMBROSINO 08G reports $[\Gamma(\phi(1020) \rightarrow \omega\pi^0)/\Gamma_{\text{total}} \times \Gamma(\phi(1020) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+\pi^-\pi^0)] = (1.22 \pm 0.13 \pm 0.08) \times 10^{-8}$ which we divide by our best (shown rounded) value $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best (shown rounded) value.

$\Gamma(\pi^0\pi^0\gamma)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{17}/\Gamma \times \Gamma_8/\Gamma$

VALUE (units 10^{-8})	DOCUMENT ID	TECN	COMMENT
3.34 ± 0.17 OUR FIT			
3.33^{+0.04+0.19}_{-0.09-0.20}	¹ AMBROSINO 07	KLOE	$e^+e^- \rightarrow \pi^0\pi^0\gamma$

¹ Calculated by the authors from the cross section at the peak.

$\Gamma(\pi^+\pi^-\pi^+\pi^-)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{18}/\Gamma \times \Gamma_8/\Gamma$

VALUE (units 10^{-9})	EVTS	DOCUMENT ID	TECN	COMMENT
1.2^{+0.8}_{-0.7} OUR FIT				
1.17 ± 0.52 ± 0.64	3285	¹ AKHMETSHIN 00E	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-$

¹ Recalculated by the authors from the cross section in the peak.

$\phi(1020)$ BRANCHING RATIOS $\Gamma(K^+ K^-)/\Gamma_{\text{total}}$ Γ_1/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.501±0.006 OUR FIT Error includes scale factor of 1.6.**0.493±0.010 OUR AVERAGE**

0.492±0.012	2913	AKHMETSHIN 95	CMD2	$e^+ e^- \rightarrow K^+ K^-$
0.44 ±0.05	321	KALBFLEISCH 76	HBC	2.18 $K^- p \rightarrow \Lambda K^+ K^-$
0.49 ±0.06	270	DEGROOT 74	HBC	4.2 $K^- p \rightarrow \Lambda \phi$
0.540±0.034	565	BALAKIN 71	OSPK	$e^+ e^- \rightarrow K^+ K^-$
0.48 ±0.04	252	LINDSEY 66	HBC	2.1–2.7 $K^- p \rightarrow \Lambda K^+ K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.493±0.003±0.007		¹ AKHMETSHIN 11	CMD2	1.02 $e^+ e^- \rightarrow K^+ K^-$
0.476±0.017	1000k	² ACHASOV 01E	SND	$e^+ e^- \rightarrow K^+ K^-, K_S K_L, \pi^+ \pi^- \pi^0$

¹ Combined analysis of the CMD-2 data on $\phi \rightarrow K^+ K^-, K_S^0 K_L^0, \pi^+ \pi^- \pi^0, \eta \gamma$ assuming that the sum of their branching fractions is 0.99741 ± 0.00007 .² Using $B(\phi \rightarrow e^+ e^-) = (2.93 \pm 0.14) \times 10^{-4}$. $\Gamma(K_L^0 K_S^0)/\Gamma_{\text{total}}$ Γ_2/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.336±0.004 OUR FIT Error includes scale factor of 1.3.**0.331±0.009 OUR AVERAGE**

0.335±0.010	40644	AKHMETSHIN 95	CMD2	$e^+ e^- \rightarrow K_L^0 K_S^0$
0.326±0.035		DOLINSKY 91	ND	$e^+ e^- \rightarrow K_L^0 K_S^0$
0.310±0.024		DRUZHININ 84	ND	$e^+ e^- \rightarrow K_L^0 K_S^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.336±0.002±0.006		¹ AKHMETSHIN 11	CMD2	1.02 $e^+ e^- \rightarrow K_S^0 K_L^0$
0.351±0.013	500k	² ACHASOV 01E	SND	$e^+ e^- \rightarrow K^+ K^-, K_S K_L, \pi^+ \pi^- \pi^0$
0.27 ±0.03	133	KALBFLEISCH 76	HBC	2.18 $K^- p \rightarrow \Lambda K_L^0 K_S^0$
0.257±0.030	95	³ BALAKIN 71	OSPK	$e^+ e^- \rightarrow K_L^0 K_S^0$
0.40 ±0.04	167	LINDSEY 66	HBC	2.1–2.7 $K^- p \rightarrow \Lambda K_L^0 K_S^0$

¹ Combined analysis of the CMD-2 data on $\phi \rightarrow K^+ K^-, K_S^0 K_L^0, \pi^+ \pi^- \pi^0, \eta \gamma$ assuming that the sum of their branching fractions is 0.99741 ± 0.00007 .² Using $B(\phi \rightarrow e^+ e^-) = (2.93 \pm 0.14) \times 10^{-4}$.³ Balakin error increased by Paul. $\Gamma(K_L^0 K_S^0)/\Gamma(K^+ K^-)$ Γ_2/Γ_1

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

67.1± 1.4 OUR FIT Error includes scale factor of 1.5.**74.0± 3.1 OUR AVERAGE**

70 ± 6	2732	BUKIN 78C	OLYA	$e^+ e^- \rightarrow K_L^0 K_S^0$
82 ± 8		LOSTY 78	HBC	4.2 $K^- p \rightarrow \phi$ hyperon
71 ± 5		LAVEN 77	HBC	10 $K^- p \rightarrow K^+ K^- \Lambda$
71 ± 8		LYONS 77	HBC	3–4 $K^- p \rightarrow \Lambda \phi$
89 ± 10	144	AGUILAR-...	72B	3.9,4.6 $K^- p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

59.3 ± 2.3 ± 2.1	665k	¹ ABLIKIM	25BR	BES3	$D_S^+ \rightarrow \phi(\rightarrow K_S^0 K_L^0)\pi^+$
64.4 ± 1.7		² DUBNICKA	24	RVUE	$e^+e^- \rightarrow K^+K^-, K_S^0 K_L^0$
63.8 ± 2.2	2.3M	³ KOZYREV	18	CMD3	$e^+e^- \rightarrow K_L^0 K_S^0, K^+K^-$
68 ± 3		⁴ AKHMETSHIN	95	CMD2	$e^+e^- \rightarrow K_L^0 K_S^0, K^+K^-$

¹ From a measurement of $D_S^+ \rightarrow \phi(\rightarrow K_S^0 K_L^0)\pi^+$ using the PDG 24 value for $B(D_S^+ \rightarrow \phi(\rightarrow K^+K^-)\pi^+)$.

² From a fit to CMD3, BABAR and BES3 data.

³ The prediction taking into account phase-space difference, radiative corrections, isospin breaking, and the Sommerfeld-Gamow-Sakharov factor gives 0.630.

⁴ Theoretical analysis of BRAMON 00 taking into account phase-space difference, electromagnetic radiative corrections, as well as isospin breaking, predicts 0.62. FLOREZ-BAEZ 08 predicts 0.63 considering also structure-dependent radiative corrections. FIS-CHBACH 02 calculates additional corrections caused by the close threshold and predicts 0.68. See also BENAYOUN 01 and DUBYNSKIY 07. BENAYOUN 12 obtains 0.71 ± 0.01 in the HLS model.

$\Gamma(K_L^0 K_S^0)/\Gamma(K\bar{K})$ $\Gamma_2/(\Gamma_1+\Gamma_2)$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.402 ± 0.005 OUR FIT	Error includes scale factor of 1.5.			
0.45 ± 0.04 OUR AVERAGE				
0.44 ± 0.07		¹ LONDON	66	HBC 2.24 $K^-p \rightarrow \Lambda K\bar{K}$
0.48 ± 0.07	52	BADIER	65B	HBC 3 K^-p
0.40 ± 0.10	34	SCHLEIN	63	HBC 1.95 $K^-p \rightarrow \Lambda K\bar{K}$

¹ This is probably not affected by their controversial background subtraction; the value is from their numbers of $K_1 K_2$ vs $K^+ K^-$ events.

$[\Gamma(\rho\pi) + \Gamma(\pi^+\pi^-\pi^0)]/\Gamma_{\text{total}}$ Γ_3/Γ

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.147 ± 0.004 OUR FIT	Error includes scale factor of 1.5.			
0.151 ± 0.009 OUR AVERAGE	Error includes scale factor of 1.7.			
0.161 ± 0.008	11761	AKHMETSHIN 95	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
0.143 ± 0.007		DOLINSKY 91	ND	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.155 ± 0.002 ± 0.005		¹ AKHMETSHIN 11	CMD2	1.02 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
0.159 ± 0.008	400k	² ACHASOV	01E	SND $e^+e^- \rightarrow K^+K^-, K_S^0 K_L^0, \pi^+\pi^-\pi^0$
0.145 ± 0.009 ± 0.003	11169	³ AKHMETSHIN 98	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
0.139 ± 0.007		⁴ PARROUR	76B	OSPK e^+e^-

¹ Combined analysis of the CMD-2 data on $\phi \rightarrow K^+K^-, K_S^0 K_L^0, \pi^+\pi^-\pi^0, \eta\gamma$ assuming that the sum of their branching fractions is 0.99741 ± 0.00007 .

² Using $B(\phi \rightarrow e^+e^-) = (2.93 \pm 0.14) \times 10^{-4}$.

³ Using $B(\phi \rightarrow e^+e^-) = (2.99 \pm 0.08) \times 10^{-4}$.

⁴ Using $\Gamma(\phi) = 4.1$ MeV. If interference between the $\rho\pi$ and 3π modes is neglected, the fraction of the $\rho\pi$ is more than 80% at the 90% confidence level.

$$[\Gamma(\rho\pi) + \Gamma(\pi^+\pi^-\pi^0)]/\Gamma(K^+K^-) \quad \Gamma_3/\Gamma_1$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.293±0.011 OUR FIT Error includes scale factor of 1.6.

0.229±0.014 OUR AVERAGE

0.230±0.014±0.010		¹ ABLIKIM	25A	BES3	$D_S^+ \rightarrow \phi\pi^+$
0.222±0.019±0.016		² ABLIKIM	25F	BES3	$D_S^+ \rightarrow \phi\rho^+$
0.28 ±0.09	34	AGUILAR-...	72B	HBC	3.9,4.6 K^-p

¹ Using ABLIKIM 21AE.

² Using ABLIKIM 21U.

$$[\Gamma(\rho\pi) + \Gamma(\pi^+\pi^-\pi^0)]/\Gamma(K\bar{K}) \quad \Gamma_3/(\Gamma_1+\Gamma_2)$$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.175±0.006 OUR FIT Error includes scale factor of 1.5.

0.24 ±0.04 OUR AVERAGE

0.237±0.039	CERRADA	77B	HBC	4.2 $K^-p \rightarrow \Lambda 3\pi$
0.30 ±0.15	LONDON	66	HBC	2.24 $K^-p \rightarrow \Lambda\pi^+\pi^-\pi^0$

$$[\Gamma(\rho\pi) + \Gamma(\pi^+\pi^-\pi^0)]/\Gamma(K_L^0 K_S^0) \quad \Gamma_3/\Gamma_2$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.437±0.013 OUR FIT Error includes scale factor of 1.4.

0.51 ±0.05 OUR AVERAGE

0.56 ±0.07	3681	BUKIN	78c	OLYA	$e^+e^- \rightarrow K_L^0 K_S^0, \pi^+\pi^-\pi^0$
0.47 ±0.06	516	COSME	74	OSPK	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$

$$\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}} \quad \Gamma_4/\Gamma$$

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
-------	-----	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

≈ 0.0087	1.98M	^{1,2} ALOISIO	03	KLOE	1.02 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
<0.0006	90	³ ACHASOV	02	SND	1.02 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
<0.23	90	³ CORDIER	80	DM1	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
<0.20	90	³ PARROUR	76B	OSPK	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$

¹ From a fit without limitations on charged and neutral ρ masses and widths.

² Adding the direct and $\omega\pi$ contributions and considering the interference between the $\rho\pi$ and $\pi^+\pi^-\pi^0$.

³ Neglecting the interference between the $\rho\pi$ and $\pi^+\pi^-\pi^0$.

$$\Gamma(\eta\gamma)/\Gamma_{\text{total}} \quad \Gamma_5/\Gamma$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

1.306±0.024 OUR FIT Error includes scale factor of 1.2.

1.26 ±0.04 OUR AVERAGE

1.246±0.025±0.057	10k	¹ ACHASOV	98F	SND	$e^+e^- \rightarrow 7\gamma$
1.18 ±0.11	279	² AKHMETSHIN	95	CMD2	$e^+e^- \rightarrow \pi^+\pi^-3\gamma$
1.30 ±0.06		³ DRUZHININ	84	ND	$e^+e^- \rightarrow 3\gamma$
1.4 ±0.2		⁴ DRUZHININ	84	ND	$e^+e^- \rightarrow 6\gamma$
0.88 ±0.20	290	KURDADZE	83c	OLYA	$e^+e^- \rightarrow 3\gamma$
1.35 ±0.29		ANDREWS	77	CNTR	6.7–10 γCu
1.5 ±0.4	54	³ COSME	76	OSPK	e^+e^-

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.38 ± 0.02 ± 0.02		⁵ AKHMETSHIN 11	CMD2	1.02 e ⁺ e ⁻ → ηγ
1.37 ± 0.05 ± 0.02	33k	⁶ ACHASOV 07B	SND	0.6–1.38 e ⁺ e ⁻ → ηγ
1.373 ± 0.014 ± 0.085	17.4k	^{7,8} AKHMETSHIN 05	CMD2	0.60–1.38 e ⁺ e ⁻ → ηγ
1.287 ± 0.013 ± 0.063		^{9,10} AKHMETSHIN 01B	CMD2	e ⁺ e ⁻ → ηγ
1.338 ± 0.012 ± 0.052		¹¹ ACHASOV 00	SND	e ⁺ e ⁻ → ηγ
1.18 ± 0.03 ± 0.06	2200	¹² AKHMETSHIN 99F	CMD2	e ⁺ e ⁻ → ηγ
1.21 ± 0.07		¹³ BENAYOUN 96	RVUE	0.54–1.04 e ⁺ e ⁻ → ηγ

¹ Using $B(\phi \rightarrow e^+e^-) = (2.99 \pm 0.08) \times 10^{-4}$ and $B(\eta \rightarrow 3\pi^0) = (32.2 \pm 0.4) \times 10^{-2}$.

² From $\pi^+\pi^-\pi^0$ decay mode of η .

³ From 2γ decay mode of η .

⁴ From $3\pi^0$ decay mode of η .

⁵ Combined analysis of the CMD-2 data on $\phi \rightarrow K^+K^-, K_S^0 K_L^0, \pi^+\pi^-\pi^0, \eta\gamma$ assuming that the sum of their branching fractions is 0.99741 ± 0.00007 .

⁶ ACHASOV 07B reports $[\Gamma(\phi(1020) \rightarrow \eta\gamma)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow e^+e^-)] = (4.050 \pm 0.067 \pm 0.118) \times 10^{-6}$ which we divide by our best (shown rounded) value $B(\phi(1020) \rightarrow e^+e^-) = (2.963 \pm 0.034) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best (shown rounded) value. Supersedes ACHASOV 00D and ACHASOV 06A.

⁷ Using $B(\phi \rightarrow e^+e^-) = (2.98 \pm 0.04) \times 10^{-4}$ and $B(\eta \rightarrow \gamma\gamma) = 39.43 \pm 0.26\%$.

⁸ Not independent of the corresponding $\Gamma(e^+e^-) \times \Gamma(\eta\gamma)/\Gamma_{\text{total}}^2$.

⁹ Using $B(\phi \rightarrow e^+e^-) = (2.99 \pm 0.08) \times 10^{-4}$ and $B(\eta \rightarrow 3\pi^0) = (32.24 \pm 0.29) \times 10^{-2}$.

¹⁰ The combined fit from 600 to 1380 MeV taking into account $\rho(770), \omega(782), \phi(1020)$, and $\rho(1450)$ (mass and width fixed at 1450 MeV and 310 MeV respectively).

¹¹ From the $\eta \rightarrow 2\gamma$ decay and using $B(\phi \rightarrow e^+e^-) = (2.99 \pm 0.08) \times 10^{-4}$.

¹² From $\pi^+\pi^-\pi^0$ decay mode of η and using $B(\phi \rightarrow e^+e^-) = (2.99 \pm 0.08) \times 10^{-4}$.

¹³ Reanalysis of DRUZHININ 84, DOLINSKY 89, and DOLINSKY 91 taking into account a triangle anomaly contribution.

$\Gamma(\pi^0\gamma)/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE (units 10^{-3}) EVTS DOCUMENT ID TECN COMMENT

1.33 ± 0.05 OUR FIT

1.31 ± 0.13 OUR AVERAGE

1.30 ± 0.13		DRUZHININ 84	ND	e ⁺ e ⁻ → 3γ
1.4 ± 0.5	32	COSME 76	OSPK	e ⁺ e ⁻

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.367 ± 0.072		¹ ACHASOV 16A	SND	0.60–1.38 e ⁺ e ⁻ → π ⁰ γ
1.258 ± 0.037 ± 0.077	18k	^{2,3} AKHMETSHIN 05	CMD2	0.60–1.38 e ⁺ e ⁻ → π ⁰ γ
1.226 ± 0.036 ^{+0.096} _{-0.089}		⁴ ACHASOV 00	SND	e ⁺ e ⁻ → π ⁰ γ
1.26 ± 0.17		⁵ BENAYOUN 96	RVUE	0.54–1.04 e ⁺ e ⁻ → π ⁰ γ

¹ Using $B(\phi \rightarrow e^+e^-)$ from PDG 15. Supersedes ACHASOV 00.

² Using $B(\phi \rightarrow e^+e^-) = (2.98 \pm 0.04) \times 10^{-4}$.

³ Not independent of the corresponding $\Gamma(e^+e^-) \times \Gamma(\pi^0\gamma)/\Gamma_{\text{total}}^2$.

⁴ From the $\pi^0 \rightarrow 2\gamma$ decay and using $B(\phi \rightarrow e^+e^-) = (2.99 \pm 0.08) \times 10^{-4}$.

⁵ Reanalysis of DRUZHININ 84, DOLINSKY 89, and DOLINSKY 91 taking into account a triangle anomaly contribution.

$\Gamma(\eta\gamma)/\Gamma(\pi^0\gamma)$ Γ_5/Γ_6

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
$10.9 \pm 0.3^{+0.7}_{-0.8}$	ACHASOV	00	SND $e^+e^- \rightarrow \eta\gamma, \pi^0\gamma$

$\Gamma(e^+e^-)/\Gamma_{\text{total}}$ Γ_8/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.963 ± 0.034 OUR FIT Error includes scale factor of 1.4.				
2.98 ± 0.07 OUR AVERAGE Error includes scale factor of 1.1.				
2.93 ± 0.14	1900k	¹ ACHASOV	01E	SND $e^+e^- \rightarrow K^+K^-, K_S K_L, \pi^+\pi^-\pi^0$
2.88 ± 0.09	55600	AKHMETSHIN	95	CMD2 $e^+e^- \rightarrow$ hadrons
3.00 ± 0.21	3681	BUKIN	78C	OLYA $e^+e^- \rightarrow$ hadrons
3.10 ± 0.14		² PARROUR	76	OSPK e^+e^-
3.3 ± 0.3		COSME	74	OSPK $e^+e^- \rightarrow$ hadrons
2.81 ± 0.25	681	BALAKIN	71	OSPK $e^+e^- \rightarrow$ hadrons
3.50 ± 0.27		CHATELUS	71	OSPK e^+e^-

¹From the combined fit assuming that the total $\phi(1020)$ production cross section is saturated by those of K^+K^- , $K_S K_L$, $\pi^+\pi^-\pi^0$, and $\eta\gamma$ decays modes and using ACHASOV 00B for the $\eta\gamma$ decay mode.

²Using total width 4.2 MeV. They detect 3π mode and observe significant interference with ω tail. This is accounted for in the result quoted above.

$\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$ Γ_9/Γ

<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.86 ± 0.22 OUR FIT Error includes scale factor of 1.2.			
2.96 ± 0.16 OUR AVERAGE Error includes scale factor of 1.1.			
2.69 ± 0.46	¹ HAYES	71	CNTR $8.3, 9.8 \gamma C \rightarrow \mu^+\mu^- X$
2.17 ± 0.60	¹ EARLES	70	CNTR $6.0 \gamma C \rightarrow \mu^+\mu^- X$
● ● ● We use the following data for averages but not for fits. ● ● ●			
$3.045 \pm 0.049 \pm 0.148$	² AAIJ	24B	LHCB $D^+, D_S^+ \rightarrow \pi^+\phi$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
$2.87 \pm 0.20 \pm 0.14$	³ ACHASOV	01G	SND $e^+e^- \rightarrow \mu^+\mu^-$
$3.30 \pm 0.45 \pm 0.32$	⁴ ACHASOV	99C	SND $e^+e^- \rightarrow \mu^+\mu^-$
4.83 ± 1.02	⁵ VASSERMAN	81	OLYA $e^+e^- \rightarrow \mu^+\mu^-$
2.87 ± 1.98	⁵ AUGUSTIN	73	OSPK $e^+e^- \rightarrow \mu^+\mu^-$

¹Neglecting interference between resonance and continuum.

²Value not independent from the measurement of $B(\phi \rightarrow \mu^+\mu^-)/B(\phi \rightarrow e^+e^-)$.

³Using $B(\phi \rightarrow e^+e^-) = (2.91 \pm 0.07) \times 10^{-4}$.

⁴Using $B(\phi \rightarrow e^+e^-) = (2.99 \pm 0.08) \times 10^{-4}$.

⁵Recalculated by us using $B(\phi \rightarrow e^+e^-) = (2.99 \pm 0.08) \times 10^{-4}$.

$\Gamma(\mu^+\mu^-)/\Gamma(e^+e^-)$ Γ_9/Γ_8

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.022 ± 0.012 ± 0.048				
	24k	AAIJ	24B	LHCB $D^+, D_S^+ \rightarrow \pi^+\phi$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$1.026 \pm 0.020 \pm 0.056$	7.5k	AAIJ	24B	LHCB $D^+ \rightarrow \pi^+\phi$
$1.017 \pm 0.013 \pm 0.051$	16.7k	AAIJ	24B	LHCB $D_S^+ \rightarrow \pi^+\phi$

$\Gamma(\eta e^+ e^-)/\Gamma_{\text{total}}$ Γ_{10}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.08 ± 0.04 OUR AVERAGE				
1.075 ± 0.007 ± 0.038	30k	¹ BABUSCI 15	KLOE	1.02 $e^+ e^- \rightarrow \eta e^+ e^-$
1.19 ± 0.19 ± 0.12	213	² ACHASOV 01B	SND	$e^+ e^- \rightarrow \eta e^+ e^-$
1.14 ± 0.10 ± 0.06	355	³ AKHMETSHIN 01	CMD2	$e^+ e^- \rightarrow \eta e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.13 ± 0.14 ± 0.07	183	⁴ AKHMETSHIN 01	CMD2	$e^+ e^- \rightarrow \eta e^+ e^-$
1.21 ± 0.14 ± 0.09	130	⁵ AKHMETSHIN 01	CMD2	$e^+ e^- \rightarrow \eta e^+ e^-$
1.04 ± 0.20 ± 0.08	42	⁶ AKHMETSHIN 01	CMD2	$e^+ e^- \rightarrow \eta e^+ e^-$
1.3 $\begin{smallmatrix} +0.8 \\ -0.6 \end{smallmatrix}$	7	GOLUBEV 85	ND	$e^+ e^- \rightarrow \eta e^+ e^-$

¹ Using $B(\eta \rightarrow 3\pi^0) = (32.57 \pm 0.23)\%$ from PDG 12.

² Using $B(\eta \rightarrow \gamma\gamma) = (39.25 \pm 0.32)\%$, $B(\phi \rightarrow \eta\gamma) = (1.26 \pm 0.06)\%$, and $B(\phi \rightarrow e^+ e^-) = (3.00 \pm 0.06) \times 10^{-4}$.

³ The average of the branching ratios separately obtained from the $\eta \rightarrow \gamma\gamma$, $3\pi^0$, $\pi^+ \pi^- \pi^0$ decays.

⁴ From $\eta \rightarrow \gamma\gamma$ decays and using $B(\eta \rightarrow \gamma\gamma) = (39.33 \pm 0.25) \times 10^{-2}$, $B(\eta \rightarrow \pi^+ \pi^- \gamma) = (4.75 \pm 11) \times 10^{-2}$, and $B(\phi \rightarrow \eta\gamma) = (1.297 \pm 0.033) \times 10^{-2}$.

⁵ From $\eta \rightarrow 3\pi^0$ decays and using $B(\pi^0 \rightarrow \gamma\gamma) = (98.798 \pm 0.033) \times 10^{-2}$, $B(\eta \rightarrow 3\pi^0) = (32.24 \pm 0.29) \times 10^{-2}$, $B(\eta \rightarrow \pi^+ \pi^- \gamma) = (4.75 \pm 0.11) \times 10^{-2}$, and $B(\phi \rightarrow \eta\gamma) = (1.297 \pm 0.033) \times 10^{-2}$.

⁶ From $\eta \rightarrow \pi^+ \pi^- \pi^0$ decays and using $B(\pi^0 \rightarrow \gamma\gamma) = (98.798 \pm 0.033) \times 10^{-2}$, $B(\pi^0 \rightarrow e^+ e^- \gamma) = (1.198 \pm 0.032) \times 10^{-2}$, $B(\eta \rightarrow \pi^+ \pi^- \pi^0) = (23.0 \pm 0.4) \times 10^{-2}$, $B(\phi \rightarrow \pi^+ \pi^- \pi^0) = (15.5 \pm 0.6) \times 10^{-2}$, and $B(\phi \rightarrow \eta\gamma) = (1.297 \pm 0.033) \times 10^{-2}$.

 $\Gamma(\pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.71 ± 0.11 ± 0.09		¹ ACHASOV 00C	SND	$e^+ e^- \rightarrow \pi^+ \pi^-$
0.65 $\begin{smallmatrix} +0.38 \\ -0.29 \end{smallmatrix}$		¹ GOLUBEV 86	ND	$e^+ e^- \rightarrow \pi^+ \pi^-$
2.01 $\begin{smallmatrix} +1.07 \\ -0.84 \end{smallmatrix}$		¹ VASSERMAN 81	OLYA	$e^+ e^- \rightarrow \pi^+ \pi^-$
<6.6	95	BUKIN 78B	OLYA	$e^+ e^- \rightarrow \pi^+ \pi^-$
<2.7	95	ALVENSLEB... 72	CNTR	6.7 $\gamma C \rightarrow C \pi^+ \pi^-$

¹ Using $B(\phi \rightarrow e^+ e^-) = (2.99 \pm 0.08) \times 10^{-4}$.

 $\Gamma(\omega \pi^0)/\Gamma_{\text{total}}$ Γ_{12}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
4.7 ± 0.5 OUR FIT			
5.2 $\begin{smallmatrix} +1.3 \\ -1.1 \end{smallmatrix}$	^{1,2} AULCHENKO 00A	SND	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 \pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
4.4 ± 0.6	³ AMBROSINO 08G	KLOE	$e^+ e^- \rightarrow \pi^+ \pi^- 2\pi^0, 2\pi^0 \gamma$
~ 5.4	⁴ ACHASOV 00E	SND	$e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$
5.5 $\begin{smallmatrix} +1.6 \\ -1.4 \end{smallmatrix} \pm 0.3$	^{2,5} AULCHENKO 00A	SND	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 \pi^0$
4.8 $\begin{smallmatrix} +1.9 \\ -1.7 \end{smallmatrix} \pm 0.8$	⁴ ACHASOV 99	SND	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 \pi^0$

¹ Using the 1996 and 1998 data.

² $(2.3 \pm 0.3)\%$ correction for other decay modes of the $\omega(782)$ applied.

³ Not independent of the corresponding $\Gamma(\omega\pi^0) \times \Gamma(e^+e^-) / \Gamma^2(\text{total})$.

⁴ Using the 1996 data.

⁵ Using the 1998 data.

$\Gamma(\omega\gamma)/\Gamma_{\text{total}}$						Γ_{13}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<0.05	84	LINDSEY	66	HBC	2.1–2.7 $K^- p \rightarrow \Lambda\pi^+\pi^-$ neutrals	

$\Gamma(\rho\gamma)/\Gamma_{\text{total}}$						Γ_{14}/Γ
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT		
< 0.12	90	¹ AKHMETSHIN 99B	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\gamma$		
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
< 7	90	AKHMETSHIN 97C	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\gamma$		
<200	84	LINDSEY	66	HBC	2.1–2.7 $K^- p \rightarrow \Lambda\pi^+\pi^-$ neutrals	

¹ Supersedes AKHMETSHIN 97C.

$\Gamma(\pi^+\pi^-\gamma)/\Gamma_{\text{total}}$						Γ_{15}/Γ
VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	
0.41±0.12±0.04		30175	¹ AKHMETSHIN 99B	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\gamma$	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
< 0.3	90		² AKHMETSHIN 97C	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\gamma$	
<600	90		KALBFLEISCH 75	HBC	2.18 $K^- p \rightarrow \Lambda\pi^+\pi^-\gamma$	
< 70	90		COSME	74	OSPK $e^+e^- \rightarrow \pi^+\pi^-\gamma$	
<400	90		LINDSEY	65	HBC 2.1–2.7 $K^- p \rightarrow \Lambda\pi^+\pi^-$ neutrals	

¹ For $E_\gamma > 20$ MeV and assuming that $B(\phi(1020) \rightarrow f_0(980)\gamma)$ is negligible. Supersedes AKHMETSHIN 97C.

² For $E_\gamma > 20$ MeV and assuming that $B(\phi(1020) \rightarrow f_0(980)\gamma)$ is negligible.

$\Gamma(f_0(980)\gamma)/\Gamma_{\text{total}}$						Γ_{16}/Γ
VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	
3.22±0.19 OUR FIT	Error includes scale factor of 1.1.					
3.21±0.19 OUR AVERAGE						
$3.21^{+0.03}_{-0.09} \pm 0.18$			¹ AMBROSINO 07	KLOE	$e^+e^- \rightarrow \pi^0\pi^0\gamma$	
$2.90 \pm 0.21 \pm 1.54$			² AKHMETSHIN 99C	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\gamma, \pi^0\pi^0\gamma$	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
4.47 ± 0.21	2438		³ ALOISIO 02D	KLOE	$e^+e^- \rightarrow \pi^0\pi^0\gamma$	
$3.5 \pm 0.3 \begin{smallmatrix} +1.3 \\ -0.5 \end{smallmatrix}$	419		^{4,5} ACHASOV 00H	SND	$e^+e^- \rightarrow \pi^0\pi^0\gamma$	
$1.93 \pm 0.46 \pm 0.50$	27188		⁶ AKHMETSHIN 99B	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\gamma$	
$3.05 \pm 0.25 \pm 0.72$	268		⁷ AKHMETSHIN 99C	CMD2	$e^+e^- \rightarrow \pi^0\pi^0\gamma$	
1.5 ± 0.5	268		⁸ AKHMETSHIN 99C	CMD2	$e^+e^- \rightarrow \pi^0\pi^0\gamma$	

$3.42 \pm 0.30 \pm 0.36$	164	⁴ ACHASOV	98I	SND	$e^+e^- \rightarrow 5\gamma$
< 1	90	⁹ AKHMETSHIN	97C	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\gamma$
< 7	90	¹⁰ AKHMETSHIN	97C	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\gamma$
< 20	90	DRUZHININ	87	ND	$e^+e^- \rightarrow \pi^0\pi^0\gamma$

¹ Obtained by the authors taking into account the $\pi^+\pi^-$ decay mode. Includes a component due to $\pi\pi$ production via the $f_0(500)$ meson. Supersedes ALOISIO 02D.

² From the combined fit of the photon spectra in the reactions $e^+e^- \rightarrow \pi^+\pi^-\gamma$, $\pi^0\pi^0\gamma$.

³ From the negative interference with the $f_0(500)$ meson of AITALA 01B using the ACHASOV 89 parameterization for the $f_0(980)$, a Breit-Wigner for the $f_0(500)$, and ACHASOV 01F for the $\rho\pi$ contribution. Superseded by AMBROSINO 07.

⁴ Assuming that the $\pi^0\pi^0\gamma$ final state is completely determined by the $f_0\gamma$ mechanism, neglecting the decay $B(\phi \rightarrow K\bar{K}\gamma)$ and using $B(f_0 \rightarrow \pi^+\pi^-) = 2B(f_0 \rightarrow \pi^0\pi^0)$.

⁵ Using the value $B(\phi \rightarrow \eta\gamma) = (1.338 \pm 0.053) \times 10^{-2}$.

⁶ For $E_\gamma > 20$ MeV. Supersedes AKHMETSHIN 97C.

⁷ Neglecting other intermediate mechanisms ($\rho\pi$, $\sigma\gamma$).

⁸ A narrow pole fit taking into account $f_0(980)$ and $f_0(1200)$ intermediate mechanisms.

⁹ For destructive interference with the Bremsstrahlung process

¹⁰ For constructive interference with the Bremsstrahlung process

$\Gamma(f_0(980)\gamma)/\Gamma(\eta\gamma)$ Γ_{16}/Γ_5

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

2.47 ± 0.15 OUR FIT Error includes scale factor of 1.1.

2.6 ± 0.2	$^{+0.8}_{-0.3}$	419	¹ ACHASOV	00H	SND	$e^+e^- \rightarrow \pi^0\pi^0\gamma$
---------------	------------------	-----	----------------------	-----	-----	---------------------------------------

¹ Assuming that the $\pi^0\pi^0\gamma$ final state is completely determined by the $f_0\gamma$ mechanism, neglecting the decay $B(\phi \rightarrow K\bar{K}\gamma)$ and using $B(f_0 \rightarrow \pi^+\pi^-) = 2B(f_0 \rightarrow \pi^0\pi^0)$.

$\Gamma(\pi^0\pi^0\gamma)/\Gamma_{total}$ Γ_{17}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	-------------	--------------------	-------------	----------------

1.07 ± 0.06 OUR AVERAGE

1.07	$^{+0.01}_{-0.03}$	$^{+0.06}_{-0.06}$		¹ AMBROSINO	07	KLOE	$e^+e^- \rightarrow \pi^0\pi^0\gamma$
------	--------------------	--------------------	--	------------------------	----	------	---------------------------------------

1.08	± 0.17	± 0.09	268	AKHMETSHIN	99C	CMD2	$e^+e^- \rightarrow \pi^0\pi^0\gamma$
------	------------	------------	-----	------------	-----	------	---------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.09	± 0.03	± 0.05	2438	ALOISIO	02D	KLOE	$e^+e^- \rightarrow \pi^0\pi^0\gamma$
------	------------	------------	------	---------	-----	------	---------------------------------------

1.158	± 0.093	± 0.052	419	^{2,3} ACHASOV	00H	SND	$e^+e^- \rightarrow \pi^0\pi^0\gamma$
-------	-------------	-------------	-----	------------------------	-----	-----	---------------------------------------

< 10			90	DRUZHININ	87	ND	$e^+e^- \rightarrow 5\gamma$
--------	--	--	----	-----------	----	----	------------------------------

¹ Supersedes ALOISIO 02D.

² Using the value $B(\phi \rightarrow \eta\gamma) = (1.338 \pm 0.053) \times 10^{-2}$.

³ Supersedes ACHASOV 98I. Excluding $\omega\pi^0$.

$\Gamma(\pi^0\pi^0\gamma)/\Gamma(\eta\gamma)$ Γ_{17}/Γ_5

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

0.86 ± 0.04 OUR FIT

0.865 ± 0.070 ± 0.017	419	¹ ACHASOV	00H	SND	$e^+e^- \rightarrow \pi^0\pi^0\gamma$
------------------------------	-----	----------------------	-----	-----	---------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.90	± 0.08	± 0.07	164	ACHASOV	98I	SND	$e^+e^- \rightarrow 5\gamma$
------	------------	------------	-----	---------	-----	-----	------------------------------

¹ Supersedes ACHASOV 98I. Excluding $\omega\pi^0$.

$\Gamma(\pi^+\pi^-\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{18}/Γ

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
6.5 ± 2.7 ± 1.6		6.8k	¹ AKHMETSHIN 17	CMD3	$e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
3.93 ± 1.74 ± 2.14		3.3k	AKHMETSHIN 00E	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-$
< 870	90		CORDIER 79	WIRE	$e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-$

¹ Using the cross section at the ϕ meson peak $\sigma(\phi) = 4172 \pm 42$ nb, the nonresonant cross section $\sigma(0) = 1.263 \pm 0.027$ nb and $\text{Re}(Z) = 0.146 \pm 0.030$, $\text{Im}(Z) = -0.002 \pm 0.024$ for the complex amplitude of the $\phi \rightarrow \pi^+\pi^-\pi^+\pi^-$ transition.

$\Gamma(\pi^+\pi^+\pi^-\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{19}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 4.6	90	AKHMETSHIN 00E	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 150	95	BARKOV 88	CMD	$e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-\pi^0$

$\Gamma(\pi^0 e^+ e^-)/\Gamma_{\text{total}}$ Γ_{20}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
1.33^{+0.07}_{-0.10} OUR AVERAGE					

1.35 ± 0.05^{+0.05}_{-0.10} 9.5k ¹ ANASTASI 16B KLOE $e^+e^- \rightarrow \pi^0 e^+ e^-$

1.01 ± 0.28 ± 0.29 52 ² ACHASOV 02D SND $e^+e^- \rightarrow \pi^0 e^+ e^-$

1.22 ± 0.34 ± 0.21 46 ³ AKHMETSHIN 01C CMD2 $e^+e^- \rightarrow \pi^0 e^+ e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 12 90 DOLINSKY 88 ND $e^+e^- \rightarrow \pi^0 e^+ e^-$

¹ Using $B(\pi^0 \rightarrow \gamma\gamma)$ from the 2014 Edition of this Review (PDG 14).

² Using various branching ratios from the 2000 Edition of this Review (PDG 00).

³ Using $B(\pi^0 \rightarrow \gamma\gamma) = 0.98798 \pm 0.00032$, $B(\phi \rightarrow \eta\gamma) = (1.297 \pm 0.033) \times 10^{-2}$, and $B(\eta \rightarrow \pi^+\pi^-\gamma) = (4.75 \pm 0.11) \times 10^{-2}$.

$\Gamma(\pi^0 \eta \gamma)/\Gamma_{\text{total}}$ Γ_{21}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
7.27 ± 0.30 OUR AVERAGE					Error includes scale factor of 1.5. See the ideogram below.

7.06 ± 0.22 16.9k ¹ AMBROSINO 09F KLOE $1.02 e^+e^- \rightarrow \eta\pi^0\gamma$

8.51 ± 0.51 ± 0.57 607 ² ALOISIO 02C KLOE $e^+e^- \rightarrow \eta\pi^0\gamma$

7.96 ± 0.60 ± 0.40 197 ³ ALOISIO 02C KLOE $e^+e^- \rightarrow \eta\pi^0\gamma$

8.8 ± 1.4 ± 0.9 36 ⁴ ACHASOV 00F SND $e^+e^- \rightarrow \eta\pi^0\gamma$

9.0 ± 2.4 ± 1.0 80 AKHMETSHIN 99C CMD2 $e^+e^- \rightarrow \eta\pi^0\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

7.01 ± 0.10 ± 0.20 13.3k ^{2,5} AMBROSINO 09F KLOE $1.02 e^+e^- \rightarrow \eta\pi^0\gamma$

7.12 ± 0.13 ± 0.22 3.6k ^{3,6} AMBROSINO 09F KLOE $1.02 e^+e^- \rightarrow \eta\pi^0\gamma$

8.3 ± 2.3 ± 1.2 20 ACHASOV 98B SND $e^+e^- \rightarrow 5\gamma$

< 250 90 DOLINSKY 91 ND $e^+e^- \rightarrow \pi^0\eta\gamma$

¹ Combined results of $\eta \rightarrow \gamma\gamma$ and $\eta \rightarrow \pi^+\pi^-\pi^0$ decay modes measurements.


² From the decay mode $\eta \rightarrow \gamma\gamma$.

³ From the decay mode $\eta \rightarrow \pi^+\pi^-\pi^0$.

⁴ Supersedes ACHASOV 98B.

⁵ Using $B(\phi \rightarrow \eta\gamma) = (1.304 \pm 0.025)\%$, $B(\eta \rightarrow 3\pi^0) = (32.56 \pm 0.23)\%$, and $B(\eta \rightarrow \gamma\gamma) = (39.31 \pm 0.20)\%$.

⁶ Using $B(\phi \rightarrow \eta\gamma) = (1.304 \pm 0.025)\%$, $B(\eta \rightarrow 3\pi^0) = (32.56 \pm 0.23)\%$, and $B(\eta \rightarrow \pi^+\pi^-\pi^0) = (22.73 \pm 0.28)\%$.



$\Gamma(a_0(980)\gamma)/\Gamma_{\text{total}}$

Γ_{22}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
7.6±0.6 OUR FIT					
7.6±0.6 OUR AVERAGE					
7.4±0.7			1 ALOISIO	02C	KLOE $e^+e^- \rightarrow \eta\pi^0\gamma$
8.8±1.7		36	2 ACHASOV	00F	SND $e^+e^- \rightarrow \eta\pi^0\gamma$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
11 ±2			3 GOKALP	02	RVUE $e^+e^- \rightarrow \eta\pi^0\gamma$
<500	90		DOLINSKY	91	ND $e^+e^- \rightarrow \pi^0\eta\gamma$

¹ Using $M_{a_0(980)}=984.8$ MeV and assuming $a_0(980)\gamma$ dominance.

² Assuming $a_0(980)\gamma$ dominance in the $\eta\pi^0\gamma$ final state.

³ Using data of ACHASOV 00F.

$\Gamma(f_0(980)\gamma)/\Gamma(a_0(980)\gamma)$

Γ_{16}/Γ_{22}

VALUE	DOCUMENT ID	TECN	COMMENT
6.1±0.6	1 ALOISIO	02C	KLOE $e^+e^- \rightarrow \eta\pi^0\gamma$

¹ Using results of ALOISIO 02D and assuming that $f_0(980)$ decays into $\pi\pi$ only and $a_0(980)$ into $\eta\pi$ only.

$\Gamma(K^0 \bar{K}^0 \gamma) / \Gamma_{\text{total}}$			Γ_{23} / Γ		
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$<1.9 \times 10^{-8}$	90	AMBROSINO 09C	KLOE	$e^+ e^- \rightarrow K_S^0 K_S^0 \gamma$	

$\Gamma(\eta'(958)\gamma) / \Gamma_{\text{total}}$			Γ_{24} / Γ		
<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>

6.23 ± 0.21 OUR FIT
6.24 ± 0.30 OUR AVERAGE

6.23 ± 0.27 ± 0.12	3407	¹ AMBROSINO 07A	KLOE	1.02 $e^+ e^- \rightarrow \pi^+ \pi^- 7\gamma$	
6.7 $^{+2.8}_{-2.4} \pm 0.8$	12	² AULCHENKO 03B	SND	$e^+ e^- \rightarrow \eta' \gamma$	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
6.7 $^{+5.0}_{-4.2} \pm 1.5$	7	AULCHENKO 03B	SND	$e^+ e^- \rightarrow 7\gamma$	
6.10 ± 0.61 ± 0.43	120	³ ALOISIO 02E	KLOE	1.02 $e^+ e^- \rightarrow \pi^+ \pi^- 3\gamma$	
8.2 $^{+2.1}_{-1.9} \pm 1.1$	21	⁴ AKHMETSHIN 00B	CMD2	$e^+ e^- \rightarrow \pi^+ \pi^- 3\gamma$	
4.9 $^{+2.2}_{-1.8} \pm 0.6$	9	⁵ AKHMETSHIN 00F	CMD2	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^+ \pi^- \geq 2\gamma$	
6.4 ± 1.6	30	⁶ AKHMETSHIN 00F	CMD2	$e^+ e^- \rightarrow \eta'(958)\gamma$	
6.7 $^{+3.4}_{-2.9} \pm 1.0$	5	⁷ AULCHENKO 99	SND	$e^+ e^- \rightarrow \pi^+ \pi^- 3\gamma$	
<11	90	AULCHENKO 98	SND	$e^+ e^- \rightarrow 7\gamma$	
12 $^{+7}_{-5} \pm 2$	6	⁴ AKHMETSHIN 97B	CMD2	$e^+ e^- \rightarrow \pi^+ \pi^- 3\gamma$	
<41	90	DRUZHININ 87	ND	$e^+ e^- \rightarrow \gamma \eta \pi^+ \pi^-$	

¹ AMBROSINO 07A reports $[\Gamma(\phi(1020) \rightarrow \eta'(958)\gamma) / \Gamma_{\text{total}}] / [B(\phi(1020) \rightarrow \eta\gamma)] = (4.77 \pm 0.09 \pm 0.19) \times 10^{-3}$ which we multiply by our best (shown rounded) value $B(\phi(1020) \rightarrow \eta\gamma) = (1.306 \pm 0.024) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best (shown rounded) value.

² Averaging AULCHENKO 03B with AULCHENKO 99.

³ Using $B(\phi \rightarrow \eta\gamma) = (1.297 \pm 0.033)\%$.

⁴ Using the value $B(\phi \rightarrow \eta\gamma) = (1.26 \pm 0.06) \times 10^{-2}$.

⁵ Using $B(\phi \rightarrow K_L^0 K_S^0) = (33.8 \pm 0.6)\%$.

⁶ Averaging AKHMETSHIN 00B with AKHMETSHIN 00F.

⁷ Using the value $B(\eta' \rightarrow \eta \pi^+ \pi^-) = (43.7 \pm 1.5) \times 10^{-2}$ and $B(\eta \rightarrow \gamma\gamma) = (39.25 \pm 0.31) \times 10^{-2}$.

$\Gamma(\eta'(958)\gamma) / \Gamma(K_L^0 K_S^0)$			Γ_{24} / Γ_2		
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	

1.85 ± 0.06 OUR FIT

1.46 $^{+0.64}_{-0.54} \pm 0.18$	9	¹ AKHMETSHIN 00F	CMD2	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^+ \pi^- \geq 2\gamma$	
----------------------------------	---	-----------------------------	------	--	--

¹ Using various branching ratios of K_S^0 , K_L^0 , η , η' from the 2000 edition (The European Physical Journal **C15** 1 (2000)) of this Review.

$\Gamma(\eta'(958)\gamma)/\Gamma(\eta\gamma)$ Γ_{24}/Γ_5

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

4.77±0.15 OUR FIT**4.78±0.20 OUR AVERAGE**4.77±0.09±0.19 3407 AMBROSINO 07A KLOE 1.02 $e^+e^- \rightarrow \pi^+\pi^-7\gamma$ 4.70±0.47±0.31 120 ¹ALOISIO 02E KLOE 1.02 $e^+e^- \rightarrow \pi^+\pi^-3\gamma$ 6.5 $\begin{smallmatrix} +1.7 \\ -1.5 \end{smallmatrix}$ ±0.8 21 AKHMETSHIN 00B CMD2 $e^+e^- \rightarrow \pi^+\pi^-3\gamma$

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

9.5 $\begin{smallmatrix} +5.2 \\ -4.0 \end{smallmatrix}$ ±1.4 6 ²AKHMETSHIN 97B CMD2 $e^+e^- \rightarrow \pi^+\pi^-3\gamma$ ¹From the decay mode $\eta' \rightarrow \eta\pi^+\pi^-$, $\eta \rightarrow \gamma\gamma$.²Superseded by AKHMETSHIN 00B. $\Gamma(\eta\pi^0\pi^0\gamma)/\Gamma_{\text{total}}$ Γ_{25}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<2 90 AULCHENKO 98 SND $e^+e^- \rightarrow 7\gamma$ $\Gamma(\mu^+\mu^-\gamma)/\Gamma_{\text{total}}$ Γ_{26}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

1.43±0.45±0.14 27188 ¹AKHMETSHIN 99B CMD2 $e^+e^- \rightarrow \mu^+\mu^-\gamma$

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

2.3 ±1.0 824 ± 33 ²AKHMETSHIN 97C CMD2 $e^+e^- \rightarrow \mu^+\mu^-\gamma$ ¹For $E_\gamma > 20$ MeV. Supersedes AKHMETSHIN 97C.²For $E_\gamma > 20$ MeV. $\Gamma(\rho\gamma\gamma)/\Gamma_{\text{total}}$ Γ_{27}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<1.2 90 AULCHENKO 08 CMD2 $\phi \rightarrow \pi^+\pi^-\gamma\gamma$

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

<5 90 AKHMETSHIN 98 CMD2 $e^+e^- \rightarrow \pi^+\pi^-\gamma\gamma$ $\Gamma(\eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{28}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

< 1.8 90 AKHMETSHIN 00E CMD2 $e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-\pi^0$

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

< 6.1 90 AULCHENKO 08 CMD2 $\phi \rightarrow \eta\pi^+\pi^-$ <30 90 AKHMETSHIN 98 CMD2 $e^+e^- \rightarrow \pi^+\pi^-\gamma\gamma$ $\Gamma(\eta\mu^+\mu^-)/\Gamma_{\text{total}}$ Γ_{29}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<9.4 90 AKHMETSHIN 01 CMD2 $e^+e^- \rightarrow \eta e^+e^-$ $\Gamma(\eta U \rightarrow \eta e^+e^-)/\Gamma_{\text{total}}$ Γ_{30}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<1 × 10⁻⁶ 90 ¹BABUSCI 13B KLOE 1.02 $e^+e^- \rightarrow \eta e^+e^-$

¹ For a narrow vector U with mass between 5 and 470 MeV, from the combined analysis of $\eta \rightarrow \pi^+ \pi^- \pi^0$ and $\eta \rightarrow \pi^0 \pi^0 \pi^0$ from ARCHILLI 12. Measured 90% CL limits as a function of m_U range from 2.2×10^{-8} to 10^{-6} .

$\Gamma(\text{invisible})/\Gamma(K^+ K^-)$					Γ_{31}/Γ_1
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<3.4 \times 10^{-4}$	90	ABLIKIM	18S BES3	$J/\psi \rightarrow \phi \eta \rightarrow \phi \pi^+ \pi^- \pi^0$	

———— Lepton Family number (LF) violating modes ————

$\Gamma(e^\pm \mu^\mp)/\Gamma_{\text{total}}$					Γ_{32}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<2 \times 10^{-6}$	90	ACHASOV	10A SND	$e^+ e^- \rightarrow e^\pm \mu^\mp$	

$\Gamma(\pi^+ \pi^+ e^- e^-)/\Gamma_{\text{total}}$					Γ_{33}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.3 \times 10^{-5}$	90	¹ ABLIKIM	25AK BES3	$J/\psi \rightarrow \phi \eta$	

¹ $B(\phi \rightarrow K^+ K^-) = (49.2 \pm 0.5)\%$ was used.

$\pi^+ \pi^- \pi^0 / \rho\pi$ AMPLITUDE RATIO a_1 IN DECAY OF $\phi \rightarrow \pi^+ \pi^- \pi^0$

NIECKNIG 12 describes final-state interactions between the three pions in a dispersive framework using data on the $\pi\pi$ P -wave scattering phase shift.

VALUE (units 10^{-2})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
9.1±1.2 OUR AVERAGE					
$10.1 \pm 4.4 \pm 1.7$		80k	¹ AKHMETSHIN 06	CMD2	$1.017-1.021 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
$9.0 \pm 1.1 \pm 0.6$		1.98M	^{2,3} ALOISIO	03 KLOE	$1.02 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$-6 < a_1 < 6$		500k	³ ACHASOV	02 SND	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
$-16 < a_1 < 11$	90	9.8k	^{1,4} AKHMETSHIN 98	CMD2	$e^+ e^- \rightarrow \pi^+ \pi^- \gamma \gamma$

¹ Dalitz plot analysis taking into account interference between the contact and $\rho\pi$ amplitudes.

² From a fit without limitations on charged and neutral ρ masses and widths.

³ Recalculated by us to match the notations of AKHMETSHIN 98.

⁴ Assuming zero phase for the contact term.

PARAMETER β IN $\phi \rightarrow P e^+ e^-$ DECAYS

In the one-pole approximation the electromagnetic transition form factor for $\phi \rightarrow P e^+ e^-$ ($P = \pi, \eta$) is given as a function of the $e^+ e^-$ invariant mass squared, q^2 , by the expression:

$$|F(q^2)|^2 = (1 - q^2/\Lambda^2)^{-2},$$

where vector meson dominance predicts parameter $\Lambda \approx 0.770$ GeV ($\Lambda^{-2} \approx 1.687$ GeV⁻²). The slope of this form factor, $\beta = dF/dq^2(q^2=0)$, equals Λ^{-2} in this approximation.

The measurements below obtain β in the one-pole approximation.

PARAMETER β IN $\phi \rightarrow \pi^0 e^+ e^-$ DECAY

VALUE (GeV^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
2.02 ± 0.11	9.5k	¹ ANASTASI	16B	KLOE $1.02 e^+ e^- \rightarrow \pi^0 e^+ e^-$

¹ The error combines statistical and systematic uncertainties.

PARAMETER β IN $\phi \rightarrow \eta e^+ e^-$ DECAY

VALUE (GeV^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
1.29 ± 0.13 OUR AVERAGE				
$1.28 \pm 0.10^{+0.09}_{-0.08}$	30k	BABUSCI	15	KLOE $1.02 e^+ e^- \rightarrow \eta e^+ e^-$
3.8 ± 1.8	213	¹ ACHASOV	01B	SND $1.02 e^+ e^- \rightarrow \eta e^+ e^-$

¹ The uncertainty is statistical only. The systematic one is negligible, in comparison.

$\phi(1020)$ REFERENCES

ABLIKIM	25A	PRL 134 011904	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	25AK	CP C49 043001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	25BR	PRL 135 161902	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	25F	PRL 134 201902	M. Ablikim <i>et al.</i>	(BESIII Collab.)
DRUZHININ	25	PPN 56 405	V.P. Druzhinin <i>et al.</i>	(SND Collab.)
AAIJ	24B	JHEP 2405 293	R. Aaij <i>et al.</i>	(LHCb Collab.)
ACHASOV	24	PR D110 072001	M.N. Achasov <i>et al.</i>	(SND Collab.)
DUBNICKA	24	PR D110 054019	S. Dubnicka <i>et al.</i>	(SLOW, CMNS)
IGNATOV	24	PR D109 112002	F.V. Ignatov <i>et al.</i>	(CMD-3 Collab.)
PDG	24	PR D110 030001	S. Navas <i>et al.</i>	(PDG Collab.)
PDG	22	PTEP 2022 083C01	R.L. Workman <i>et al.</i>	(PDG Collab.)
ABLIKIM	21AE	PR D104 012016	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21U	PR D104 032011	M. Ablikim <i>et al.</i>	(BESIII Collab.)
LEES	21B	PR D104 112003	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ALBRECHT	20	EPJ C80 453	M. Albrecht <i>et al.</i>	(Crystal Barrel Collab.)
HOID	20	EPJ C80 988	B.-L. Hoid, M. Hoferichter, B. Kubis	(BONN, BERN)
AAIJ	19H	JHEP 1904 063	R. Aaij <i>et al.</i>	(LHCb Collab.)
HOFERICH... 19	JHEP 1908 137	M. Hoferichter, B.-L. Hoid, B. Kubis	(WASH, BONN)	
ABLIKIM	18S	PR D98 032001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
AOUDE	18	PR D98 056021	R.T. Aoude <i>et al.</i>	
KOZYREV	18	PL B779 64	E.A. Kozyrev <i>et al.</i>	(CMD-3 Collab.)
AKHMETSHIN	17	PL B768 345	R.R. Akhmetshin <i>et al.</i>	(CMD-3 Collab.)
ACHASOV	16A	PR D93 092001	M.N. Achasov <i>et al.</i>	(SND Collab.)
ANASTASI	16B	PL B757 362	A. Anastasi <i>et al.</i>	(KLOE-2 Collab.)
KOZYREV	16	PL B760 314	E.A. Kozyrev <i>et al.</i>	(CMD-3 Collab.)
BABUSCI	15	PL B742 1	D. Babusci <i>et al.</i>	(KLOE-2 Collab.)
PDG	15	RPP 2015 at pdg.lbl.gov		(PDG Collab.)
LEES	14H	PR D89 092002	J.P. Lees <i>et al.</i>	(BABAR Collab.)
PDG	14	CP C38 070001	K. Olive <i>et al.</i>	(PDG Collab.)
BABUSCI	13B	PL B720 111	D. Babusci <i>et al.</i>	(KLOE-2 Collab.)
BENAYOUN	13	EPJ C73 2453	M. Benayoun, P. David, L. DelBuono	(PARIN, BERLIN+)
LEES	13F	PR D87 052010	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	13Q	PR D88 032013	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ARCHILLI	12	PL B706 251	F. Archilli <i>et al.</i>	(KLOE-2 Collab.)
BENAYOUN	12	EPJ C72 1848	M. Benayoun <i>et al.</i>	
NIECKNIG	12	EPJ C72 2014	F. Niecknig, B. Kubis, S.P. Schneider	(BONN)
PDG	12	PR D86 010001	J. Beringer <i>et al.</i>	(PDG Collab.)
AKHMETSHIN	11	PL B695 412	R.R. Akhmetshin <i>et al.</i>	(CMD-2 Collab.)
ACHASOV	10A	PR D81 057102	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
BENAYOUN	10	EPJ C65 211	M. Benayoun <i>et al.</i>	
AMBROSINO	09C	PL B679 10	F. Ambrosino <i>et al.</i>	(KLOE Collab.)
AMBROSINO	09F	PL B681 5	F. Ambrosino <i>et al.</i>	(KLOE Collab.)
AKHMETSHIN	08	PL B669 217	R.R. Akhmetshin <i>et al.</i>	(CMD-2 Collab.)
AMBROSINO	08G	PL B669 223	F. Ambrosino <i>et al.</i>	(KLOE Collab.)
AULCHENKO	08	JETPL 88 85	V. Aulchenko <i>et al.</i>	(CMD-2 Collab.)

Translated from ZETFP 88 93.

FLOREZ-BAEZ	08	PR D78 077301	F.V. Florez-Baez, G. Lopez Castro	
ACHASOV	07B	PR D76 077101	M.N. Achasov <i>et al.</i>	(SND Collab.)
AMBROSINO	07	EPJ C49 473	F. Ambrosino <i>et al.</i>	(KLOE Collab.)
AMBROSINO	07A	PL B648 267	F. Ambrosino <i>et al.</i>	(KLOE Collab.)
DUBYSKIY	07	PR D75 113001	S. Dubynskiy <i>et al.</i>	
ACHASOV	06A	PR D74 014016	M.N. Achasov <i>et al.</i>	(SND Collab.)
AKHMETSHIN	06	PL B642 203	R.R. Akhmetshin <i>et al.</i>	(CMD-2 Collab.)
AKHMETSHIN	05	PL B605 26	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
AMBROSINO	05	PL B608 199	F. Ambrosino <i>et al.</i>	(KLOE Collab.)
AUBERT,B	05J	PR D72 052008	B. Aubert <i>et al.</i>	(BABAR Collab.)
AKHMETSHIN	04	PL B578 285	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
AUBERT,B	04N	PR D70 072004	B. Aubert <i>et al.</i>	(BABAR Collab.)
ALOISIO	03	PL B561 55	A. Aloisio <i>et al.</i>	(KLOE Collab.)
AULCHENKO	03B	JETP 97 24	V.M. Aulchenko <i>et al.</i>	(Novosibirsk SND Collab.)
		Translated from ZETF 124 28.		
ACHASOV	02	PR D65 032002	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
ACHASOV	02D	JETPL 75 449	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
		Translated from ZETFP 75 539.		
ALOISIO	02C	PL B536 209	A. Aloisio <i>et al.</i>	(KLOE Collab.)
ALOISIO	02D	PL B537 21	A. Aloisio <i>et al.</i>	(KLOE Collab.)
ALOISIO	02E	PL B541 45	A. Aloisio <i>et al.</i>	(KLOE Collab.)
FISCHBACH	02	PL B526 355	E. Fischbach, A.W. Overhauser, B. Woodahl	
GOKALP	02	JP G28 2783	A. Gokalp <i>et al.</i>	
ACHASOV	01B	PL B504 275	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
ACHASOV	01E	PR D63 072002	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
ACHASOV	01F	PR D63 094007	N.N. Achasov, V.V. Gubin	(Novosibirsk SND Collab.)
ACHASOV	01G	PRL 86 1698	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
AITALA	01B	PRL 86 770	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
AKHMETSHIN	01	PL B501 191	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
AKHMETSHIN	01B	PL B509 217	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
AKHMETSHIN	01C	PL B503 237	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
BENAYOUN	01	EPJ C22 503	M. Benayoun, H.B. O'Connell	
ACHASOV	00	EPJ C12 25	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
ACHASOV	00B	JETP 90 17	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
		Translated from ZETF 117 22.		
ACHASOV	00C	PL B474 188	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
ACHASOV	00D	JETPL 72 282	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
		Translated from ZETFP 72 411.		
ACHASOV	00E	NP B569 158	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
ACHASOV	00F	PL B479 53	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
ACHASOV	00H	PL B485 349	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
AKHMETSHIN	00B	PL B473 337	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
AKHMETSHIN	00E	PL B491 81	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
AKHMETSHIN	00F	PL B494 26	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
AULCHENKO	00A	JETP 90 927	V.M. Aulchenko <i>et al.</i>	(Novosibirsk SND Collab.)
		Translated from ZETF 117 1067.		
BRAMON	00	PL B486 406	A. Bramon <i>et al.</i>	
PDG	00	EPJ C15 1	D.E. Groom <i>et al.</i>	(PDG Collab.)
ACHASOV	99	PL B449 122	M.N. Achasov <i>et al.</i>	
ACHASOV	99C	PL B456 304	M.N. Achasov <i>et al.</i>	
AKHMETSHIN	99B	PL B462 371	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
AKHMETSHIN	99C	PL B462 380	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
AKHMETSHIN	99D	PL B466 385	R.R. Akhmetshin <i>et al.</i>	
		Also PL B508 217 (errat.)	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
AKHMETSHIN	99F	PL B460 242	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
AULCHENKO	99	JETPL 69 97	V.M. Aulchenko <i>et al.</i>	
		Translated from ZETFP 69 87.		
ACHASOV	98B	PL B438 441	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
ACHASOV	98F	JETPL 68 573	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
ACHASOV	98I	PL B440 442	M.N. Achasov <i>et al.</i>	
AKHMETSHIN	98	PL B434 426	R.R. Akhmetshin <i>et al.</i>	(CMD-2 Collab.)
AULCHENKO	98	PL B436 199	V.M. Aulchenko <i>et al.</i>	(Novosibirsk SND Collab.)
BARBERIS	98	PL B432 436	D. Barberis <i>et al.</i>	(Omega Expt.)
AKHMETSHIN	97B	PL B415 445	R.R. Akhmetshin <i>et al.</i>	(NOVO, BOST, PITT+)
AKHMETSHIN	97C	PL B415 452	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
BENAYOUN	96	ZPHY C72 221	M. Benayoun <i>et al.</i>	(IPNP, NOVO)
AKHMETSHIN	95	PL B364 199	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
DOLINSKY	91	PRPL 202 99	S.I. Dolinsky <i>et al.</i>	(NOVO)
KUHN	90	ZPHY C48 445	J.H. Kuhn <i>et al.</i>	(MPIM)
ACHASOV	89	NP B315 465	N.N. Achasov, V.N. Ivanchenko	
DOLINSKY	89	ZPHY C42 511	S.I. Dolinsky <i>et al.</i>	(NOVO)
BARKOV	88	SJNP 47 248	L.M. Barkov <i>et al.</i>	(NOVO)
		Translated from YAF 47 393.		

DOLINSKY	88	SJNP 48 277	S.I. Dolinsky <i>et al.</i>	(NOVO)
		Translated from YAF 48	442.	
DRUZHININ	87	ZPHY C37 1	V.P. Druzhinin <i>et al.</i>	(NOVO)
ARMSTRONG	86	PL 166B 245	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+)
ATKINSON	86	ZPHY C30 521	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)
BEBEK	86	PRL 56 1893	C. Bebek <i>et al.</i>	(CLEO Collab.)
DAVENPORT	86	PR D33 2519	T.F. Davenport	(TUFTS, ARIZ, FNAL, FSU, NDAM+)
DIJKSTRA	86	ZPHY C31 375	H. Dijkstra <i>et al.</i>	(ANIK, BRIS, CERN+)
FRAME	86	NP B276 667	D. Frame <i>et al.</i>	(GLAS)
GOLUBEV	86	SJNP 44 409	V.B. Golubev <i>et al.</i>	(NOVO)
		Translated from YAF 44	633.	
ALBRECHT	85D	PL 153B 343	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
GOLUBEV	85	SJNP 41 756	V.B. Golubev <i>et al.</i>	(NOVO)
		Translated from YAF 41	1183.	
DRUZHININ	84	PL 144B 136	V.P. Druzhinin <i>et al.</i>	(NOVO)
ARMSTRONG	83B	NP B224 193	T.A. Armstrong <i>et al.</i>	(BARI, BIRM, CERN+)
BARATE	83	PL 121B 449	R. Barate <i>et al.</i>	(SACL, LOIC, SHMP, IND)
KURDADZE	83C	JETPL 38 366	L.M. Kurdadze <i>et al.</i>	(NOVO)
		Translated from ZETFP 38	306.	
ARENTON	82	PR D25 2241	M.W. Arenton <i>et al.</i>	(ANL, ILL)
PELLINEN	82	PS 25 599	A. Pellinen, M. Roos	(HELS)
DAUM	81	PL 100B 439	C. Daum <i>et al.</i>	(AMST, BRIS, CERN, CRAC+)
IVANOV	81	PL 107B 297	P.M. Ivanov <i>et al.</i>	(NOVO)
Also		Private Comm.	S.I. Eidelman	(NOVO)
VASSERMAN	81	PL 99B 62	I.B. Vasserman <i>et al.</i>	(NOVO)
Also		SJNP 35 240	L.M. Kurdadze <i>et al.</i>	
		Translated from YAF 35	352.	
CORDIER	80	NP B172 13	A. Cordier <i>et al.</i>	(LALO)
CORDIER	79	PL 81B 389	A. Cordier <i>et al.</i>	(LALO)
BUKIN	78B	SJNP 27 521	A.D. Bukin <i>et al.</i>	(NOVO)
		Translated from YAF 27	985.	
BUKIN	78C	SJNP 27 516	A.D. Bukin <i>et al.</i>	(NOVO)
		Translated from YAF 27	976.	
COOPER	78B	NP B146 1	A.M. Cooper <i>et al.</i>	(TATA, CERN, CDEF+)
LOSTY	78	NP B133 38	M.J. Losty <i>et al.</i>	(CERN, AMST, NIJM+)
AKERLOF	77	PRL 39 861	C.W. Akerlof <i>et al.</i>	(FNAL, MICH, PURD)
ANDREWS	77	PRL 38 198	D.E. Andrews <i>et al.</i>	(ROCH)
BALDI	77	PL 68B 381	R. Baldi <i>et al.</i>	(GEVA)
CERRADA	77B	NP B126 241	M. Cerrada <i>et al.</i>	(AMST, CERN, NIJM+)
COHEN	77	PRL 38 269	D. Cohen <i>et al.</i>	(ANL)
LAVEN	77	NP B127 43	H. Laven <i>et al.</i>	(AACH3, BERL, CERN, LOIC+)
LYONS	77	NP B125 207	L. Lyons, A.M. Cooper, A.G. Clark	(OXF)
COSME	76	PL 63B 352	G. Cosme <i>et al.</i>	(ORSAY)
KALBFLEISCH	76	PR D13 22	G.R. Kalbfleisch, R.C. Strand, J.W. Chapman	(BNL+)
PARROUR	76	PL 63B 357	G. Parrou <i>et al.</i>	(ORSAY)
PARROUR	76B	PL 63B 362	G. Parrou <i>et al.</i>	(ORSAY)
KALBFLEISCH	75	PR D11 987	G.R. Kalbfleisch, R.C. Strand, J.W. Chapman	(BNL+)
AYRES	74	PRL 32 1463	D.S. Ayres <i>et al.</i>	(ANL)
BESCH	74	NP B70 257	H.J. Besch <i>et al.</i>	(BONN)
COSME	74	PL 48B 155	G. Cosme <i>et al.</i>	(ORSAY)
COSME	74B	PL 48B 159	G. Cosme <i>et al.</i>	(ORSAY)
DEGROOT	74	NP B74 77	A.J. de Groot <i>et al.</i>	(AMST, NIJM)
AUGUSTIN	73	PRL 30 462	J.E. Augustin <i>et al.</i>	(ORSAY)
BALLAM	73	PR D7 3150	J. Ballam <i>et al.</i>	(SLAC, LBL)
BINNIE	73B	PR D8 2789	D.M. Binnie <i>et al.</i>	(LOIC, SHMP)
AGUILAR...	72B	PR D6 29	M. Aguilar-Benitez <i>et al.</i>	(BNL)
ALVENSLEB...	72	PRL 28 66	H. Alvensleben <i>et al.</i>	(MIT, DESY)
BORENSTEIN	72	PR D5 1559	S.R. Borenstein <i>et al.</i>	(BNL, MICH)
COLLEY	72	NP B50 1	D.C. Colley <i>et al.</i>	(BIRM, GLAS)
BALAKIN	71	PL 34B 328	V.E. Balakin <i>et al.</i>	(NOVO)
CHATELUS	71	Thesis LAL 1247	Y. Chatelus	(STRB)
Also		PL 32B 416	J.C. Bizot <i>et al.</i>	(ORSAY)
HAYES	71	PR D4 899	S. Hayes <i>et al.</i>	(CORN)
STOTTLE...	71	Thesis ORO 2504 170	A.R. Stottlemyer	(UMD)
BIZOT	70	PL 32B 416	J.C. Bizot <i>et al.</i>	(ORSAY)
Also		Liverpool Sym. 69	J.P. Perez-y-Jorba	
EARLES	70	PRL 25 1312	D.R. Earles <i>et al.</i>	(NEAS)
GOUNARIS	68	PRL 21 244	G.J. Gounaris, J.J. Sakurai	
LINDSEY	66	PR 147 913	J.S. Lindsey, G. Smith	(LRL)
LONDON	66	PR 143 1034	G.W. London <i>et al.</i>	(BNL, SYRA) IGJPC
BADIER	65B	PL 17 337	J. Badier <i>et al.</i>	(EPOL, SACL, AMST)

LINDSEY	65	PRL 15 221	J.S. Lindsey, G.A. Smith	(LRL)
LINDSEY 65 data included in LINDSEY 66.				
SCHLEIN	63	PRL 10 368	P.E. Schlein <i>et al.</i>	(UCLA) IGJP
