

$\Upsilon(1S)$ 

$$J^{PC} = 0^{-}(1^{-}-)$$

### $\Upsilon(1S)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>9460.40 ± 0.10 OUR AVERAGE</b>				
9460.37 ± 0.01 ± 2.85	15M	<sup>1</sup> AAIJ	24AC	LHCB $\Upsilon(1S) \rightarrow \mu^+ \mu^-$
9460.40 ± 0.09 ± 0.04		<sup>2</sup> SHAMOV	23	RVUE $e^+ e^- \rightarrow$ hadrons
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
9460.11 ± 0.11 ± 0.07		<sup>3</sup> SHAMOV	23	RVUE $e^+ e^- \rightarrow$ hadrons
9460.51 ± 0.09 ± 0.05		<sup>4,5</sup> ARTAMONOV	00	MD1 $e^+ e^- \rightarrow$ hadrons
9460.60 ± 0.09 ± 0.05		<sup>6,7</sup> BARU	92B	MD1 $e^+ e^- \rightarrow$ hadrons
9460.59 ± 0.12		BARU	86	MD1 $e^+ e^- \rightarrow$ hadrons
9460.6 ± 0.4		<sup>7,8</sup> ARTAMONOV	84	MD1 $e^+ e^- \rightarrow$ hadrons
9459.97 ± 0.11 ± 0.07		<sup>9</sup> MACKAY	84	CUSB $e^+ e^- \rightarrow$ hadrons

<sup>1</sup> Observed in prompt  $pp$  production.<sup>2</sup> Reanalysis of MD1 data using the electron mass from COHEN 87, the radiative corrections from KURAEV 85 and interference effects.<sup>3</sup> Obtained by reanalysing CUSB data (MACKAY 84), but not authored by the CUSB collaboration.<sup>4</sup> Reanalysis of BARU 92B and ARTAMONOV 84 using new electron mass (COHEN 87).<sup>5</sup> Superseded by SHAMOV 23.<sup>6</sup> Supersedes BARU 86.<sup>7</sup> Superseded by ARTAMONOV 00.<sup>8</sup> Value includes data of ARTAMONOV 82.<sup>9</sup> Reanalysed by SHAMOV 23.

### $\Upsilon(1S)$ WIDTH

VALUE (keV)	DOCUMENT ID
<b>54.02 ± 1.25 OUR EVALUATION</b>	See the Note on "Width Determinations of the $\Upsilon$ States"

### $\Upsilon(1S)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Scale factor/ Confidence level
$\Gamma_1$ $\tau^+ \tau^-$	( 2.60 ± 0.10 ) %	
$\Gamma_2$ $e^+ e^-$	( 2.39 ± 0.08 ) %	
$\Gamma_3$ $\mu^+ \mu^-$	( 2.48 ± 0.04 ) %	

#### Hadronic decays

$\Gamma_4$ $ggg$	( 81.7 ± 0.7 ) %	
$\Gamma_5$ $\gamma gg$	( 2.2 ± 0.6 ) %	
$\Gamma_6$ $\eta'(958)$ anything	( 2.94 ± 0.24 ) %	
$\Gamma_7$ $J/\psi(1S)$ anything	( 5.4 ± 0.4 ) × 10 <sup>-4</sup>	S=1.4
$\Gamma_8$ $J/\psi(1S)\eta_c$	< 2.2	× 10 <sup>-6</sup> CL=90%
$\Gamma_9$ $J/\psi(1S)\chi_{c0}$	< 3.4	× 10 <sup>-6</sup> CL=90%

$\Gamma_{10}$	$J/\psi(1S)\chi_{c1}$	$( 3.9 \pm 1.2 ) \times 10^{-6}$	
$\Gamma_{11}$	$J/\psi(1S)\chi_{c2}$	$< 1.4$	$\times 10^{-6}$ CL=90%
$\Gamma_{12}$	$J/\psi(1S)\eta_c(2S)$	$< 2.2$	$\times 10^{-6}$ CL=90%
$\Gamma_{13}$	$J/\psi(1S)X(3940)$	$< 5.4$	$\times 10^{-6}$ CL=90%
$\Gamma_{14}$	$J/\psi(1S)X(4160)$	$< 5.4$	$\times 10^{-6}$ CL=90%
$\Gamma_{15}$	$X(4350)$ anything, $X \rightarrow$ $J/\psi(1S)\phi$	$< 8.1$	$\times 10^{-6}$ CL=90%
$\Gamma_{16}$	$T_{c\bar{c}1}(3900)^\pm$ anything, $T_{c\bar{c}1} \rightarrow J/\psi(1S)\pi^\pm$	$< 1.3$	$\times 10^{-5}$ CL=90%
$\Gamma_{17}$	$T_{c\bar{c}1}(4200)^\pm$ anything, $Z_c \rightarrow$ $J/\psi(1S)\pi^\pm$	$< 6.0$	$\times 10^{-5}$ CL=90%
$\Gamma_{18}$	$T_{c\bar{c}1}(4430)^\pm$ anything, $T_{c\bar{c}1} \rightarrow J/\psi(1S)\pi^\pm$	$< 4.9$	$\times 10^{-5}$ CL=90%
$\Gamma_{19}$	$P_{c\bar{c}s}(4459)/\bar{P}_{c\bar{c}s}(4459)$ anything	$( 3.5 \pm 2.0 ) \times 10^{-6}$	
$\Gamma_{20}$	$P_{c\bar{c}s}(4338)/\bar{P}_{c\bar{c}s}(4338)$ anything	$< 1.8$	$\times 10^{-6}$ CL=90%
$\Gamma_{21}$	$P_c(4312)$ anything, $P_c(4312) \rightarrow$ $pJ/\psi(1S)$	$< 5.7$	$\times 10^{-6}$ CL=90%
$\Gamma_{22}$	$P_c(4440)$ anything, $P_c(4440) \rightarrow$ $pJ/\psi(1S)$	$< 1.00$	$\times 10^{-5}$ CL=90%
$\Gamma_{23}$	$P_c(4457)$ anything, $P_c(4457) \rightarrow$ $pJ/\psi(1S)$	$< 7.6$	$\times 10^{-6}$ CL=90%
$\Gamma_{24}$	$p/\bar{p}J/\psi(1S)$ anything	$( 8.1 \pm 0.8 ) \times 10^{-5}$	
$\Gamma_{25}$	$J/\psi(1S)\Lambda/\bar{\Lambda}$ anything	$( 3.7 \pm 0.6 ) \times 10^{-5}$	
$\Gamma_{26}$	$X_{cs}^\pm$ anything, $X \rightarrow J/\psi K^\pm$	$< 5.7$	$\times 10^{-6}$ CL=90%
$\Gamma_{27}$	$\psi(4230)$ anything, $\psi \rightarrow$ $J/\psi(1S)\pi^+\pi^-$	$< 3.8$	$\times 10^{-5}$ CL=90%
$\Gamma_{28}$	$\psi(4230)$ anything, $\psi \rightarrow$ $J/\psi(1S)K^+K^-$	$< 7.5$	$\times 10^{-6}$ CL=90%
$\Gamma_{29}$	$\chi_{c1}(4140)$ anything, $\chi_{c1} \rightarrow$ $J/\psi(1S)\phi$	$< 5.2$	$\times 10^{-6}$ CL=90%
$\Gamma_{30}$	$\chi_{c0}$ anything	$< 4$	$\times 10^{-3}$ CL=90%
$\Gamma_{31}$	$\chi_{c1}$ anything	$( 1.90 \pm 0.35 ) \times 10^{-4}$	
$\Gamma_{32}$	$\chi_{c1}(1P)X_{tetra}$	$< 3.78$	$\times 10^{-5}$ CL=90%
$\Gamma_{33}$	$\chi_{c2}$ anything	$( 2.8 \pm 0.8 ) \times 10^{-4}$	
$\Gamma_{34}$	$\psi(2S)$ anything	$( 1.23 \pm 0.20 ) \times 10^{-4}$	
$\Gamma_{35}$	$\psi(2S)\eta_c$	$< 3.6$	$\times 10^{-6}$ CL=90%
$\Gamma_{36}$	$\psi(2S)\chi_{c0}$	$< 6.5$	$\times 10^{-6}$ CL=90%
$\Gamma_{37}$	$\psi(2S)\chi_{c1}$	$< 4.5$	$\times 10^{-6}$ CL=90%
$\Gamma_{38}$	$\psi(2S)\chi_{c2}$	$< 2.1$	$\times 10^{-6}$ CL=90%
$\Gamma_{39}$	$\psi(2S)\eta_c(2S)$	$< 3.2$	$\times 10^{-6}$ CL=90%
$\Gamma_{40}$	$\psi(2S)X(3940)$	$< 2.9$	$\times 10^{-6}$ CL=90%
$\Gamma_{41}$	$\psi(2S)X(4160)$	$< 2.9$	$\times 10^{-6}$ CL=90%
$\Gamma_{42}$	$\psi(4230)$ anything, $\psi \rightarrow$ $\psi(2S)\pi^+\pi^-$	$< 7.9$	$\times 10^{-5}$ CL=90%

$\Gamma_{43}$	$\psi(4360)$ anything, $\psi \rightarrow \psi(2S)\pi^+\pi^-$	$< 5.2$	$\times 10^{-5}$	CL=90%
$\Gamma_{44}$	$\psi(4660)$ anything, $\psi \rightarrow \psi(2S)\pi^+\pi^-$	$< 2.2$	$\times 10^{-5}$	CL=90%
$\Gamma_{45}$	$T_{c\bar{c}}(4050)^\pm$ anything, $X \rightarrow \psi(2S)\pi^\pm$	$< 8.8$	$\times 10^{-5}$	CL=90%
$\Gamma_{46}$	$T_{c\bar{c}1}(4430)^\pm$ anything, $T_{c\bar{c}1} \rightarrow \psi(2S)\pi^\pm$	$< 6.7$	$\times 10^{-5}$	CL=90%
$\Gamma_{47}$	$\chi_{c1}(3872)$ anything	$< 2.2$	$\times 10^{-4}$	CL=90%
$\Gamma_{48}$	$T_{c\bar{c}1}(4200)^+ T_{c\bar{c}1}(4200)^-$	$< 2.23$	$\times 10^{-5}$	CL=90%
$\Gamma_{49}$	$T_{c\bar{c}1}(3900)^\pm T_{c\bar{c}1}(4200)^\mp$	$< 8.1$	$\times 10^{-6}$	CL=90%
$\Gamma_{50}$	$T_{c\bar{c}1}(3900)^+ T_{c\bar{c}1}(3900)^-$	$< 1.8$	$\times 10^{-6}$	CL=90%
$\Gamma_{51}$	$T_{c\bar{c}}(4050)^+ T_{c\bar{c}}(4050)^-$	$< 1.58$	$\times 10^{-5}$	CL=90%
$\Gamma_{52}$	$T_{c\bar{c}}(4250)^+ T_{c\bar{c}}(4250)^-$	$< 2.66$	$\times 10^{-5}$	CL=90%
$\Gamma_{53}$	$T_{c\bar{c}}(4050)^\pm T_{c\bar{c}}(4250)^\mp$	$< 4.42$	$\times 10^{-5}$	CL=90%
$\Gamma_{54}$	$T_{c\bar{c}1}(4430)^+ T_{c\bar{c}1}(4430)^-$	$< 2.03$	$\times 10^{-5}$	CL=90%
$\Gamma_{55}$	$T_{c\bar{c}}(4055)^\pm T_{c\bar{c}}(4055)^\mp$	$< 2.33$	$\times 10^{-5}$	CL=90%
$\Gamma_{56}$	$T_{c\bar{c}}(4055)^\pm T_{c\bar{c}1}(4430)^\mp$	$< 4.55$	$\times 10^{-5}$	CL=90%
$\Gamma_{57}$	$\rho\pi$	$< 3.68$	$\times 10^{-6}$	CL=90%
$\Gamma_{58}$	$\omega\pi^0$	$< 3.90$	$\times 10^{-6}$	CL=90%
$\Gamma_{59}$	$\pi^+\pi^-$	$< 5$	$\times 10^{-4}$	CL=90%
$\Gamma_{60}$	$K^+K^-$	$< 5$	$\times 10^{-4}$	CL=90%
$\Gamma_{61}$	$\rho\bar{\rho}$	$< 5$	$\times 10^{-4}$	CL=90%
$\Gamma_{62}$	$\pi^+\pi^-\pi^0$	$(2.1 \pm 0.8)$	$\times 10^{-6}$	
$\Gamma_{63}$	$\phi K^+K^-$	$(2.4 \pm 0.5)$	$\times 10^{-6}$	
$\Gamma_{64}$	$\omega\pi^+\pi^-$	$(4.5 \pm 1.0)$	$\times 10^{-6}$	
$\Gamma_{65}$	$K^*(892)^0 K^- \pi^+ + \text{c.c.}$	$(4.4 \pm 0.8)$	$\times 10^{-6}$	
$\Gamma_{66}$	$\phi f_2'(1525)$	$< 1.63$	$\times 10^{-6}$	CL=90%
$\Gamma_{67}$	$\omega f_2(1270)$	$< 1.79$	$\times 10^{-6}$	CL=90%
$\Gamma_{68}$	$\rho(770) a_2(1320)$	$< 2.24$	$\times 10^{-6}$	CL=90%
$\Gamma_{69}$	$K^*(892)^0 \bar{K}_2^*(1430)^0 + \text{c.c.}$	$(3.0 \pm 0.8)$	$\times 10^{-6}$	
$\Gamma_{70}$	$K_1(1270)^\pm K^\mp$	$< 2.41$	$\times 10^{-6}$	CL=90%
$\Gamma_{71}$	$K_1(1400)^\pm K^\mp$	$(1.0 \pm 0.4)$	$\times 10^{-6}$	
$\Gamma_{72}$	$b_1(1235)^\pm \pi^\mp$	$< 1.25$	$\times 10^{-6}$	CL=90%
$\Gamma_{73}$	$\pi^+\pi^-\pi^0\pi^0$	$(1.28 \pm 0.30)$	$\times 10^{-5}$	
$\Gamma_{74}$	$K_S^0 K^+ \pi^- + \text{c.c.}$	$(1.6 \pm 0.4)$	$\times 10^{-6}$	
$\Gamma_{75}$	$K^*(892)^0 \bar{K}^0 + \text{c.c.}$	$(2.9 \pm 0.9)$	$\times 10^{-6}$	
$\Gamma_{76}$	$K^*(892)^- K^+ + \text{c.c.}$	$< 1.11$	$\times 10^{-6}$	CL=90%
$\Gamma_{77}$	$f_1(1285)$ anything	$(4.6 \pm 3.1)$	$\times 10^{-3}$	
$\Gamma_{78}$	$D^*(2010)^\pm$ anything	$(2.52 \pm 0.20)$	%	
$\Gamma_{79}$	$\underline{f}_1(1285) X_{tetra}$	$< 6.24$	$\times 10^{-5}$	CL=90%
$\Gamma_{80}$	${}^2H$ anything	$(2.85 \pm 0.25)$	$\times 10^{-5}$	
$\Gamma_{81}$	Sum of 100 exclusive modes	$(1.200 \pm 0.017)$	%	

### Radiative decays

$\Gamma_{82}$	$\gamma\pi^+\pi^-$	$(6.3 \pm 1.8) \times 10^{-5}$	
$\Gamma_{83}$	$\gamma\pi^0\pi^0$	$(1.7 \pm 0.7) \times 10^{-5}$	
$\Gamma_{84}$	$\gamma\pi\pi$ (S-wave)	$(4.6 \pm 0.7) \times 10^{-5}$	
$\Gamma_{85}$	$\gamma\pi^0\eta$	$< 2.4$	$\times 10^{-6}$ CL=90%
$\Gamma_{86}$	$\gamma K^+ K^-$	[a] $(1.14 \pm 0.13) \times 10^{-5}$	
$\Gamma_{87}$	$\gamma p\bar{p}$	[b] $< 6$	$\times 10^{-6}$ CL=90%
$\Gamma_{88}$	$\gamma 2h^+ 2h^-$	$(7.0 \pm 1.5) \times 10^{-4}$	
$\Gamma_{89}$	$\gamma 3h^+ 3h^-$	$(5.4 \pm 2.0) \times 10^{-4}$	
$\Gamma_{90}$	$\gamma 4h^+ 4h^-$	$(7.4 \pm 3.5) \times 10^{-4}$	
$\Gamma_{91}$	$\gamma\pi^+\pi^- K^+ K^-$	$(2.9 \pm 0.9) \times 10^{-4}$	
$\Gamma_{92}$	$\gamma 2\pi^+ 2\pi^-$	$(2.5 \pm 0.9) \times 10^{-4}$	
$\Gamma_{93}$	$\gamma 3\pi^+ 3\pi^-$	$(2.5 \pm 1.2) \times 10^{-4}$	
$\Gamma_{94}$	$\gamma 2\pi^+ 2\pi^- K^+ K^-$	$(2.4 \pm 1.2) \times 10^{-4}$	
$\Gamma_{95}$	$\gamma\pi^+\pi^- p\bar{p}$	$(1.5 \pm 0.6) \times 10^{-4}$	
$\Gamma_{96}$	$\gamma 2\pi^+ 2\pi^- p\bar{p}$	$(4 \pm 6) \times 10^{-5}$	
$\Gamma_{97}$	$\gamma 2K^+ 2K^-$	$(2.0 \pm 2.0) \times 10^{-5}$	
$\Gamma_{98}$	$\gamma\eta'(958)$	$< 1.9$	$\times 10^{-6}$ CL=90%
$\Gamma_{99}$	$\gamma\eta$	$< 1.0$	$\times 10^{-6}$ CL=90%
$\Gamma_{100}$	$\gamma f_0(980)$	$< 3$	$\times 10^{-5}$ CL=90%
$\Gamma_{101}$	$\gamma f_2'(1525)$	$(2.9 \pm 0.6) \times 10^{-5}$	
$\Gamma_{102}$	$\gamma f_2(1270)$	$(1.01 \pm 0.06) \times 10^{-4}$	
$\Gamma_{103}$	$\gamma\eta(1405)$	$< 8.2$	$\times 10^{-5}$ CL=90%
$\Gamma_{104}$	$\gamma f_0(1500)$	$< 1.5$	$\times 10^{-5}$ CL=90%
$\Gamma_{105}$	$\gamma f_0(1500) \rightarrow \gamma K^+ K^-$	$(1.0 \pm 0.4) \times 10^{-5}$	
$\Gamma_{106}$	$\gamma f_0(1710)$	$< 2.6$	$\times 10^{-4}$ CL=90%
$\Gamma_{107}$	$\gamma f_0(1710) \rightarrow \gamma K^+ K^-$	$(1.01 \pm 0.32) \times 10^{-5}$	
$\Gamma_{108}$	$\gamma f_0(1710) \rightarrow \gamma\pi^+\pi^-$	$(5.3 \pm 2.0) \times 10^{-6}$	
$\Gamma_{109}$	$\gamma f_0(1710) \rightarrow \gamma\pi^0\pi^0$	$< 1.4$	$\times 10^{-6}$ CL=90%
$\Gamma_{110}$	$\gamma f_0(1710) \rightarrow \gamma\eta\eta$	$< 1.8$	$\times 10^{-6}$ CL=90%
$\Gamma_{111}$	$\gamma f_4(2050)$	$< 5.3$	$\times 10^{-5}$ CL=90%
$\Gamma_{112}$	$\gamma f_0(2200) \rightarrow \gamma K^+ K^-$	$< 2$	$\times 10^{-4}$ CL=90%
$\Gamma_{113}$	$\gamma f_J(2220) \rightarrow \gamma K^+ K^-$	$< 8$	$\times 10^{-7}$ CL=90%
$\Gamma_{114}$	$\gamma f_J(2220) \rightarrow \gamma\pi^+\pi^-$	$< 6$	$\times 10^{-7}$ CL=90%
$\Gamma_{115}$	$\gamma f_J(2220) \rightarrow \gamma p\bar{p}$	$< 1.1$	$\times 10^{-6}$ CL=90%
$\Gamma_{116}$	$\gamma\eta(2225) \rightarrow \gamma\phi\phi$	$< 3$	$\times 10^{-3}$ CL=90%
$\Gamma_{117}$	$\gamma\eta_c(1S)$	$< 2.9$	$\times 10^{-5}$ CL=90%
$\Gamma_{118}$	$\gamma\eta_c(2S)$	$< 4$	$\times 10^{-4}$ CL=90%
$\Gamma_{119}$	$\gamma\chi_{c0}$	$< 6.6$	$\times 10^{-5}$ CL=90%
$\Gamma_{120}$	$\gamma\chi_{c1}$	$(4.7^{+2.4}_{-1.9}) \times 10^{-5}$	
$\Gamma_{121}$	$\gamma\chi_{c2}$	$< 7.6$	$\times 10^{-6}$ CL=90%
$\Gamma_{122}$	$\gamma\chi_{c1}(3872)$	$< 4$	$\times 10^{-5}$ CL=90%
$\Gamma_{123}$	$\gamma\chi_{c1}(3872), \chi_{c1} \rightarrow \pi^+\pi^-\pi^0 J/\psi$	$< 2.8$	$\times 10^{-6}$ CL=90%

$\Gamma_{124}$	$\gamma\chi_{c0}(3915) \rightarrow \omega J/\psi$	$< 3.0$	$\times 10^{-6}$	CL=90%
$\Gamma_{125}$	$\gamma\chi_{c1}(4140) \rightarrow \phi J/\psi$	$< 2.2$	$\times 10^{-6}$	CL=90%
$\Gamma_{126}$	$\gamma X \bar{X} (m_X < 3.1 \text{ GeV})$	[c] $< 1$	$\times 10^{-3}$	CL=90%
$\Gamma_{127}$	$\gamma X \bar{X} (m_X < 4.5 \text{ GeV})$	[d] $< 2.4$	$\times 10^{-4}$	CL=90%
$\Gamma_{128}$	$\gamma X \rightarrow \gamma + \geq 4 \text{ prongs}$	[e] $< 1.78$	$\times 10^{-4}$	CL=95%
$\Gamma_{129}$	$\gamma A^0$	[f]		
$\Gamma_{130}$	$\gamma A^0 \rightarrow \gamma \mu^+ \mu^-$	[g] $< 9$	$\times 10^{-6}$	CL=90%
$\Gamma_{131}$	$\gamma A^0 \rightarrow \gamma \tau^+ \tau^-$	[a] $< 1.30$	$\times 10^{-4}$	CL=90%
$\Gamma_{132}$	$\gamma A^0 \rightarrow \gamma g g$	[h] $< 1$	%	CL=90%
$\Gamma_{133}$	$\gamma A^0 \rightarrow \gamma s \bar{s}$	[h] $< 1$	$\times 10^{-3}$	CL=90%

**Lepton Family number (LF) violating modes**

$\Gamma_{134}$	$e^\pm \mu^\mp$	LF	$< 3.9$	$\times 10^{-7}$	CL=90%
$\Gamma_{135}$	$\mu^\pm \tau^\mp$	LF	$< 2.7$	$\times 10^{-6}$	CL=90%
$\Gamma_{136}$	$e^\pm \tau^\mp$	LF	$< 2.7$	$\times 10^{-6}$	CL=90%
$\Gamma_{137}$	$\gamma e^\pm \mu^\mp$	LF	$< 4.2$	$\times 10^{-7}$	CL=90%
$\Gamma_{138}$	$\gamma \mu^\pm \tau^\mp$	LF	$< 6.1$	$\times 10^{-6}$	CL=90%
$\Gamma_{139}$	$\gamma e^\pm \tau^\mp$	LF	$< 6.5$	$\times 10^{-6}$	CL=90%

**Other decays**

$\Gamma_{140}$	invisible	$< 3.0$	$\times 10^{-4}$	CL=90%
$\Gamma_{141}$	hadrons	(96 $\pm$ 4 ) %		

[a]  $2m_\tau < M(\tau^+ \tau^-) < 9.2 \text{ GeV}$

[b]  $2 \text{ GeV} < m_{K^+ K^-} < 3 \text{ GeV}$

[c]  $X \bar{X} = \text{vectors with } m < 3.1 \text{ GeV}$

[d]  $X \text{ and } \bar{X} = \text{zero spin with } m < 4.5 \text{ GeV}$

[e]  $1.5 \text{ GeV} < m_X < 5.0 \text{ GeV}$

[f]  $A^0 = \text{scalar with } m < 8.0 \text{ GeV}$

[g]  $201 \text{ MeV} < M(\mu^+ \mu^-) < 3565 \text{ MeV}$

[h]  $0.5 \text{ GeV} < m_X < 9.0 \text{ GeV}$ , where  $m_X$  is the invariant mass of the hadronic final state.

**$\mathcal{R}(1S) \Gamma(i)\Gamma(e^+ e^-)/\Gamma(\text{total})$**

<b><math>\Gamma(\mu^+ \mu^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}</math></b>				<b><math>\Gamma_3 \Gamma_2 / \Gamma</math></b>
<u>VALUE (eV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<b>31.2<math>\pm</math>1.6<math>\pm</math>1.7</b>	KOBEL	92	CBAL	$e^+ e^- \rightarrow \mu^+ \mu^-$

<b><math>\Gamma(\text{hadrons}) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}</math></b>				<b><math>\Gamma_{141} \Gamma_2 / \Gamma</math></b>
<u>VALUE (keV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<b>1.240<math>\pm</math>0.016 OUR AVERAGE</b>				
1.252 $\pm$ 0.004 $\pm$ 0.019	<sup>1</sup> ROSNER	06	CLEO	9.5 $e^+ e^- \rightarrow \text{hadrons}$
1.187 $\pm$ 0.023 $\pm$ 0.031	<sup>1</sup> BARU	92B	MD1	$e^+ e^- \rightarrow \text{hadrons}$
1.23 $\pm$ 0.02 $\pm$ 0.05	<sup>1</sup> JAKUBOWSKI	88	CBAL	$e^+ e^- \rightarrow \text{hadrons}$

1.37 ±0.06 ±0.09	<sup>2</sup> GILES	84B	CLEO	$e^+e^- \rightarrow$	hadrons
1.23 ±0.08 ±0.04	<sup>2</sup> ALBRECHT	82	DASP	$e^+e^- \rightarrow$	hadrons
1.13 ±0.07 ±0.11	<sup>2</sup> NICZYPORUK	82	LENA	$e^+e^- \rightarrow$	hadrons
1.09 ±0.25	<sup>2</sup> BOCK	80	CNTR	$e^+e^- \rightarrow$	hadrons
1.35 ±0.14	<sup>3</sup> BERGER	79	PLUT	$e^+e^- \rightarrow$	hadrons

<sup>1</sup> Radiative corrections evaluated following KURAEV 85.

<sup>2</sup> Radiative corrections reevaluated by BUCHMUELLER 88 following KURAEV 85.

<sup>3</sup> Radiative corrections reevaluated by ALEXANDER 89 using  $B(\mu\mu) = 0.026$ .

## $\Upsilon(1S)$ PARTIAL WIDTHS

$\Gamma(e^+e^-)$

$\Gamma_2$

VALUE (keV)

DOCUMENT ID

**1.340±0.018 OUR EVALUATION**

## $\Upsilon(1S)$ BRANCHING RATIOS

$\Gamma(\tau^+\tau^-)/\Gamma_{\text{total}}$

$\Gamma_1/\Gamma$

VALUE (units  $10^{-2}$ )

EVTS

DOCUMENT ID

TECN

COMMENT

**2.60±0.10 OUR AVERAGE**

2.53±0.13±0.04    60k    <sup>1</sup> BESSON    07    CLEO     $e^+e^- \rightarrow \Upsilon(1S) \rightarrow \tau^+\tau^-$

2.61±0.12<sup>+0.09</sup><sub>-0.13</sub>    25k    CINABRO    94B    CLE2     $e^+e^- \rightarrow \tau^+\tau^-$

2.7 ±0.4 ±0.2    <sup>2</sup> ALBRECHT    85C    ARG     $\Upsilon(2S) \rightarrow \pi^+\pi^-\tau^+\tau^-$

3.4 ±0.4 ±0.4    GILES    83    CLEO     $e^+e^- \rightarrow \tau^+\tau^-$

<sup>1</sup> BESSON 07 reports  $[\Gamma(\Upsilon(1S) \rightarrow \tau^+\tau^-)/\Gamma_{\text{total}}] / [B(\Upsilon(1S) \rightarrow \mu^+\mu^-)] = 1.02 \pm 0.02 \pm 0.05$  which we multiply by our best (shown rounded) value  $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.04) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best (shown rounded) value.

<sup>2</sup> Using  $B(\Upsilon(1S) \rightarrow ee) = B(\Upsilon(1S) \rightarrow \mu\mu) = 0.0256$ ; not used for width evaluations.

$\Gamma(e^+e^-)/\Gamma_{\text{total}}$

$\Gamma_2/\Gamma$

VALUE (units  $10^{-2}$ )

EVTS

DOCUMENT ID

TECN

COMMENT

**2.39±0.08 OUR AVERAGE**

2.40±0.01±0.12    191k    PATRA    22    BELL     $\Upsilon(2S) \rightarrow \pi^+\pi^-e^+e^-$

2.29±0.08±0.11    ALEXANDER    98    CLE2     $\Upsilon(2S) \rightarrow \pi^+\pi^-e^+e^-$

2.42±0.14±0.14    307    ALBRECHT    87    ARG     $\Upsilon(2S) \rightarrow \pi^+\pi^-e^+e^-$

2.8 ±0.3 ±0.2    826    BESSON    84    CLEO     $\Upsilon(2S) \rightarrow \pi^+\pi^-e^+e^-$

5.1 ±3.0    BERGER    80C    PLUT     $e^+e^- \rightarrow e^+e^-$

$\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$

$\Gamma_3/\Gamma$

VALUE (units  $10^{-2}$ )

EVTS

DOCUMENT ID

TECN

COMMENT

**2.48±0.04 OUR AVERAGE**

2.46±0.01±0.11    246k    PATRA    22    BELL     $\Upsilon(2S) \rightarrow \pi^+\pi^-\mu^+\mu^-$

2.49±0.02±0.07    345k    ADAMS    05    CLEO     $e^+e^- \rightarrow \mu^+\mu^-$

2.49±0.08±0.13    ALEXANDER    98    CLE2     $\Upsilon(2S) \rightarrow \pi^+\pi^-\mu^+\mu^-$

2.12±0.20±0.10    <sup>1</sup> BARU    92    MD1     $e^+e^- \rightarrow \mu^+\mu^-$

$2.31 \pm 0.12 \pm 0.10$		<sup>1</sup> KOBEL	92	CBAL	$e^+ e^- \rightarrow \mu^+ \mu^-$
$2.52 \pm 0.07 \pm 0.07$		CHEN	89B	CLEO	$e^+ e^- \rightarrow \mu^+ \mu^-$
$2.61 \pm 0.09 \pm 0.11$		KAARSBERG	89	CSB2	$e^+ e^- \rightarrow \mu^+ \mu^-$
$2.30 \pm 0.25 \pm 0.13$	86	ALBRECHT	87	ARG	$\Upsilon(2S) \rightarrow \pi^+ \pi^- \mu^+ \mu^-$
$2.9 \pm 0.3 \pm 0.2$	864	BESSON	84	CLEO	$\Upsilon(2S) \rightarrow \pi^+ \pi^- \mu^+ \mu^-$
$2.7 \pm 0.3 \pm 0.3$		ANDREWS	83	CLEO	$e^+ e^- \rightarrow \mu^+ \mu^-$
$3.2 \pm 1.3 \pm 0.3$		ALBRECHT	82	DASP	$e^+ e^- \rightarrow \mu^+ \mu^-$
$3.8 \pm 1.5 \pm 0.2$		NICZYPORUK	82	LENA	$e^+ e^- \rightarrow \mu^+ \mu^-$
$1.4 \begin{smallmatrix} +3.4 \\ -1.4 \end{smallmatrix}$		BOCK	80	CNTR	$e^+ e^- \rightarrow \mu^+ \mu^-$
$2.2 \pm 2.0$		BERGER	79	PLUT	$e^+ e^- \rightarrow \mu^+ \mu^-$

<sup>1</sup> Taking into account interference between the resonance and continuum.

### $\Gamma(\tau^+ \tau^-) / \Gamma(\mu^+ \mu^-)$ $\Gamma_1 / \Gamma_3$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>1.008 ± 0.023 OUR AVERAGE</b>				
$1.005 \pm 0.013 \pm 0.022$	0.7M	<sup>1</sup> DEL-AMO-SA..10C	BABR	$\Upsilon(3S) \rightarrow \pi^+ \pi^- \Upsilon(1S)$
$1.02 \pm 0.02 \pm 0.05$	60k	BESSON	07	CLEO $e^+ e^- \rightarrow \Upsilon(1S)$

<sup>1</sup> Allows any number of extra photons with total energy < 500 MeV.

### $\Gamma(g g g) / \Gamma_{\text{total}}$ $\Gamma_4 / \Gamma$

<u>VALUE (units <math>10^{-2}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>81.7 ± 0.7</b>	20M	<sup>1</sup> BESSON	06A	CLEO $\Upsilon(1S) \rightarrow \text{hadrons}$

<sup>1</sup> Calculated using the value  $\Gamma(\gamma g g) / \Gamma(g g g) = (2.70 \pm 0.01 \pm 0.13 \pm 0.24)\%$  from BESSON 06A and PDG 08 values of  $B(\mu^+ \mu^-) = (2.48 \pm 0.05)\%$  and  $R_{\text{hadrons}} = 3.51$ . The statistical error is negligible and the systematic error is partially correlated with that of  $\Gamma(\gamma g g) / \Gamma_{\text{total}}$  measurement of BESSON 06A.

### $\Gamma(\gamma g g) / \Gamma_{\text{total}}$ $\Gamma_5 / \Gamma$

<u>VALUE (units <math>10^{-2}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>2.20 ± 0.60</b>	400k	<sup>1</sup> BESSON	06A	CLEO $\Upsilon(1S) \rightarrow \gamma + \text{hadrons}$

<sup>1</sup> Calculated using BESSON 06A values of  $\Gamma(\gamma g g) / \Gamma(g g g) = (2.70 \pm 0.01 \pm 0.13 \pm 0.24)\%$  and  $\Gamma(g g g) / \Gamma_{\text{total}}$ . The statistical error is negligible and the systematic error is partially correlated with that of  $\Gamma(g g g) / \Gamma_{\text{total}}$  measurement of BESSON 06A.

### $\Gamma(\gamma g g) / \Gamma(g g g)$ $\Gamma_5 / \Gamma_4$

<u>VALUE (units <math>10^{-2}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>2.70 ± 0.01 ± 0.27</b>	20M	BESSON	06A	CLEO $\Upsilon(1S) \rightarrow (\gamma +) \text{hadrons}$

### $\Gamma(\eta'(958) \text{ anything}) / \Gamma_{\text{total}}$ $\Gamma_6 / \Gamma$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.0294 ± 0.0024 OUR AVERAGE</b>			
$0.030 \pm 0.002 \pm 0.002$	AQUINES	06A	CLE3 $\Upsilon(1S) \rightarrow \eta' \text{ anything}$
$0.028 \pm 0.004 \pm 0.002$	ARTUSO	03	CLE2 $\Upsilon(1S) \rightarrow \eta' \text{ anything}$

$\Gamma(J/\psi(1S) \text{ anything})/\Gamma_{\text{total}}$   $\Gamma_7/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>5.4 \pm 0.4</math></b>	<b>OUR FIT</b>	Error includes scale factor of 1.4.			
<b><math>5.4 \pm 0.4</math></b>	<b>OUR AVERAGE</b>	Error includes scale factor of 1.5.			
$5.25 \pm 0.13 \pm 0.25$		3k	SHEN	16	BELL $e^+e^- \rightarrow J/\psi X$
$6.4 \pm 0.4 \pm 0.6$		730	BRIERE	04	CLEO $e^+e^- \rightarrow J/\psi X$
$11 \pm 4 \pm 2$			<sup>1</sup> FULTON	89	CLEO $e^+e^- \rightarrow \mu^+\mu^- X$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
<6.8	90		ALBRECHT	92J	ARG $e^+e^- \rightarrow e^+e^- X, \mu^+\mu^- X$
<17	90		MASCHMANN	90	CBAL $e^+e^- \rightarrow \text{hadrons}$
<200	90		NICZYPORUK	83	LENA

<sup>1</sup> Using  $B((J/\psi) \rightarrow \mu^+\mu^-) = (6.9 \pm 0.9)\%$ .

 $\Gamma(J/\psi(1S)\eta_c)/\Gamma_{\text{total}}$   $\Gamma_8/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< <b><math>2.2 \times 10^{-6}</math></b>	90	YANG	14	BELL $e^+e^- \rightarrow J/\psi X$

 $\Gamma(J/\psi(1S)\chi_{c0})/\Gamma_{\text{total}}$   $\Gamma_9/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< <b><math>3.4 \times 10^{-6}</math></b>	90	YANG	14	BELL $e^+e^- \rightarrow J/\psi X$

 $\Gamma(J/\psi(1S)\chi_{c1})/\Gamma_{\text{total}}$   $\Gamma_{10}/\Gamma$ 

VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>3.90 \pm 1.21 \pm 0.23</math></b>	20	YANG	14	BELL $e^+e^- \rightarrow J/\psi X$

 $\Gamma(J/\psi(1S)\chi_{c2})/\Gamma_{\text{total}}$   $\Gamma_{11}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< <b><math>1.4 \times 10^{-6}</math></b>	90	YANG	14	BELL $e^+e^- \rightarrow J/\psi X$

 $\Gamma(J/\psi(1S)\eta_c(2S))/\Gamma_{\text{total}}$   $\Gamma_{12}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< <b><math>2.2 \times 10^{-6}</math></b>	90	YANG	14	BELL $e^+e^- \rightarrow J/\psi X$

 $\Gamma(J/\psi(1S)X(3940))/\Gamma_{\text{total}}$   $\Gamma_{13}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< <b><math>5.4 \times 10^{-6}</math></b>	90	YANG	14	BELL $e^+e^- \rightarrow J/\psi X$

 $\Gamma(J/\psi(1S)X(4160))/\Gamma_{\text{total}}$   $\Gamma_{14}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< <b><math>5.4 \times 10^{-6}</math></b>	90	YANG	14	BELL $e^+e^- \rightarrow J/\psi X$

 $\Gamma(X(4350) \text{ anything}, X \rightarrow J/\psi(1S)\phi)/\Gamma_{\text{total}}$   $\Gamma_{15}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< <b><math>8.1 \times 10^{-6}</math></b>	90	SHEN	16	BELL $\Upsilon(1S) \rightarrow J/\psi K^+ K^- X$

 $\Gamma(T_{c\bar{c}1}(3900)^\pm \text{ anything}, T_{c\bar{c}1} \rightarrow J/\psi(1S)\pi^\pm)/\Gamma_{\text{total}}$   $\Gamma_{16}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< <b><math>1.3 \times 10^{-5}</math></b>	90	SHEN	16	BELL $\Upsilon(1S) \rightarrow J/\psi \pi^\pm X$

$\Gamma(T_{c\bar{c}1}(4200)^\pm \text{ anything}, Z_c \rightarrow J/\psi(1S)\pi^\pm)/\Gamma_{\text{total}}$					$\Gamma_{17}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<6.0 \times 10^{-5}$	90	SHEN	16	BELL	$\Upsilon(1S) \rightarrow J/\psi\pi^\pm X$
$\Gamma(T_{c\bar{c}1}(4430)^\pm \text{ anything}, T_{c\bar{c}1} \rightarrow J/\psi(1S)\pi^\pm)/\Gamma_{\text{total}}$					$\Gamma_{18}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<4.9 \times 10^{-5}$	90	SHEN	16	BELL	$\Upsilon(1S) \rightarrow J/\psi\pi^\pm X$
$\Gamma(P_{c\bar{c}s}(4459)/\bar{P}_{c\bar{c}s}(4459) \text{ anything})/\Gamma_{\text{total}}$					$\Gamma_{19}/\Gamma$
VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	
$3.5 \pm 2.0 \pm 0.2$	21	<sup>1</sup> ADACHI	25A	BELL	$e^+e^- \rightarrow \Upsilon(1S)$
<sup>1</sup> $21 \pm 5$ corresponds to the number of $P_{c\bar{c}s}(4459)$ events in the combined $\Upsilon(1S)$ and $\Upsilon(2S)$ samples.					
$\Gamma(P_{c\bar{c}s}(4338)/\bar{P}_{c\bar{c}s}(4338) \text{ anything})/\Gamma_{\text{total}}$					$\Gamma_{20}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.8 \times 10^{-6}$	90	ADACHI	25A	BELL	$e^+e^- \rightarrow \Upsilon(1S)$
$\Gamma(P_c(4312) \text{ anything}, P_c(4312) \rightarrow pJ/\psi(1S))/\Gamma_{\text{total}}$					$\Gamma_{21}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<5.7 \times 10^{-6}$	90	DONG	25	BELL	$e^+e^- \rightarrow \Upsilon(1S)$
$\Gamma(P_c(4440) \text{ anything}, P_c(4440) \rightarrow pJ/\psi(1S))/\Gamma_{\text{total}}$					$\Gamma_{22}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<10.0 \times 10^{-6}$	90	DONG	25	BELL	$e^+e^- \rightarrow \Upsilon(1S)$
$\Gamma(P_c(4457) \text{ anything}, P_c(4457) \rightarrow pJ/\psi(1S))/\Gamma_{\text{total}}$					$\Gamma_{23}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<7.6 \times 10^{-6}$	90	DONG	25	BELL	$e^+e^- \rightarrow \Upsilon(1S)$
$\Gamma(p/\bar{p}J/\psi(1S) \text{ anything})/\Gamma_{\text{total}}$					$\Gamma_{24}/\Gamma$
VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	
$8.1 \pm 0.6 \pm 0.5$	377	DONG	25	BELL	$e^+e^- \rightarrow \Upsilon(1S)$
$\Gamma(J/\psi(1S)\Lambda/\bar{\Lambda} \text{ anything})/\Gamma_{\text{total}}$					$\Gamma_{25}/\Gamma$
VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	
$36.9 \pm 5.3 \pm 2.4$	84	ADACHI	25A	BELL	$e^+e^- \rightarrow \Upsilon(1S)$
$\Gamma(X_{cs}^\pm \text{ anything}, X \rightarrow J/\psi K^\pm)/\Gamma_{\text{total}}$					$\Gamma_{26}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<5.7 \times 10^{-6}$	90	SHEN	16	BELL	$\Upsilon(1S) \rightarrow J/\psi K^- X$
$\Gamma(\psi(4230) \text{ anything}, \psi \rightarrow J/\psi(1S)\pi^+\pi^-)/\Gamma_{\text{total}}$					$\Gamma_{27}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<3.8 \times 10^{-5}$	90	SHEN	16	BELL	$\Upsilon(1S) \rightarrow J/\psi\pi^+\pi^- X$
$\Gamma(\psi(4230) \text{ anything}, \psi \rightarrow J/\psi(1S)K^+K^-)/\Gamma_{\text{total}}$					$\Gamma_{28}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<7.5 \times 10^{-6}$	90	SHEN	16	BELL	$\Upsilon(1S) \rightarrow J/\psi K^+ K^- X$

$\Gamma(\chi_{c1}(4140) \text{ anything}, \chi_{c1} \rightarrow J/\psi(1S)\phi)/\Gamma_{\text{total}}$   $\Gamma_{29}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5.2 \times 10^{-6}$	90	SHEN	16	BELL $\Upsilon(1S) \rightarrow J/\psi K^+ K^- X$

 $\Gamma(\chi_{c0} \text{ anything})/\Gamma(J/\psi(1S) \text{ anything})$   $\Gamma_{30}/\Gamma_7$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<7.4$	90	BRIERE	04	CLEO $e^+ e^- \rightarrow J/\psi X$

 $\Gamma(\chi_{c1} \text{ anything})/\Gamma_{\text{total}}$   $\Gamma_{31}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>1.90 \pm 0.35</math> OUR FIT</b>				
<b><math>1.90 \pm 0.43 \pm 0.14</math></b>	215	JIA	17	BELL $\Upsilon(1S) \rightarrow \gamma J/\psi(1S)$

 $\Gamma(\chi_{c1} \text{ anything})/\Gamma(J/\psi(1S) \text{ anything})$   $\Gamma_{31}/\Gamma_7$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>0.35 \pm 0.07</math> OUR FIT</b>				
<b><math>0.35 \pm 0.08 \pm 0.06</math></b>	$52 \pm 12$	BRIERE	04	CLEO $e^+ e^- \rightarrow J/\psi X$

 $\Gamma(\chi_{c1}(1P) X_{tetra})/\Gamma_{\text{total}}$   $\Gamma_{32}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<37.8 \times 10^{-6}$	90	<sup>1</sup> JIA	17A	BELL $e^+ e^- \rightarrow \text{hadrons}$

<sup>1</sup> For a tetraquark state  $X_{tetra}$ , with mass in the range 1.16–2.46 GeV and width in the range 0–0.3 GeV. Measured 90% CL limits as a function of  $X_{tetra}$  mass and width range from  $4.4 \times 10^{-6}$  to  $37.8 \times 10^{-6}$ .

 $\Gamma(\chi_{c2} \text{ anything})/\Gamma(J/\psi(1S) \text{ anything})$   $\Gamma_{33}/\Gamma_7$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>0.52 \pm 0.12 \pm 0.09</math></b>	$47 \pm 11$	BRIERE	04	CLEO $e^+ e^- \rightarrow J/\psi X$

 $\Gamma(\psi(2S) \text{ anything})/\Gamma_{\text{total}}$   $\Gamma_{34}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>1.23 \pm 0.17 \pm 0.11</math></b>	215	SHEN	16	BELL $e^+ e^- \rightarrow \psi(2S) X$

 $\Gamma(\psi(2S) \text{ anything})/\Gamma(J/\psi(1S) \text{ anything})$   $\Gamma_{34}/\Gamma_7$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>0.41 \pm 0.11 \pm 0.08</math></b>	$42 \pm 11$	BRIERE	04	CLEO $e^+ e^- \rightarrow J/\psi \pi^+ \pi^- X$

 $\Gamma(\psi(2S) \eta_c)/\Gamma_{\text{total}}$   $\Gamma_{35}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.6 \times 10^{-6}$	90	YANG	14	BELL $e^+ e^- \rightarrow \psi(2S) X$

 $\Gamma(\psi(2S) \chi_{c0})/\Gamma_{\text{total}}$   $\Gamma_{36}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<6.5 \times 10^{-6}$	90	YANG	14	BELL $e^+ e^- \rightarrow \psi(2S) X$

 $\Gamma(\psi(2S) \chi_{c1})/\Gamma_{\text{total}}$   $\Gamma_{37}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<4.5 \times 10^{-6}$	90	YANG	14	BELL $e^+ e^- \rightarrow \psi(2S) X$

$\Gamma(\psi(2S)\chi_{c2})/\Gamma_{\text{total}}$					$\Gamma_{38}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<2.1 \times 10^{-6}$	90	YANG 14	BELL	$e^+e^- \rightarrow \psi(2S)X$	
$\Gamma(\psi(2S)\eta_c(2S))/\Gamma_{\text{total}}$					$\Gamma_{39}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<3.2 \times 10^{-6}$	90	YANG 14	BELL	$e^+e^- \rightarrow \psi(2S)X$	
$\Gamma(\psi(2S)X(3940))/\Gamma_{\text{total}}$					$\Gamma_{40}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<2.9 \times 10^{-6}$	90	YANG 14	BELL	$e^+e^- \rightarrow \psi(2S)X$	
$\Gamma(\psi(2S)X(4160))/\Gamma_{\text{total}}$					$\Gamma_{41}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<2.9 \times 10^{-6}$	90	YANG 14	BELL	$e^+e^- \rightarrow \psi(2S)X$	
$\Gamma(\psi(4230) \text{ anything, } \psi \rightarrow \psi(2S)\pi^+\pi^-)/\Gamma_{\text{total}}$					$\Gamma_{42}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<7.9 \times 10^{-5}$	90	SHEN 16	BELL	$\Upsilon(1S) \rightarrow \psi(2S)\pi^+\pi^-X$	
$\Gamma(\psi(4360) \text{ anything, } \psi \rightarrow \psi(2S)\pi^+\pi^-)/\Gamma_{\text{total}}$					$\Gamma_{43}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<5.2 \times 10^{-5}$	90	SHEN 16	BELL	$\Upsilon(1S) \rightarrow \psi(2S)\pi^+\pi^-X$	
$\Gamma(\psi(4660) \text{ anything, } \psi \rightarrow \psi(2S)\pi^+\pi^-)/\Gamma_{\text{total}}$					$\Gamma_{44}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<2.2 \times 10^{-5}$	90	SHEN 16	BELL	$\Upsilon(1S) \rightarrow \psi(2S)\pi^+\pi^-X$	
$\Gamma(T_{c\bar{c}}(4050)^\pm \text{ anything, } X \rightarrow \psi(2S)\pi^\pm)/\Gamma_{\text{total}}$					$\Gamma_{45}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<8.8 \times 10^{-5}$	90	SHEN 16	BELL	$\Upsilon(1S) \rightarrow \psi(2S)\pi^\pm X$	
$\Gamma(T_{c\bar{c}}(4430)^\pm \text{ anything, } T_{c\bar{c}} \rightarrow \psi(2S)\pi^\pm)/\Gamma_{\text{total}}$					$\Gamma_{46}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<6.7 \times 10^{-5}$	90	SHEN 16	BELL	$\Upsilon(1S) \rightarrow \psi(2S)\pi^\pm X$	
$\Gamma(\chi_{c1}(3872) \text{ anything})/\Gamma_{\text{total}}$					$\Gamma_{47}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<2.2 \times 10^{-4}$	90	<sup>1</sup> SHEN 16	BELL	$\Upsilon(1S) \rightarrow J/\psi\pi^+\pi^-X$	
<sup>1</sup> SHEN 16 reports $[\Gamma(\Upsilon(1S) \rightarrow \chi_{c1}(3872) \text{ anything})/\Gamma_{\text{total}}] \times [B(\chi_{c1}(3872) \rightarrow \pi^+\pi^-J/\psi(1S))] < 9.5 \times 10^{-6}$ which we divide by our best (shown rounded) value $B(\chi_{c1}(3872) \rightarrow \pi^+\pi^-J/\psi(1S)) = 4.3 \times 10^{-2}$ .					
$\Gamma(T_{c\bar{c}}(4200)^+ T_{c\bar{c}}(4200)^-)/\Gamma_{\text{total}}$					$\Gamma_{48}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<22.3 \times 10^{-6}$	90	<sup>1</sup> JIA 18	BELL	$\Upsilon(1S) \rightarrow J/\psi\pi^\pm X$	
<sup>1</sup> Assuming $B(T_{c\bar{c}}(4200)^\pm \rightarrow J/\psi\pi^\pm) = 1$ .					

$\Gamma(T_{c\bar{c}1}(3900)^\pm T_{c\bar{c}1}(4200)^\mp)/\Gamma_{\text{total}}$   $\Gamma_{49}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<8.1 \times 10^{-6}$	90	<sup>1</sup> JIA	18	BELL $\Upsilon(1S) \rightarrow J/\psi \pi^\pm X$

<sup>1</sup> Assuming  $B(T_{c\bar{c}1}(4200)^\pm \rightarrow J/\psi \pi^\pm) = 1 = B(T_{c\bar{c}1}(3900)^\pm \rightarrow J/\psi \pi^\pm)$ .

 $\Gamma(T_{c\bar{c}1}(3900)^+ T_{c\bar{c}1}(3900)^-)/\Gamma_{\text{total}}$   $\Gamma_{50}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.8 \times 10^{-6}$	90	<sup>1</sup> JIA	18	BELL $\Upsilon(1S) \rightarrow J/\psi \pi^\pm X$

<sup>1</sup> Assuming  $B(T_{c\bar{c}1}(3900)^\pm \rightarrow J/\psi \pi^\pm) = 1$

 $\Gamma(T_{c\bar{c}}(4050)^+ T_{c\bar{c}}(4050)^-)/\Gamma_{\text{total}}$   $\Gamma_{51}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<15.8 \times 10^{-6}$	90	<sup>1</sup> JIA	18	BELL $\Upsilon(1S) \rightarrow \chi_{c1}(1P) \pi^\pm X$

<sup>1</sup> Assuming  $B(T_{c\bar{c}}(4050)^\pm \rightarrow \chi_{c1}(1P) \pi^\pm) = 1$

 $\Gamma(T_{c\bar{c}}(4250)^+ T_{c\bar{c}}(4250)^-)/\Gamma_{\text{total}}$   $\Gamma_{52}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<26.6 \times 10^{-6}$	90	<sup>1</sup> JIA	18	BELL $\Upsilon(1S) \rightarrow \chi_{c1}(1P) \pi^\pm X$

<sup>1</sup> Assuming  $B(T_{c\bar{c}}(4250)^\pm \rightarrow \chi_{c1}(1P) \pi^\pm) = 1$

 $\Gamma(T_{c\bar{c}}(4050)^\pm T_{c\bar{c}}(4250)^\mp)/\Gamma_{\text{total}}$   $\Gamma_{53}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<44.2 \times 10^{-6}$	90	<sup>1</sup> JIA	18	BELL $\Upsilon(1S) \rightarrow \chi_{c1}(1P) \pi^\pm X$

<sup>1</sup> Assuming  $B(T_{c\bar{c}}(4050)^\pm \rightarrow \chi_{c1}(1P) \pi^\pm) = 1 = B(T_{c\bar{c}}(4250)^\pm \rightarrow \chi_{c1}(1P) \pi^\pm)$

 $\Gamma(T_{c\bar{c}1}(4430)^+ T_{c\bar{c}1}(4430)^-)/\Gamma_{\text{total}}$   $\Gamma_{54}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<20.3 \times 10^{-6}$	90	<sup>1</sup> JIA	18	BELL $\Upsilon(2S) \rightarrow \psi(2S) \pi^\pm X$

<sup>1</sup> Assuming  $B(T_{c\bar{c}1}(4430)^\pm \rightarrow \psi(2S) \pi^\pm) = 1$

 $\Gamma(T_{c\bar{c}}(4055)^\pm T_{c\bar{c}}(4055)^\mp)/\Gamma_{\text{total}}$   $\Gamma_{55}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<23.3 \times 10^{-6}$	90	<sup>1</sup> JIA	18	BELL $\Upsilon(1S) \rightarrow \psi(2S) \pi^\pm X$

<sup>1</sup> Assuming  $B(T_{c\bar{c}}(4055)^\pm \rightarrow \psi(2S) \pi^\pm) = 1$

 $\Gamma(T_{c\bar{c}}(4055)^\pm T_{c\bar{c}1}(4430)^\mp)/\Gamma_{\text{total}}$   $\Gamma_{56}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<45.5 \times 10^{-6}$	90	<sup>1</sup> JIA	18	BELL $\Upsilon(1S) \rightarrow \psi(2S) \pi^\pm X$

<sup>1</sup> Assuming  $B(T_{c\bar{c}}(4055)^\pm \rightarrow \psi(2S) \pi^\pm) = 1 = B(T_{c\bar{c}1}(4430)^\pm \rightarrow \psi(2S) \pi^\pm)$

 $\Gamma(\rho\pi)/\Gamma_{\text{total}}$   $\Gamma_{57}/\Gamma$ 

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$<3.68$	90	SHEN	13	BELL $\Upsilon(1S) \rightarrow \pi^+ \pi^- \pi^0$
$<1 \times 10^3$	90	BLINOV	90	MD1 $\Upsilon(1S) \rightarrow \rho^0 \pi^0$
$<2 \times 10^2$	90	FULTON	90B	$\Upsilon(1S) \rightarrow \rho^0 \pi^0$
$<2.1 \times 10^3$	90	NICZYPORUK	83	LENA $\Upsilon(1S) \rightarrow \rho^0 \pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$\Gamma(\omega\pi^0)/\Gamma_{\text{total}}$					$\Gamma_{58}/\Gamma$
<u>VALUE (units <math>10^{-6}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<b>&lt;3.90</b>	90	SHEN	13	BELL	$\Upsilon(1S) \rightarrow \pi^+\pi^-\pi^0\pi^0$

$\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$					$\Gamma_{59}/\Gamma$
<u>VALUE (units <math>10^{-4}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<b>&lt;5</b>	90	BARU	92	MD1	$\Upsilon(1S) \rightarrow \pi^+\pi^-$

$\Gamma(K^+K^-)/\Gamma_{\text{total}}$					$\Gamma_{60}/\Gamma$
<u>VALUE (units <math>10^{-4}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<b>&lt;5</b>	90	BARU	92	MD1	$\Upsilon(1S) \rightarrow K^+K^-$

$\Gamma(\rho\bar{\rho})/\Gamma_{\text{total}}$					$\Gamma_{61}/\Gamma$
<u>VALUE (units <math>10^{-4}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<b>&lt;5</b>	90	<sup>1</sup> BARU	96	MD1	$\Upsilon(1S) \rightarrow \rho\bar{\rho}$

<sup>1</sup>Supersedes BARU 92 in this node.

$\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$						$\Gamma_{62}/\Gamma$
<u>VALUE (units <math>10^{-6}</math>)</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<b><math>2.14 \pm 0.72 \pm 0.34</math></b>		$26 \pm 9$	SHEN	13	BELL	$\Upsilon(1S) \rightarrow \pi^+\pi^-\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<18.4	90		ANASTASSOV 99	CLE2	$e^+e^- \rightarrow \text{hadrons}$	

$\Gamma(\phi K^+K^-)/\Gamma_{\text{total}}$					$\Gamma_{63}/\Gamma$
<u>VALUE (units <math>10^{-6}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<b><math>2.36 \pm 0.37 \pm 0.29</math></b>	56	SHEN	12A	BELL	$\Upsilon(1S) \rightarrow 2(K^+K^-)$

$\Gamma(\omega\pi^+\pi^-)/\Gamma_{\text{total}}$					$\Gamma_{64}/\Gamma$
<u>VALUE (units <math>10^{-6}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<b><math>4.46 \pm 0.67 \pm 0.72</math></b>	64	SHEN	12A	BELL	$\Upsilon(1S) \rightarrow 2(\pi^+\pi^-)\pi^0$

$\Gamma(K^*(892)^0 K^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$					$\Gamma_{65}/\Gamma$
<u>VALUE (units <math>10^{-6}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<b><math>4.42 \pm 0.50 \pm 0.58</math></b>	173	SHEN	12A	BELL	$\Upsilon(1S) \rightarrow K^+K^-\pi^+\pi^-$

$\Gamma(\phi f'_2(1525))/\Gamma_{\text{total}}$					$\Gamma_{66}/\Gamma$
<u>VALUE (units <math>10^{-6}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<b>&lt;1.63</b>	90	SHEN	12A	BELL	$\Upsilon(1S) \rightarrow 2(K^+K^-)$

$\Gamma(\omega f_2(1270))/\Gamma_{\text{total}}$					$\Gamma_{67}/\Gamma$
<u>VALUE (units <math>10^{-6}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<b>&lt;1.79</b>	90	SHEN	12A	BELL	$\Upsilon(1S) \rightarrow 2(\pi^+\pi^-)\pi^0$

$\Gamma(\rho(770) a_2(1320))/\Gamma_{\text{total}}$					$\Gamma_{68}/\Gamma$
<u>VALUE (units <math>10^{-6}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<b>&lt;2.24</b>	90	SHEN	12A	BELL	$\Upsilon(1S) \rightarrow 2(\pi^+\pi^-)\pi^0$

$\Gamma(K^*(892)^0 \bar{K}_2^*(1430)^0 + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{69}/\Gamma$

<u>VALUE (units <math>10^{-6}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>3.02 \pm 0.68 \pm 0.34</math></b>	42	SHEN	12A	BELL $\Upsilon(1S) \rightarrow K^+ K^- \pi^+ \pi^-$

$\Gamma(K_1(1270)^\pm K^\mp)/\Gamma_{\text{total}}$   $\Gamma_{70}/\Gamma$

<u>VALUE (units <math>10^{-6}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>&lt; 2.41</math></b>	90	SHEN	12A	BELL $\Upsilon(1S) \rightarrow K^+ K^- \pi^+ \pi^-$

$\Gamma(K_1(1400)^\pm K^\mp)/\Gamma_{\text{total}}$   $\Gamma_{71}/\Gamma$

<u>VALUE (units <math>10^{-6}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>1.02 \pm 0.35 \pm 0.22</math></b>	24	SHEN	12A	BELL $\Upsilon(1S) \rightarrow K^+ K^- \pi^+ \pi^-$

$\Gamma(b_1(1235)^\pm \pi^\mp)/\Gamma_{\text{total}}$   $\Gamma_{72}/\Gamma$

<u>VALUE (units <math>10^{-6}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>&lt; 1.25</math></b>	90	SHEN	12A	BELL $\Upsilon(1S) \rightarrow 2(\pi^+ \pi^-) \pi^0$

$\Gamma(\pi^+ \pi^- \pi^0 \pi^0)/\Gamma_{\text{total}}$   $\Gamma_{73}/\Gamma$

<u>VALUE (units <math>10^{-6}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>12.8 \pm 2.0 \pm 2.3</math></b>	$143 \pm 22$	SHEN	13	BELL $\Upsilon(1S) \rightarrow \pi^+ \pi^- \pi^0 \pi^0$

$\Gamma(K_S^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{74}/\Gamma$

<u>VALUE (units <math>10^{-6}</math>)</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>1.59 \pm 0.33 \pm 0.18</math></b>	$37 \pm 8$	SHEN	13	BELL	$\Upsilon(1S) \rightarrow K_S^0 K^- \pi^+$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 3.4$	90	<sup>1</sup> DOBBS	12A		$\Upsilon(1S) \rightarrow K_S^0 K^- \pi^+$
---------	----	--------------------	-----	--	--

<sup>1</sup> Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.

$\Gamma(K^*(892)^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{75}/\Gamma$

<u>VALUE (units <math>10^{-6}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>2.92 \pm 0.85 \pm 0.37</math></b>	$16 \pm 5$	SHEN	13	BELL $\Upsilon(1S) \rightarrow K_S^0 K^- \pi^+$

$\Gamma(K^*(892)^- K^+ + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{76}/\Gamma$

<u>VALUE (units <math>10^{-6}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>&lt; 1.11</math></b>	90	SHEN	13	BELL $\Upsilon(1S) \rightarrow K_S^0 K^- \pi^+$

$\Gamma(f_1(1285) \text{ anything})/\Gamma_{\text{total}}$   $\Gamma_{77}/\Gamma$

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>4.6 \pm 2.8 \pm 1.3</math></b>	3.1k	JIA	17A	BELL $e^+ e^- \rightarrow \text{hadrons}$

$\Gamma(D^*(2010)^\pm \text{ anything})/\Gamma_{\text{total}}$   $\Gamma_{78}/\Gamma$

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>25.2 \pm 1.3 \pm 1.5</math></b>	$\approx 2k$	<sup>1</sup> AUBERT	10c	BABR	$\Upsilon(2S) \rightarrow \pi^+ \pi^- \Upsilon(1S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 19$	90	<sup>2</sup> ALBRECHT	92J	ARG	$e^+ e^- \rightarrow D^0 \pi^\pm X$
--------	----	-----------------------	-----	-----	-------------------------------------

<sup>1</sup> For  $x_p > 0.1$ .

<sup>2</sup> For  $x_p > 0.2$ .

$\Gamma(f_1(1285)X_{tetra})/\Gamma_{total}$					$\Gamma_{79}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<b>&lt;62.4 × 10<sup>-6</sup></b>	90	<sup>1</sup> JIA	17A	BELL	$e^+e^- \rightarrow \text{hadrons}$

<sup>1</sup> For a tetraquark state  $X_{tetra}$ , with mass in the range 1.16–2.46 GeV and width in the range 0–0.3 GeV. Measured 90% CL limits as a function of  $X_{tetra}$  mass and width range from  $4.6 \times 10^{-6}$  to  $62.4 \times 10^{-6}$ .

$\Gamma(\overline{2H} \text{ anything})/\Gamma_{total}$					$\Gamma_{80}/\Gamma$
VALUE (units 10 <sup>-5</sup> )	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>2.85 ± 0.25 OUR AVERAGE</b>					

2.81 ± 0.49 <sup>+0.20</sup> <sub>-0.24</sub>		LEES	14G	BABR	$e^+e^- \rightarrow \overline{2H} X$
---	--	------	-----	------	--------------------------------------

2.86 ± 0.19 ± 0.21	455	ASNER	07	CLEO	$e^+e^- \rightarrow \overline{2H} X$
--------------------	-----	-------	----	------	--------------------------------------

$\Gamma(\text{Sum of 100 exclusive modes})/\Gamma_{total}$					$\Gamma_{81}/\Gamma$
VALUE (units 10 <sup>-2</sup> )		DOCUMENT ID		COMMENT	
<b>1.200 ± 0.017</b>		<sup>1,2</sup> DOBBS	12A	$\Upsilon(1S) \rightarrow \text{hadrons}$	

<sup>1</sup> DOBBS 12A presents individual exclusive branching fractions or upper limits for 100 modes of four to ten pions, kaons, or protons.

<sup>2</sup> Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.

$\Gamma(ggg, \gamma gg \rightarrow \overline{d} \text{ anything})/\Gamma(ggg, \gamma gg \rightarrow \text{anything})$					
VALUE (units 10 <sup>-5</sup> )	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>3.36 ± 0.23 ± 0.25</b>	455	ASNER	07	CLEO	$e^+e^- \rightarrow \overline{d} X$

$\Gamma(\gamma\pi^+\pi^-)/\Gamma_{total}$					$\Gamma_{82}/\Gamma$
VALUE (units 10 <sup>-5</sup> )		DOCUMENT ID	TECN	COMMENT	
<b>6.3 ± 1.2 ± 1.3</b>		<sup>1</sup> ANASTASSOV 99	CLE2	$e^+e^- \rightarrow \text{hadrons}$	

<sup>1</sup> For  $m_{\pi\pi} > 1$  GeV.

$\Gamma(\gamma\pi^0\pi^0)/\Gamma_{total}$					$\Gamma_{83}/\Gamma$
VALUE (units 10 <sup>-5</sup> )		DOCUMENT ID	TECN	COMMENT	
<b>1.7 ± 0.6 ± 0.3</b>		<sup>1</sup> ANASTASSOV 99	CLE2	$e^+e^- \rightarrow \text{hadrons}$	

<sup>1</sup> For  $m_{\pi\pi} > 1$  GeV.

$\Gamma(\gamma\pi\pi(\text{S-wave}))/\Gamma_{total}$					$\Gamma_{84}/\Gamma$
VALUE (units 10 <sup>-5</sup> )		DOCUMENT ID	TECN	COMMENT	
<b>4.63 ± 0.56 ± 0.48</b>		LEES	18A	BABR	$\Upsilon(1S) \rightarrow \gamma\pi^+\pi^-$

$\Gamma(\gamma\pi^0\eta)/\Gamma_{total}$					$\Gamma_{85}/\Gamma$
VALUE (units 10 <sup>-6</sup> )	CL%	DOCUMENT ID	TECN	COMMENT	
<b>&lt;2.4</b>	90	<sup>1</sup> BESSON	07A	CLEO	$e^+e^- \rightarrow \Upsilon(1S)$

<sup>1</sup> BESSON 07A obtained this limit for  $0.7 < m_{\pi^0\eta} < 3$  GeV.

$\Gamma(\gamma K^+ K^-)/\Gamma_{total}$ ( $2 < m_{K^+ K^-} < 3$ GeV)					$\Gamma_{86}/\Gamma$
VALUE (units 10 <sup>-5</sup> )	CL%	DOCUMENT ID	TECN	COMMENT	
<b>1.14 ± 0.08 ± 0.10</b>	90	ATHAR	06	CLE3	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$

$\Gamma(\gamma\rho\bar{\rho})/\Gamma_{\text{total}}$   $\Gamma_{87}/\Gamma$   
 ( $2 < m_{\rho\bar{\rho}} < 3 \text{ GeV}$ )

<u>VALUE (units <math>10^{-5}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt;0.6</b>	90	ATHAR	06 CLE3	$\Upsilon(1S) \rightarrow \gamma\rho\bar{\rho}$

$\Gamma(\gamma 2h^+ 2h^-)/\Gamma_{\text{total}}$   $\Gamma_{88}/\Gamma$

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>7.0 \pm 1.1 \pm 1.0</math></b>	$80 \pm 12$	FULTON	90B CLEO	$e^+e^- \rightarrow \text{hadrons}$

$\Gamma(\gamma 3h^+ 3h^-)/\Gamma_{\text{total}}$   $\Gamma_{89}/\Gamma$

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>5.4 \pm 1.5 \pm 1.3</math></b>	$39 \pm 11$	FULTON	90B CLEO	$e^+e^- \rightarrow \text{hadrons}$

$\Gamma(\gamma 4h^+ 4h^-)/\Gamma_{\text{total}}$   $\Gamma_{90}/\Gamma$

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>7.4 \pm 2.5 \pm 2.5</math></b>	$36 \pm 12$	FULTON	90B CLEO	$e^+e^- \rightarrow \text{hadrons}$

$\Gamma(\gamma\pi^+\pi^-K^+K^-)/\Gamma_{\text{total}}$   $\Gamma_{91}/\Gamma$

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>2.9 \pm 0.7 \pm 0.6</math></b>	$29 \pm 8$	FULTON	90B CLEO	$e^+e^- \rightarrow \text{hadrons}$

$\Gamma(\gamma 2\pi^+ 2\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{92}/\Gamma$

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>2.5 \pm 0.7 \pm 0.5</math></b>	$26 \pm 7$	FULTON	90B CLEO	$e^+e^- \rightarrow \text{hadrons}$

$\Gamma(\gamma 3\pi^+ 3\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{93}/\Gamma$

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>2.5 \pm 0.9 \pm 0.8</math></b>	$17 \pm 5$	FULTON	90B CLEO	$e^+e^- \rightarrow \text{hadrons}$

$\Gamma(\gamma 2\pi^+ 2\pi^- K^+ K^-)/\Gamma_{\text{total}}$   $\Gamma_{94}/\Gamma$

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>2.4 \pm 0.9 \pm 0.8</math></b>	$18 \pm 7$	FULTON	90B CLEO	$e^+e^- \rightarrow \text{hadrons}$

$\Gamma(\gamma\pi^+\pi^-\rho\bar{\rho})/\Gamma_{\text{total}}$   $\Gamma_{95}/\Gamma$

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>1.5 \pm 0.5 \pm 0.3</math></b>	$22 \pm 6$	FULTON	90B CLEO	$e^+e^- \rightarrow \text{hadrons}$

$\Gamma(\gamma 2\pi^+ 2\pi^- \rho\bar{\rho})/\Gamma_{\text{total}}$   $\Gamma_{96}/\Gamma$

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>0.4 \pm 0.4 \pm 0.4</math></b>	$7 \pm 6$	FULTON	90B CLEO	$e^+e^- \rightarrow \text{hadrons}$

$\Gamma(\gamma 2K^+ 2K^-)/\Gamma_{\text{total}}$   $\Gamma_{97}/\Gamma$

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>0.2 \pm 0.2</math></b>	$2 \pm 2$	FULTON	90B CLEO	$e^+e^- \rightarrow \text{hadrons}$

**$\Gamma(\gamma\eta'(958))/\Gamma_{\text{total}}$**   **$\Gamma_{98}/\Gamma$**

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt; 1.9</b>	90	ATHAR	07A CLEO	$\Upsilon(1S) \rightarrow \gamma\eta' \rightarrow \gamma\pi^+\pi^-\eta, \gamma\rho$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<16	90	RICHICHI	01B CLE2	$\Upsilon(1S) \rightarrow \gamma\eta' \rightarrow \gamma\eta\pi^+\pi^-$

**$\Gamma(\gamma\eta)/\Gamma_{\text{total}}$**   **$\Gamma_{99}/\Gamma$**

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt; 1.0</b>	90	ATHAR	07A CLEO	$\Upsilon(1S) \rightarrow \gamma\eta \rightarrow \gamma\gamma\gamma, \gamma\pi^+\pi^-\pi^0, \gamma3\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<21	90	MASEK	02 CLEO	$\Upsilon(1S) \rightarrow \gamma\eta$

**$\Gamma(\gamma f_0(980))/\Gamma_{\text{total}}$**   **$\Gamma_{100}/\Gamma$**

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;3</b>	90	<sup>1</sup> ATHAR	06 CLE3	$\Upsilon(1S) \rightarrow \gamma\pi^+\pi^-$
<sup>1</sup> Assuming $B(f_0(980) \rightarrow \pi\pi) = 1$ .				

**$\Gamma(\gamma f'_2(1525))/\Gamma_{\text{total}}$**   **$\Gamma_{101}/\Gamma$**

VALUE (units $10^{-5}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.9 ± 0.6 OUR AVERAGE</b>					
2.13 ± 0.28 ± 0.72			<sup>1</sup> LEES	18A BABR	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$
4.0 ± 1.4 ± 0.1		17	<sup>2</sup> BESSION	11 CLEO	$\Upsilon(1S) \rightarrow K_S^0 K_S^0$
3.7 <sup>+0.9</sup> / <sub>-0.7</sub> ± 0.8			ATHAR	06 CLE3	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$

- • • We do not use the following data for averages, fits, limits, etc. • • •
- <14                      90                      <sup>3</sup>FULTON                      90B CLEO                       $\Upsilon(1S) \rightarrow \gamma K^+ K^-$
- <19.4                      90                      <sup>3</sup>ALBRECHT                      89 ARG                       $\Upsilon(1S) \rightarrow \gamma K^+ K^-$

<sup>1</sup> Using  $B(f'_2(1525) \rightarrow K\bar{K}) = 0.887 \pm 0.022$  and  $B(K^0\bar{K}^0) = 1/2 B(K\bar{K})$ .  
<sup>2</sup> BESSION 11 reports  $(4.0 \pm 1.3 \pm 0.6) \times 10^{-5}$  from a measurement of  $[\Gamma(\Upsilon(1S) \rightarrow \gamma f'_2(1525))/\Gamma_{\text{total}}] \times [B(f'_2(1525) \rightarrow K\bar{K})]$  assuming  $B(f'_2(1525) \rightarrow K\bar{K}) = (88.8 \pm 3.1) \times 10^{-2}$ , which we rescale to our best (shown rounded) value  $B(f'_2(1525) \rightarrow K\bar{K}) = (88.8 \pm 2.2) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best (shown rounded) value. The result also assumes  $B(K_S^0 \rightarrow \pi^+\pi^-) = (69.20 \pm 0.05)\%$  and  $B(f'_2(1525) \rightarrow K\bar{K}) = 4 B(f'_2(1525) \rightarrow K_S^0 K_S^0)$ .  
<sup>3</sup> Assuming  $B(f'_2(1525) \rightarrow K\bar{K}) = 0.71$ .

**$\Gamma(\gamma f_2(1270))/\Gamma_{\text{total}}$**   **$\Gamma_{102}/\Gamma$**

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>10.1 ± 0.6 OUR AVERAGE</b>				
10.15 ± 0.59 <sup>+0.54</sup> / <sub>-0.43</sub>		<sup>1</sup> LEES	18A BABR	$\Upsilon(1S) \rightarrow \gamma\pi^+\pi^-$
10.5 ± 1.6 <sup>+1.9</sup> / <sub>-1.8</sub>		<sup>2</sup> BESSION	07A CLE3	$\Upsilon(1S) \rightarrow \gamma\pi^0\pi^0$
10.2 ± 0.8 ± 0.7		ATHAR	06 CLE3	$\Upsilon(1S) \rightarrow \gamma\pi^+\pi^-$
8.1 ± 2.3 <sup>+2.9</sup> / <sub>-2.7</sub>		<sup>3</sup> ANASTASSOV 99	CLE2	$e^+e^- \rightarrow \text{hadrons}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<21	90	<sup>3</sup> FULTON	90B	CLEO	$\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$
<13	90	<sup>3</sup> ALBRECHT	89	ARG	$\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$
<81	90	SCHMITT	88	CBAL	$\Upsilon(1S) \rightarrow \gamma X$
<sup>1</sup> Using $B(f_2(1270) \rightarrow \pi^0 \pi^0) = 1/3 B(f_2(1270) \rightarrow \pi \pi)$ and $B(f_2(1270) \rightarrow \pi \pi) = (84.2^{+2.9}_{-0.9})\%$ .					
<sup>2</sup> Using $B(f_2(1270) \rightarrow \pi^0 \pi^0) = B(f_2(1270) \rightarrow \pi \pi)/3$ and $B(f_2(1270) \rightarrow \pi \pi) = (84.7^{+2.5}_{-1.2})\%$ .					
<sup>3</sup> Using $B(f_2(1270) \rightarrow \pi \pi) = 0.84$ .					

$\Gamma(\gamma \eta(1405))/\Gamma_{\text{total}}$   $\Gamma_{103}/\Gamma$

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;8.2</b>	90	<sup>1</sup> FULTON	90B	CLEO $\Upsilon(1S) \rightarrow \gamma K^\pm \pi^\mp K_S^0$
<sup>1</sup> Includes unknown branching ratio of $\eta(1405) \rightarrow K^\pm \pi^\mp K_S^0$ .				

$\Gamma(\gamma f_0(1500))/\Gamma_{\text{total}}$   $\Gamma_{104}/\Gamma$

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;1.5</b>	90	<sup>1</sup> BESSON	07A	CLEO $e^+ e^- \rightarrow \Upsilon(1S) \rightarrow \gamma \pi^0 \pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<6.1	90	<sup>2</sup> BESSON	07A	CLEO $e^+ e^- \rightarrow \Upsilon(1S) \rightarrow \gamma \eta \eta$
<sup>1</sup> Using $B(f_0(1500) \rightarrow \pi^0 \pi^0) = B(f_0(1500) \rightarrow \pi \pi)/3$ and $B(f_0(1500) \rightarrow \pi \pi) = (0.349 \pm 0.023)\%$ .				
<sup>2</sup> Calculated by us using $B(f_0(1500) \rightarrow \eta \eta) = (5.1 \pm 0.9)\%$ .				

$\Gamma(\gamma f_0(1500) \rightarrow \gamma K^+ K^-)/\Gamma_{\text{total}}$   $\Gamma_{105}/\Gamma$

VALUE (units $10^{-5}$ )	DOCUMENT ID	TECN	COMMENT
<b><math>1.04 \pm 0.14 \pm 0.33</math></b>	<sup>1</sup> LEES	18A	BABR $e^+ e^- \rightarrow \Upsilon(1S) \rightarrow \gamma K^+ K^-$
<sup>1</sup> LEES 18A quotes $B(\Upsilon(1S) \rightarrow \gamma f_0(1500) \rightarrow \gamma K \bar{K}) = (2.08 \pm 0.27 \pm 0.65) \times 10^{-5}$ assuming $B(K^0 \bar{K}^0) = 1/2 B(K \bar{K})$ .			

$\Gamma(\gamma f_0(1710))/\Gamma_{\text{total}}$   $\Gamma_{106}/\Gamma$

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt; 2.6</b>	90	<sup>1</sup> ALBRECHT	89	ARG $\Upsilon(1S) \rightarrow \gamma K^+ K^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 6.3	90	<sup>1</sup> FULTON	90B	CLEO $\Upsilon(1S) \rightarrow \gamma K^+ K^-$
<19	90	<sup>1</sup> FULTON	90B	CLEO $\Upsilon(1S) \rightarrow \gamma K_S^0 K_S^0$
< 8	90	<sup>2</sup> ALBRECHT	89	ARG $\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$
<24	90	<sup>3</sup> SCHMITT	88	CBAL $\Upsilon(1S) \rightarrow \gamma X$
<sup>1</sup> Assuming $B(f_0(1710) \rightarrow K \bar{K}) = 0.38$ .				
<sup>2</sup> Assuming $B(f_0(1710) \rightarrow \pi \pi) = 0.04$ .				
<sup>3</sup> Assuming $B(f_0(1710) \rightarrow \eta \eta) = 0.18$ .				

$\Gamma(\gamma f_0(1710) \rightarrow \gamma K^+ K^-)/\Gamma_{\text{total}}$   $\Gamma_{107}/\Gamma$

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b><math>1.01 \pm 0.26 \pm 0.18</math></b>		<sup>1</sup> LEES	18A	BABR $e^+ e^- \rightarrow \Upsilon(1S) \rightarrow \gamma K^+ K^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.7	90	ATHAR	06	CLEO $e^+ e^- \rightarrow \Upsilon(1S) \rightarrow \gamma K^+ K^-$

<sup>1</sup> LEES 18A quotes  $B(\Upsilon(1S) \rightarrow \gamma f_0(1710) \rightarrow \gamma K \bar{K}) = (2.02 \pm 0.51 \pm 0.35) \times 10^{-5}$   
assuming  $B(K^0 \bar{K}^0) = 1/2 B(K \bar{K})$ .

### $\Gamma(\gamma f_0(1710) \rightarrow \gamma \pi^+ \pi^-) / \Gamma_{\text{total}}$ $\Gamma_{108} / \Gamma$

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b><math>0.53 \pm 0.17 \pm 0.11</math></b>		<sup>1</sup> LEES	18A	BABR $\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$

<sup>1</sup> LEES 18A quotes  $B(\Upsilon(1S) \rightarrow \gamma f_0(1710) \rightarrow \gamma \pi \pi) = (0.79 \pm 0.26 \pm 0.17) \times 10^{-5}$   
assuming  $B(\pi^0 \pi^0) = 1/3 B(\pi \pi)$ .

### $\Gamma(\gamma f_0(1710) \rightarrow \gamma \pi^0 \pi^0) / \Gamma_{\text{total}}$ $\Gamma_{109} / \Gamma$

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt; 1.4</b>	90	BESSON	07A	CLEO $e^+ e^- \rightarrow \Upsilon(1S) \rightarrow \gamma \pi^0 \pi^0$

### $\Gamma(\gamma f_0(1710) \rightarrow \gamma \eta \eta) / \Gamma_{\text{total}}$ $\Gamma_{110} / \Gamma$

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt; 1.8</b>	90	BESSON	07A	CLEO $e^+ e^- \rightarrow \Upsilon(1S) \rightarrow \gamma \eta \eta$

### $\Gamma(\gamma f_4(2050)) / \Gamma_{\text{total}}$ $\Gamma_{111} / \Gamma$

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt; 5.3</b>	90	<sup>1</sup> ATHAR	06	CLE3 $\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$

<sup>1</sup> Assuming  $B(f_4(2050) \rightarrow \pi \pi) = 0.17$ .

### $\Gamma(\gamma f_0(2200) \rightarrow \gamma K^+ K^-) / \Gamma_{\text{total}}$ $\Gamma_{112} / \Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt; 0.0002</b>	90	BARU	89	MD1 $\Upsilon(1S) \rightarrow \gamma K^+ K^-$

### $\Gamma(\gamma f_J(2220) \rightarrow \gamma K^+ K^-) / \Gamma_{\text{total}}$ $\Gamma_{113} / \Gamma$

VALUE (units $10^{-7}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt; 8</b>	90	ATHAR	06	CLE3 $\Upsilon(1S) \rightarrow \gamma K^+ K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 160	90	MASEK	02	CLEO $\Upsilon(1S) \rightarrow \gamma K^+ K^-$
< 150	90	FULTON	90B	CLEO $\Upsilon(1S) \rightarrow \gamma K^+ K^-$
< 290	90	ALBRECHT	89	ARG $\Upsilon(1S) \rightarrow \gamma K^+ K^-$
< 2000	90	BARU	89	MD1 $\Upsilon(1S) \rightarrow \gamma K^+ K^-$

### $\Gamma(\gamma f_J(2220) \rightarrow \gamma \pi^+ \pi^-) / \Gamma_{\text{total}}$ $\Gamma_{114} / \Gamma$

VALUE (units $10^{-7}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt; 6</b>	90	ATHAR	06	CLE3 $\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 120	90	MASEK	02	CLEO $\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$
-------	----	-------	----	--

### $\Gamma(\gamma f_J(2220) \rightarrow \gamma p \bar{p}) / \Gamma_{\text{total}}$ $\Gamma_{115} / \Gamma$

VALUE (units $10^{-7}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt; 11</b>	90	ATHAR	06	CLE3 $\Upsilon(1S) \rightarrow \gamma p \bar{p}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 160	90	MASEK	02	CLEO $\Upsilon(1S) \rightarrow \gamma p \bar{p}$
-------	----	-------	----	--

$\Gamma(\gamma\eta(2225) \rightarrow \gamma\phi\phi)/\Gamma_{\text{total}}$   $\Gamma_{116}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.003	90	BARU	89 MD1	$\Upsilon(1S) \rightarrow \gamma K^+ K^- K^+ K^-$

 $\Gamma(\gamma\eta_c(1S))/\Gamma_{\text{total}}$   $\Gamma_{117}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<2.9 × 10 <sup>-5</sup>	90	<sup>1</sup> KATRENKO	20 BELL	$e^+e^- \rightarrow \gamma + \text{hadrons}$
<5.7 × 10 <sup>-5</sup>	90	SHEN	10A BELL	$\Upsilon(1S) \rightarrow \gamma X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<sup>1</sup> Using  $\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-$  decays.

 $\Gamma(\gamma\eta_c(2S))/\Gamma_{\text{total}}$   $\Gamma_{118}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<4 × 10 <sup>-4</sup>	90	<sup>1</sup> KATRENKO	20 BELL	$e^+e^- \rightarrow \gamma + \text{hadrons}$

<sup>1</sup> Using  $\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-$  decays.

 $\Gamma(\gamma\chi_{c0})/\Gamma_{\text{total}}$   $\Gamma_{119}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<6.6 × 10 <sup>-5</sup>	90	<sup>1</sup> KATRENKO	20 BELL	$\Upsilon(1S) \rightarrow \gamma + \text{hadrons}$
<6.5 × 10 <sup>-4</sup>	90	SHEN	10A BELL	$\Upsilon(1S) \rightarrow \gamma X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<sup>1</sup> Using  $\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-$  decays.

 $\Gamma(\gamma\chi_{c1})/\Gamma_{\text{total}}$   $\Gamma_{120}/\Gamma$ 

VALUE (units 10 <sup>-5</sup> )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$4.7^{+2.4+0.4}_{-1.8-0.5}$		5	<sup>1</sup> KATRENKO	20 BELL	$\Upsilon(1S) \rightarrow \gamma + \text{hadrons}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<2.3      90      SHEN      10A BELL       $\Upsilon(1S) \rightarrow \gamma X$

<sup>1</sup> Using  $\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-$  decays.

 $\Gamma(\gamma\chi_{c2})/\Gamma_{\text{total}}$   $\Gamma_{121}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<7.6 × 10 <sup>-6</sup>	90	SHEN	10A BELL	$\Upsilon(1S) \rightarrow \gamma X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<3.3 × 10<sup>-5</sup>      90      <sup>1</sup> KATRENKO      20 BELL       $\Upsilon(1S) \rightarrow \gamma + \text{hadrons}$

<sup>1</sup> Using  $\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-$  decays.

 $\Gamma(\gamma\chi_{c1}(3872))/\Gamma_{\text{total}}$   $\Gamma_{122}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<4 × 10 <sup>-5</sup>	90	<sup>1</sup> SHEN	10A BELL	$\Upsilon(1S) \rightarrow \gamma X$

<sup>1</sup> SHEN 10A reports  $[\Gamma(\Upsilon(1S) \rightarrow \gamma\chi_{c1}(3872))/\Gamma_{\text{total}}] \times [B(\chi_{c1}(3872) \rightarrow \pi^+\pi^- J/\psi(1S))] < 1.6 \times 10^{-6}$  which we divide by our best (shown rounded) value  $B(\chi_{c1}(3872) \rightarrow \pi^+\pi^- J/\psi(1S)) = 4.3 \times 10^{-2}$ .

$\Gamma(\gamma\chi_{c1}(3872), \chi_{c1} \rightarrow \pi^+\pi^-\pi^0 J/\psi)/\Gamma_{\text{total}}$   $\Gamma_{123}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.8 \times 10^{-6}$	90	SHEN	10A	BELL $\Upsilon(1S) \rightarrow \gamma X$

$\Gamma(\gamma\chi_{c0}(3915) \rightarrow \omega J/\psi)/\Gamma_{\text{total}}$   $\Gamma_{124}/\Gamma$

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$<3.0$	90	SHEN	10A	BELL $\Upsilon(1S) \rightarrow \gamma X$

$\Gamma(\gamma\chi_{c1}(4140) \rightarrow \phi J/\psi)/\Gamma_{\text{total}}$   $\Gamma_{125}/\Gamma$

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$<2.2$	90	SHEN	10A	BELL $\Upsilon(1S) \rightarrow \gamma X$

$\Gamma(\gamma X \bar{X} (m_X < 3.1 \text{ GeV}))/\Gamma_{\text{total}}$   $\Gamma_{126}/\Gamma$

( $X \bar{X}$  = vectors with  $m < 3.1$  GeV)

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$<1$	90	<sup>1</sup> BALEST	95	CLEO $e^+e^- \rightarrow \gamma + X \bar{X}$

<sup>1</sup>For a noninteracting vector  $X$  with mass  $< 3.1$  GeV.

$\Gamma(\gamma X \bar{X} (m_X < 4.5 \text{ GeV}))/\Gamma_{\text{total}}$   $\Gamma_{127}/\Gamma$

$X$  and  $\bar{X}$  = zero spin with  $m < 4.5$  GeV

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$<24$	90	<sup>1</sup> DEL-AMO-SA..11J	BABR	$e^+e^- \rightarrow \gamma + X \bar{X}$

<sup>1</sup>For a noninteracting scalar  $X$  with mass  $m < 4.5$  GeV.

$\Gamma(\gamma X \rightarrow \gamma + \geq 4 \text{ prongs})/\Gamma_{\text{total}}$   $\Gamma_{128}/\Gamma$

( $1.5 \text{ GeV} < m_X < 5.0 \text{ GeV}$ )

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$<1.78$	95	ROSNER	07A	CLEO $e^+e^- \rightarrow \gamma X$

$\Gamma(\gamma A^0)/\Gamma_{\text{total}}$   $\Gamma_{129}/\Gamma$

( $A^0$  = scalar with  $m < 8.0$  GeV)

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

$<4.5 \times 10^{-6}$	90	<sup>1</sup> DEL-AMO-SA..11J	BABR	$e^+e^- \rightarrow \gamma + X$
$<3 \times 10^{-5}$	90	<sup>2</sup> BALEST	95	CLEO $e^+e^- \rightarrow \gamma + X$
$<5.6 \times 10^{-5}$	90	<sup>2</sup> ANTREASYAN 90C	CBAL	$e^+e^- \rightarrow \gamma + X$

<sup>1</sup>For a non-interacting scalar or pseudoscalar,  $A^0$ , with mass  $m_{A^0} < 8.0$  GeV. 90% CL upper limits range from  $1.9 \times 10^{-6}$  to  $4.5 \times 10^{-6}$ .

<sup>2</sup>For any non-interacting long-lived particle with mass  $< 7.2$  GeV.

$\Gamma(\gamma A^0 \rightarrow \gamma \mu^+ \mu^-)/\Gamma_{\text{total}}$   $\Gamma_{130}/\Gamma$

( $201 < M(\mu^+ \mu^-) < 3565 \text{ MeV}$ )

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$<9$	90	<sup>1</sup> LOVE	08	CLEO $e^+e^- \rightarrow \gamma A^0 \rightarrow \gamma \mu^+ \mu^-$

••• We do not use the following data for averages, fits, limits, etc. •••

$<16$	90	<sup>2</sup> JIA	22	BELL $\Upsilon(2S) \rightarrow \gamma \mu^+ \mu^- \pi^+ \pi^-$
$<9.7$	90	<sup>3</sup> LEES	13C	BABR $e^+e^- \rightarrow \gamma A^0 \rightarrow \gamma \mu^+ \mu^-$

- <sup>1</sup> For a narrow scalar or pseudoscalar,  $A^0$ , with  $201 < M(\mu^+ \mu^-) < 3565$  MeV, excluding  $J/\psi$ . Measured 90% CL limits as a function of  $M(\mu^+ \mu^-)$  range from  $1-9 \times 10^{-6}$ .
- <sup>2</sup> For a narrow scalar or pseudoscalar,  $A^0$ , with  $0.22 < M(A^0) < 9.2$  GeV, resulting in 90% CL upper limits ranging from  $3.1 \times 10^{-7}$  at  $M(A^0) = 0.22$  GeV to  $1.6 \times 10^{-5}$  at  $M(A^0) = 9.2$  GeV.
- <sup>3</sup> For a narrow scalar or pseudoscalar,  $A^0$ , with mass in the range 0.212–9.2 GeV, excluding  $J/\psi$  and  $\psi(2S)$ . Measured 90% CL limits as a function of  $m_{A^0}$  are in the range  $0.28-9.7 \times 10^{-6}$ .

**$\Gamma(\gamma A^0 \rightarrow \gamma \tau^+ \tau^-) / \Gamma_{\text{total}}$**   **$\Gamma_{131} / \Gamma$**   
 ( $2m_\tau < M(\tau^+ \tau^-) < 9.2$  GeV)

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;130</b>	90	<sup>1</sup> LEES	13R	BABR $\Upsilon(2S) \rightarrow \gamma \tau^+ \tau^- \pi^+ \pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<150	90	<sup>2</sup> JIA	22	BELL $\Upsilon(2S) \rightarrow \gamma \tau^+ \tau^- \pi^+ \pi^-$
< 50	90	<sup>3</sup> LOVE	08	CLEO $e^+ e^- \rightarrow \gamma A^0 \rightarrow \gamma \tau^+ \tau^-$

- <sup>1</sup> For a narrow scalar or pseudoscalar,  $A^0$ , with  $2m_\tau < M(A^0) < 9.2$  GeV, resulting in 90% CL upper limits of  $0.9 \times 10^{-5}$  at  $M(A^0) = 2m_\tau$ ,  $\approx 1.5 \times 10^{-5}$  at  $M(A^0) = 7.5$  GeV, and  $13 \times 10^{-5}$  at  $M(A^0) = 9.2$  GeV.
- <sup>2</sup> For a narrow scalar or pseudoscalar,  $A^0$ , with  $2m_\tau < M(A^0) < 9.2$  GeV, resulting in 90% CL upper limits ranging from  $3.8 \times 10^{-6}$  at  $M(A^0) = 2m_\tau$  to  $1.5 \times 10^{-4}$  at  $M(A^0) = 9.2$  GeV.
- <sup>3</sup> For a narrow scalar or pseudoscalar,  $A^0$ , with  $2m_\tau < M(A^0) < 7.5$  GeV, resulting in 90% CL limits ranging from  $1 \times 10^{-5}$  at  $M(A^0) = 2m_\tau$  to  $5 \times 10^{-5}$  at  $M(A^0) = 7.5$  GeV.

**$\Gamma(\gamma A^0 \rightarrow \gamma g g) / \Gamma_{\text{total}}$**   **$\Gamma_{132} / \Gamma$**   
 ( $0.5 \text{ GeV} < m < 9.0 \text{ GeV}$ )

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;1 x 10<sup>-2</sup></b>	90	<sup>1</sup> LEES	13L	BABR $\Upsilon(1S) \rightarrow \gamma X$

- <sup>1</sup> For a narrow,  $CP$ -odd pseudoscalar,  $A^0$ , searched for in 26 hadronic decay modes with invariant mass  $0.5 \text{ GeV} < m_{A^0} < 9.0 \text{ GeV}$ . Measured 90% CL limits as a function of  $m_{A^0}$  range from  $10^{-6}$  to  $10^{-2}$ .

**$\Gamma(\gamma A^0 \rightarrow \gamma s \bar{s}) / \Gamma_{\text{total}}$**   **$\Gamma_{133} / \Gamma$**   
 ( $0.5 \text{ GeV} < m < 9.0 \text{ GeV}$ )

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;1 x 10<sup>-3</sup></b>	90	<sup>1</sup> LEES	13L	BABR $\Upsilon(1S) \rightarrow \gamma X$

- <sup>1</sup> For a narrow,  $CP$ -odd pseudoscalar,  $A^0$ , searched for in 14 hadronic decay modes with invariant mass  $1.5 \text{ GeV} < m_{A^0} < 9.0 \text{ GeV}$ . Measured 90% CL limits as a function of  $m_{A^0}$  range from  $10^{-5}$  to  $10^{-3}$ .

————— **LEPTON FAMILY NUMBER (LF) VIOLATING MODES** —————

**$\Gamma(e^\pm \mu^\mp) / \Gamma_{\text{total}}$**   **$\Gamma_{134} / \Gamma$**

VALUE (units $10^{-7}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;3.9</b>	90	PATRA	22	BELL $\Upsilon(2S) \rightarrow \pi^+ \pi^- e^\pm \mu^\mp$

$\Gamma(\mu^\pm \tau^\mp)/\Gamma_{\text{total}}$   $\Gamma_{135}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.7 \times 10^{-6}$	90	PATRA	22	BELL $\Upsilon(2S) \rightarrow \pi^+ \pi^- \mu^\pm \tau^\mp$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$<6.0 \times 10^{-6}$	95	LOVE	08A	CLEO $e^+ e^- \rightarrow \mu^\pm \tau^\mp$

$\Gamma(e^\pm \tau^\mp)/\Gamma_{\text{total}}$   $\Gamma_{136}/\Gamma$

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$<2.7$	90	PATRA	22	BELL $\Upsilon(2S) \rightarrow \pi^+ \pi^- e^\pm \tau^\mp$

$\Gamma(\gamma e^\pm \mu^\mp)/\Gamma_{\text{total}}$   $\Gamma_{137}/\Gamma$

VALUE (units $10^{-7}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$<4.2$	90	PATRA	22	BELL $\Upsilon(2S) \rightarrow \pi^+ \pi^- \gamma e^\pm \mu^\mp$

$\Gamma(\gamma \mu^\pm \tau^\mp)/\Gamma_{\text{total}}$   $\Gamma_{138}/\Gamma$

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$<6.1$	90	PATRA	22	BELL $\Upsilon(2S) \rightarrow \pi^+ \pi^- \gamma \mu^\pm \tau^\mp$

$\Gamma(\gamma e^\pm \tau^\mp)/\Gamma_{\text{total}}$   $\Gamma_{139}/\Gamma$

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$<6.5$	90	PATRA	22	BELL $\Upsilon(2S) \rightarrow \pi^+ \pi^- \gamma e^\pm \tau^\mp$

————— OTHER DECAYS —————

$\Gamma(\text{invisible})/\Gamma_{\text{total}}$   $\Gamma_{140}/\Gamma$

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$< 3.0$	90	AUBERT	09AX	BABR $\Upsilon(3S) \rightarrow \pi^+ \pi^- \Upsilon(1S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$<39$	90	RUBIN	07	CLEO $\Upsilon(2S) \rightarrow \pi^+ \pi^- \Upsilon(1S)$
$<25$	90	TAJIMA	07	BELL $\Upsilon(3S) \rightarrow \pi^+ \pi^- \Upsilon(1S)$

**$\Upsilon(1S)$  REFERENCES**

ADACHI	25A	PRL 135 041901	I. Adachi <i>et al.</i>	(BELLE and BELLE II Collab.)
DONG	25	JHEP 2509 048	X. Dong <i>et al.</i>	(BELLE Collab.)
AAIJ	24AC	JHEP 2410 122	R. Aaij <i>et al.</i>	(LHCb Collab.)
SHAMOV	23	PL B839 137766	A.G. Shamov, O.L. Rezanova	(NOVO, NOVOU)
JIA	22	PRL 128 081804	S. Jia <i>et al.</i>	(BELLE Collab.)
PATRA	22	JHEP 2205 095	S. Patra <i>et al.</i>	(BELLE Collab.)
KATRENKO	20	PRL 124 122001	P. Katrenko <i>et al.</i>	(BELLE Collab.)
JIA	18	PR D97 112004	S. Jia <i>et al.</i>	(BELLE Collab.)
LEES	18A	PR D97 112006	J.P. Lees <i>et al.</i>	(BABAR Collab.)
JIA	17	PR D95 012001	S. Jia <i>et al.</i>	(BELLE Collab.)
JIA	17A	PR D96 112002	S. Jia <i>et al.</i>	(BELLE Collab.)
SHEN	16	PR D93 112013	C.P. Shen <i>et al.</i>	(BELLE Collab.)
LEES	14G	PR D89 111102	J.P. Lees <i>et al.</i>	(BABAR Collab.)
YANG	14	PR D90 112008	S.D. Yang <i>et al.</i>	(BELLE Collab.)
LEES	13C	PR D87 031102	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	13L	PR D88 031701	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	13R	PR D88 071102	J.P. Lees <i>et al.</i>	(BABAR Collab.)
SHEN	13	PR D88 011102	C.P. Shen <i>et al.</i>	(BELLE Collab.)
DOBBS	12A	PR D86 052003	S. Dobbs <i>et al.</i>	
SHEN	12A	PR D86 031102	C.P. Shen <i>et al.</i>	(BELLE Collab.)
BESSION	11	PR D83 037101	D. Besson <i>et al.</i>	(CLEO Collab.)

DEL-AMO-SA...	11J	PRL 107 021804	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
AUBERT	10C	PR D81 011102	B. Aubert <i>et al.</i>	(BABAR Collab.)
DEL-AMO-SA...	10C	PRL 104 191801	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
SHEN	10A	PR D82 051504	C.P. Shen <i>et al.</i>	(BELLE Collab.)
AUBERT	09AX	PRL 103 251801	B. Aubert <i>et al.</i>	(BABAR Collab.)
LOVE	08	PRL 101 151802	W. Love <i>et al.</i>	(CLEO Collab.)
LOVE	08A	PRL 101 201601	W. Love <i>et al.</i>	(CLEO Collab.)
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)
ASNER	07	PR D75 012009	D.M. Asner <i>et al.</i>	(CLEO Collab.)
ATHAR	07A	PR D76 072003	S.B. Athar <i>et al.</i>	(CLEO Collab.)
BESSON	07	PRL 98 052002	D. Besson <i>et al.</i>	(CLEO Collab.)
BESSON	07A	PR D75 072001	D. Besson <i>et al.</i>	(CLEO Collab.)
ROSNER	07A	PR D76 117102	J.L. Rosner <i>et al.</i>	(CLEO Collab.)
RUBIN	07	PR D75 031104	P. Rubin <i>et al.</i>	(CLEO Collab.)
TAJIMA	07	PRL 98 132001	O. Tajima <i>et al.</i>	(BELLE Collab.)
AQUINES	06A	PR D74 092006	O. Aquines <i>et al.</i>	(CLEO Collab.)
ATHAR	06	PR D73 032001	S.B. Athar <i>et al.</i>	(CLEO Collab.)
BESSON	06A	PR D74 012003	D. Besson <i>et al.</i>	(CLEO Collab.)
ROSNER	06	PRL 96 092003	J.L. Rosner <i>et al.</i>	(CLEO Collab.)
ADAMS	05	PRL 94 012001	G.S. Adams <i>et al.</i>	(CLEO Collab.)
BRIERE	04	PR D70 072001	R.A. Briere <i>et al.</i>	(CLEO Collab.)
ARTUSO	03	PR D67 052003	M. Artuso <i>et al.</i>	(CLEO Collab.)
MASEK	02	PR D65 072002	G. Masek <i>et al.</i>	(CLEO Collab.)
RICHICHI	01B	PRL 87 141801	S.J. Richichi <i>et al.</i>	(CLEO Collab.)
ARTAMONOV	00	PL B474 427	A.S. Artamonov <i>et al.</i>	
ANASTASSOV	99	PRL 82 286	A. Anastassov <i>et al.</i>	(CLEO Collab.)
ALEXANDER	98	PR D58 052004	J.P. Alexander <i>et al.</i>	(CLEO Collab.)
BARU	96	PRPL 267 71	S.E. Baru <i>et al.</i>	(NOVO)
BALEST	95	PR D51 2053	R. Balest <i>et al.</i>	(CLEO Collab.)
CINABRO	94B	PL B340 129	D. Cinabro <i>et al.</i>	(CLEO Collab.)
ALBRECHT	92J	ZPHY C55 25	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
BARU	92	ZPHY C54 229	S.E. Baru <i>et al.</i>	(NOVO)
BARU	92B	ZPHY C56 547	S.E. Baru <i>et al.</i>	(NOVO)
KOBEL	92	ZPHY C53 193	M. Kobel <i>et al.</i>	(Crystal Ball Collab.)
ANTREASYAN	90C	PL B251 204	D. Antreasyan <i>et al.</i>	(Crystal Ball Collab.)
BLINOV	90	PL B245 311	A.E. Blinov <i>et al.</i>	(NOVO)
FULTON	90B	PR D41 1401	R. Fulton <i>et al.</i>	(CLEO Collab.)
MASCHMANN	90	ZPHY C46 555	W.S. Maschmann <i>et al.</i>	(Crystal Ball Collab.)
ALBRECHT	89	ZPHY C42 349	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALEXANDER	89	NP B320 45	J.P. Alexander <i>et al.</i>	(LBL, MICH, SLAC)
BARU	89	ZPHY C42 505	S.E. Baru <i>et al.</i>	(NOVO)
CHEN	89B	PR D39 3528	W.Y. Chen <i>et al.</i>	(CLEO Collab.)
FULTON	89	PL B224 445	R. Fulton <i>et al.</i>	(CLEO Collab.)
KAARSBERG	89	PRL 62 2077	T.M. Kaarsberg <i>et al.</i>	(CUSB Collab.)
BUCHMUEL...	88	HE $e^+e^-$ Physics 412	W. Buchmueller, S. Cooper	(HANN, DESY, MIT)
Editors: A. Ali and P. Soeding, World Scientific, Singapore				
JAKUBOWSKI	88	ZPHY C40 49	Z. Jakubowski <i>et al.</i>	(Crystal Ball Collab.) IGJPC
SCHMITT	88	ZPHY C40 199	P. Schmitt <i>et al.</i>	(Crystal Ball Collab.)
ALBRECHT	87	ZPHY C35 283	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
COHEN	87	RMP 59 1121	E.R. Cohen, B.N. Taylor	(RISC, NBS)
BARU	86	ZPHY C30 551	S.E. Baru <i>et al.</i>	(NOVO)
ALBRECHT	85C	PL 154B 452	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
KURAEV	85	SJNP 41 466	E.A. Kuraev, V.S. Fadin	(NOVO)
Translated from YAF 41 733.				
ARTAMONOV	84	PL 137B 272	A.S. Artamonov <i>et al.</i>	(NOVO)
BESSON	84	PR D30 1433	D. Besson <i>et al.</i>	(CLEO Collab.)
GILES	84B	PR D29 1285	R. Giles <i>et al.</i>	(CLEO Collab.)
MACKAY	84	PR D29 2483	W.W. MacKay <i>et al.</i>	(CUSB Collab.)
ANDREWS	83	PRL 50 807	D.E. Andrews <i>et al.</i>	(CLEO Collab.)
GILES	83	PRL 50 877	R. Giles <i>et al.</i>	(HARV, OSU, ROCH, RUTG+)
NICZYPORUK	83	ZPHY C17 197	B. Niczyporuk <i>et al.</i>	(LENA Collab.)
ALBRECHT	82	PL 116B 383	H. Albrecht <i>et al.</i>	(DESY, DORT, HEIDH+)
ARTAMONOV	82	PL 118B 225	A.S. Artamonov <i>et al.</i>	(NOVO)
NICZYPORUK	82	ZPHY C15 299	B. Niczyporuk <i>et al.</i>	(LENA Collab.)
BERGER	80C	PL 93B 497	C. Berger <i>et al.</i>	(PLUTO Collab.)
BOCK	80	ZPHY C6 125	P. Bock <i>et al.</i>	(HEIDP, MPIM, DESY, HAMB)
BERGER	79	ZPHY C1 343	C. Berger <i>et al.</i>	(PLUTO Collab.)