

$\Upsilon_2(1D)$

$$I^G(J^{PC}) = 0^-(2^{--})$$

was $\Upsilon(1D)$

First observed by BONVICINI 04 in the decay to $\gamma\gamma \Upsilon(1S)$ and confirmed by DEL-AMO-SANCHEZ 10R in the decay to $\pi^+\pi^- \Upsilon(1S)$.

Data consistent with $J^P = 2^-$. The states with $J = 1$ and 3 also possibly seen, but need confirmation.

$\Upsilon_2(1D)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
10163.7 ± 1.4 OUR AVERAGE		Error includes scale factor of 1.7.		
$10164.5 \pm 0.8 \pm 0.5$		DEL-AMO-SA...10R	BABR	$\Upsilon(3S) \rightarrow \gamma\gamma\pi^+\pi^-\ell^+\ell^-$
$10161.1 \pm 0.6 \pm 1.6$	38	BONVICINI 04	CLE3	$\Upsilon(3S) \rightarrow 4\gamma\ell^+\ell^-$

$\Upsilon_2(1D)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \quad \gamma\gamma \Upsilon(1S)$	seen
$\Gamma_2 \quad \gamma\chi_{bJ}(1P)$	seen
$\Gamma_3 \quad \eta \Upsilon(1S)$	not seen
$\Gamma_4 \quad \pi^+\pi^- \Upsilon(1S)$	$(6.6 \pm 1.6) \times 10^{-3}$

$\Upsilon_2(1D)$ BRANCHING RATIOS

$\Gamma(\eta \Upsilon(1S))/\Gamma(\gamma\gamma \Upsilon(1S))$	Γ_3/Γ_1										
<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="text-align: left;">VALUE</th> <th style="text-align: left;">CL%</th> <th style="text-align: left;">DOCUMENT ID</th> <th style="text-align: left;">TECN</th> <th style="text-align: left;">COMMENT</th> </tr> </thead> <tbody> <tr> <td><0.25</td> <td>90</td> <td>BONVICINI 04</td> <td>CLE3</td> <td>$\Upsilon(3S) \rightarrow 4\gamma\ell^+\ell^-$</td> </tr> </tbody> </table>	VALUE	CL%	DOCUMENT ID	TECN	COMMENT	<0.25	90	BONVICINI 04	CLE3	$\Upsilon(3S) \rightarrow 4\gamma\ell^+\ell^-$	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT							
<0.25	90	BONVICINI 04	CLE3	$\Upsilon(3S) \rightarrow 4\gamma\ell^+\ell^-$							

$\Gamma(\pi^+\pi^- \Upsilon(1S))/\Gamma_{\text{total}}$	Γ_4/Γ								
<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="text-align: left;">VALUE (units 10^{-2})</th> <th style="text-align: left;">DOCUMENT ID</th> <th style="text-align: left;">TECN</th> <th style="text-align: left;">COMMENT</th> </tr> </thead> <tbody> <tr> <td>$0.66^{+0.15}_{-0.14} \pm 0.06$</td> <td>¹ DEL-AMO-SA...10R</td> <td>BABR</td> <td>$\Upsilon(3S) \rightarrow \gamma\gamma\pi^+\pi^-\ell^+\ell^-$</td> </tr> </tbody> </table>	VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT	$0.66^{+0.15}_{-0.14} \pm 0.06$	¹ DEL-AMO-SA...10R	BABR	$\Upsilon(3S) \rightarrow \gamma\gamma\pi^+\pi^-\ell^+\ell^-$	
VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT						
$0.66^{+0.15}_{-0.14} \pm 0.06$	¹ DEL-AMO-SA...10R	BABR	$\Upsilon(3S) \rightarrow \gamma\gamma\pi^+\pi^-\ell^+\ell^-$						

¹ Using theoretical predictions for $B(\chi_{bJ}(2P) \rightarrow \gamma \Upsilon_2(1D))$.

$\Gamma(\pi^+\pi^- \Upsilon(1S))/\Gamma(\gamma\gamma \Upsilon(1S))$	Γ_4/Γ_1										
<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="text-align: left;">VALUE</th> <th style="text-align: left;">CL%</th> <th style="text-align: left;">DOCUMENT ID</th> <th style="text-align: left;">TECN</th> <th style="text-align: left;">COMMENT</th> </tr> </thead> <tbody> <tr> <td><1.2</td> <td>90</td> <td>² BONVICINI 04</td> <td>CLE3</td> <td>$\Upsilon(3S) \rightarrow 4\gamma\ell^+\ell^-$</td> </tr> </tbody> </table>	VALUE	CL%	DOCUMENT ID	TECN	COMMENT	<1.2	90	² BONVICINI 04	CLE3	$\Upsilon(3S) \rightarrow 4\gamma\ell^+\ell^-$	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT							
<1.2	90	² BONVICINI 04	CLE3	$\Upsilon(3S) \rightarrow 4\gamma\ell^+\ell^-$							

² Assuming $J = 2$.

$\Upsilon_2(1D)$ REFERENCES

DEL-AMO-SA... 10R	PR D82 111102	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
BONVICINI 04	PR D70 032001	G. Bonvicini <i>et al.</i>	(CLEO Collab.)